

• **Problem 1**

(a)

$$30030 = 257 (116) + 218$$

$$257 = 218 (1) + 39$$

$$218 = 39 (5) + 23$$

$$39 = 23 (1) + 16$$

$$23 = 16 (1) + 7$$

$$16 = 7 (2) + 2$$

$$7 = 2 (3) + 1$$

Which leads us to the fact that $\gcd(30030, 257) = 1$. by assumption we have the unique prime factorization of $30030 = 2 \times 3 \times 5 \times 7 \times 11 \times 13$. By definition of \gcd we know that any of $A = \{2, 3, 5, 7, 11, 13\}$ is not a prime factor for 257. and on the other hand we have the fact that $\sqrt{257} \approx 16$ and none of primes up to 16 that are elements of set A divide 257 so there could not be any prime factor for 257 which yields that 257 is prime ■

(b)

$$4883 = 4369 (1) + 514$$

$$4369 = 514 (8) + 257$$

$$514 = 257 (2) + 0$$

By part a we know that 257 is prime and thus we factor numbers as follows. $\frac{4883}{257} = 19$, and because 19 is prime, for the factorization we have: $4883 = 257 \times 19$. With the same argument for 4369 we have the factorization, $4369 = 257 \times 17$.

• **Problem 2**

(a) We know that every linear combination of a and b in the form $ax + by$ for $x, y \in \mathbb{Z}$ is a multiple of their greatest common divisor $\gcd(a, b)$, by knowing this fact and that there is a linear combination of a, b such that $ax + by = 1$ it is trivial that $\gcd(a, b) = 1$ ■

(b) Assume that a is invertible mod b then the equation $ax \equiv 1 \pmod{b}$ has a solution for x , which by definition of modular equation leads us to the equation $ax = by + 1$ for some value of b , which means that $ax + by = 1$ for some values of x and y which means that $\gcd(a, b) = 1$.

Conversely assume that $\gcd(a, b) = 1$ which means that $ax + by = 1$ for some value of x and y now consider this equation modular b which gives us $\overline{ax} + \overline{by} \equiv \overline{1} \pmod{b}$ which means a is invertible modular b ■

(c)

$$101 = 17 (5) + 16 \quad (I)$$

$$17 = 16 (1) + 1 \quad (II)$$

By the equation (II) we can write 16 as follows $16 = 17 - 1$ by putting this into the equation (I) we have the equation $101 = 17(5) + (17 - 1)$ which leads us to $101 = 17(6) - 1$, Thus $1 = 17(6) - 101(1)$. Hence $17^{-1} \pmod{101} = 6$ ■

• **Problem 3**

(a)

$$\begin{cases} x \equiv 1 \pmod{4} \\ x \equiv 2 \pmod{6} \end{cases} \longrightarrow \gcd(4, 6) = 2$$

Assume for the sake of contradiction that there is a solution for this system. Means that there is a x such that.

$$\begin{cases} x \equiv 1 \pmod{4} \Rightarrow x = 4k + 1 \ (k \in \mathbb{Z}) \\ x \equiv 2 \pmod{6} \Rightarrow x = 6l + 2 \ (l \in \mathbb{Z}) \end{cases} \Rightarrow 4k + 1 = 6l + 2 \Rightarrow 4k = 6l + 1$$

But in the last equation the Left-Hand-Side is always even and the Right-Hand-Side is always odd which contradicts. Thus there is no such a solution for the system this completes the proof of our counterexample. ■

(b)

$$\begin{cases} x \equiv 2 \pmod{17} \\ x \equiv 9 \pmod{101} \end{cases}$$

By the first equation we have $x = 17k + 2$, then by putting this together with the second equation we have that $17k + 2 \equiv 9 \pmod{101}$ which leads us to $17k \equiv 7 \pmod{101}$. Since we have already calculated the inverse of 17 modulo 101 we can now evaluate the value of k modulo 101.

$$6 \times 17k \equiv 6 \times 7 \pmod{101} \Rightarrow k \equiv 42 \pmod{101} \Rightarrow k = 101u + 42$$

Now by putting this together with the last equation we have $x = 17(101u + 42) + 2$. Thus $x = 1717u + 716$. ■

• **Problem 4**

(a) Assume for the sake of contradiction that there are two identity elements e and e' , we get the contradiction by just applying the definition of identity element on each:

$$m(e, e') = \begin{cases} e & e' \text{ is identity} \\ e' & e \text{ is identity} \end{cases} \implies e = e'$$

Which contradicts so the identity element must be unique. ■

Assume for the sake of contradiction that arbitrary element x of our group has two inverses x_1^{-1} and x_2^{-1} .

$$\begin{cases} m(x, x_1^{-1}) = e \\ m(x, x_2^{-1}) = e \end{cases} \xrightarrow[\text{identity element}]{\text{uniqueness of}} m(x, x_1^{-1}) = m(x, x_2^{-1}) \Rightarrow x_1^{-1} = x_2^{-1} = x^{-1}$$

(b) We need to show (I)-Injectivity and (II)-Surjectivity

(I) Assume that $hg = h'g$ by multiplying g^{-1} to the both sides of equation we have $h = h'$.

(II) Take arbitrary element $h \in G$ we claim that $g^{-1}h$ is the element that maps under the function $m_g(h)$ to the element h . To prove the claim we have $m_g(g^{-1}h) = gg^{-1}h = h$ as desired.

So function is both injective and surjective together means that the function is Bijection. ■

- (c) (I) Closed under operator: This is easy to see that $ax + bz, \dots, cy + dt$ are real numbers since $a, b, c, d, x, y, z, t \in \mathbb{R}$

$$A \times B = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \times \begin{pmatrix} x & y \\ z & t \end{pmatrix} \Rightarrow \begin{cases} A \in GL_2(\mathbb{R}) \\ B \in GL_2(\mathbb{R}) \end{cases} \Rightarrow A \times B \in GL_2(\mathbb{R})$$

- (II) Existence of Inverse: This follows immediately from the fact the determinant is non-zero and basic Linear-Algebra. Identity element is just identity matrix I_2

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \frac{1}{1} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

- (III) $(A \times B) \times C = A \times (B \times C)$: This is easy to verify by Linear-Algebra. Just write it down it is straightforward.

This is not a commutative group. Counterexample is as follows:

$$\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 3 & 2 \end{pmatrix} \quad \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 3 & -1 \\ 2 & 0 \end{pmatrix}$$

- (d) (I) Closed under operator: $\bar{a} \times \bar{b} = \overline{a \times b}$, Thus it is closed.
 (II) Identity element is : $\bar{1}$ and inverse for every element exists since p is considered to be prime: $\gcd(p, a)$ is either p which means p divide a , This is not the case because every element of our group is to be considered lower than p , or $\gcd(p, a) = 1$ which by bezout's coefficients we know that there is an inverse for a modulo p .
 (III) $\bar{a} \times (\bar{b} \times \bar{c}) = (\bar{a} \times \bar{b}) \times \bar{c}$: which is trivially true by the definition.

Under addition is similar.

- (e) No it is not closed under operator: $\bar{3} \times \bar{5} = \bar{0}$
 The largest prime number before 15 is 13 so $\frac{\mathbb{Z}}{13\mathbb{Z}} \equiv \mathbb{F}_{13}$ is the maximal subset of $\frac{\mathbb{Z}}{15\mathbb{Z}}$ which is a group under multiplication.
 (f) We know that φ is a multiplicative function. Let's evaluate the φ for prime values:

$$\begin{aligned} \varphi(p) &= \#\{1, \dots, p-1\} = p-1 \\ \varphi(p^\alpha) &= \#\{1, \dots, p-1, p+1, \dots, p^2-1, p^2+1, \dots, p^\alpha-1\} = p^{\alpha-1}(p-1) \end{aligned}$$

Now let's evaluate the φ for arbitrary N with the factorization $N = p_1^{\alpha_1} \dots p_s^{\alpha_s}$.

$$\begin{aligned} \varphi(N) &= [\varphi(p_1^{\alpha_1})] \dots [\varphi(p_s^{\alpha_s})] = [p_1^{\alpha_1-1}(p_1-1)] \dots [p_s^{\alpha_s-1}(p_s-1)] \\ &\xrightarrow{\text{Closed form}} \prod_{\substack{p_i | N \\ p_i \text{ prime}}} p_i^{\alpha_i-1}(p_i-1) = \prod_{\substack{p_i | N \\ p_i \text{ prime}}} p_i^{\alpha_i} \left(1 - \frac{1}{p_i}\right) \end{aligned}$$

By multiplying $p_i^{\alpha_i}$ we get the formula as desired.

$$\varphi(N) = N \prod_{\substack{p_i | N \\ p_i \text{ prime}}} \left(1 - \frac{1}{p_i}\right)$$

- **Problem 5**

- (a) By definition of φ we know that $\varphi(N)$ is the number of number less than N and relatively prime to N . Assume that order of element a is d by Lagrange's Theorem this d should dividde $\varphi(N)$ so $d \times k = \varphi(N)$,

$$\text{Hence } a^{\varphi(N)} \stackrel{N}{\equiv} a^{d.k} \stackrel{N}{\equiv} 1$$

- (b)

$$31 = 7(4) + 3 \Rightarrow$$

$$3 = 31 + 7(-4)$$

$$7 = 3(2) + 1 \Rightarrow$$

$$1 = 7 + 3(-2)$$

$$1 = 7 + [31 + 7(-4)](-2) = 31(-2) + 7(9)$$

Hence inverse of 7 modulo 31 is 9 i.e $7 \times 9 \stackrel{31}{\equiv} 1$

$$m^7 \stackrel{31}{\equiv} x \Rightarrow (m^7)^e \stackrel{31}{\equiv} x^e \Rightarrow 7 \times e \stackrel{\varphi(31)}{\equiv} 1 \Rightarrow \xi_t$$

- **Problem 6**

- **Problem 7**

- **Problem 8**