

Parsing Binomials & Multinomials in Probability

Andrew Jones

Introduction

The use of the Binomial and Multinomial theorems in Probability can often not be intuitively obvious. This text intends to highlight the relations between computing polynomials from binomials with combinations, total probability, and probability mass functions.

1 Binomial Theorem

The binomial theorem expresses the expansion of two monomial terms such as $(x + y)^2 = x^2 + 2xy + y^2$. However, in probability the binomial theorem can express the probability of all combinations of two independent events.

Example 1. Let an unfair coin be flipped twice with $P(Tails) = 0.3$ and $P(Heads) = 0.7$

We know the probability must sum to 1. In two flips then, $(T + H)^2 = T^2 + 2TH + H^2$. This aligns with the outcomes of TT , TH , HT , and HH for 2 flips. Substituting in the probabilities we have $0.3^2 + 2 * 0.3 * 0.7 + 0.7^2 = 1$.

To compute the expansion of two monomials the theorem uses:

Definition 1 (Factorial $n!$). Count every way to permute a set of n distinct objects

$$n! = \prod_{i=1}^n i$$

where $0! = 1$ and $n \geq 0$.

Building on factorials, the theorem uses the

Definition 2 (Binomial Coefficient). Count every way to combine a set of n objects of k size

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Note that the binomial coefficient can be used to determine the number of combinations of a particular size the binomial theorem produces.

Example 2. 3 flips of a coin yields:

$$\begin{aligned}(T + H)^3 &= TTT + TTH + THT + HTT + HHT + HTH + THH + HHH \\&= T^3 + 3T^2H + 3H^2T + H^3 \\ \binom{3}{3} &= 1 \text{ hence } T^3 = TTT \text{ or } H^3 = HHH \\ \binom{3}{2} &= 3 \text{ hence } 3T^2H = TTH + THT + HTT \text{ or } 3H^2T = HHT + HTH + THH \\ 2^3 &= T^3 + 3T^2H + 3H^2T + H^3 = 1 + 3 + 3 + 1 = 8 \text{ Total Combinations}\end{aligned}$$

Many combinations can be simplified by the use of Pascal's identity:

Pascal's Identity.

$$\begin{aligned}
\binom{n-1}{k} + \binom{n-1}{k-1} &= \frac{(n-1)!}{k!(n-1-k)!} + \frac{(n-1)!}{(k-1)!(n-k)!} \\
&= (n-1)! \left[\frac{n-k}{k!(n-k)!} + \frac{k}{k(n-k)!} \right] \\
&= (n-1)! \frac{n}{k!(n-k)!} \\
&= \frac{n!}{k!(n-k)!} \\
&= \binom{n}{k}
\end{aligned}$$

□

Binomial Theorem. Assume that $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$ and by the definition of the binomial coefficient $n \geq 0$. For the case $(n=0) \rightarrow (a+b)^0 = 1$. For the case $n \geq 0$.

$$\begin{aligned}
(a+b)^{n+1} &= (a+b)(a+b)^n = (a+b) \sum_{k=0}^n \binom{n}{k} a^k b^{n-k} \\
&= \sum_{k=0}^n \binom{n}{k} a^{k+1} b^{n-k} + \sum_{k=0}^n \binom{n}{k} a^k b^{n-k+1} \\
&\quad m = k+1 \\
&= \sum_{m=1}^{n+1} \binom{n}{m-1} a^m b^{n-m+1} + \sum_{k=0}^n \binom{n}{k} a^k b^{n-k+1} \\
&= b^{n+1} + \sum_{k=1}^n \left[\binom{n}{k} + \binom{n}{k-1} \right] a^k b^{n-k+1} + a^{n+1} \\
&= b^{n+1} + \sum_{k=1}^n \binom{n+1}{k} a^k b^{n-k+1} + a^{n+1}
\end{aligned}$$

□

2 Multinomial Theorem

While the binomial theorem works for 2 independent events, the multinomial theorem generalizes to any number of groups/events.

Definition 3 (Multinomial Coefficient).

$$\binom{N}{n_1, \dots, n_r} = \frac{N!}{n_1! \dots n_r!}$$

Where n_1 to n_r are different group sizes.

Example 3. For 13 items we want to know how many combinations of 5, 5, and 3 can be made

$$\begin{aligned}\binom{13}{5,5,3} &= \binom{13}{5} \binom{8}{5} \binom{3}{3} \\ &= \frac{13!}{5!(13-5)!} \frac{8!}{5!(8-5)!} \frac{3!}{3!(3-3)!} \\ &= \frac{13!}{5!5!3!}\end{aligned}$$

Proof.

□

Remark 1. This is interesting since...

3 Possible Outcomes to Equations