Synthetic Homotopy Theory

Exercises and Solutions for the

Midlands Graduate School (MGS)
13–17 April 2026, Nottingham, UK

by
Axel Ljungström and Aref Mohammadzadeh
School of Computer Science
University of Nottingham

Heavily based on notes from MGS 2024 by Ulrik Buchholtz and Mark Williams

NB Subject to change: Refresh if needed

Contents

1 Truncated and connected types **1**
1.1 Exercises 1
1.2 Solutions 3

2 A menagerie of higher inductive types **4**
2.1 Exercises 4
2.2 Solutions 7

3 The Hopf fibration and the LES **10**
3.1 Exercises 10

4 Freudenthal and Degrees **11**
4.1 Exercises 11

Chapter 1

Truncated and connected types

1.1 Exercises

1. Show that for $n \geq m$ and any type A we have

$$\|\|A\|_n\|_m \simeq \|A\|_m.$$

2. Show that a type X is n-connected if and only if, for every n-type Y, the diagonal map

$$Y \to (X \to Y), \quad y \mapsto (x \mapsto y),$$

is an equivalence.

3. Show that the map $|_|_n : X \to \|X\|_n$ is n-connected.

4. Consider a map $f : X \to Y$. We define its **n-image** to be the type $\mathrm{im}_n(f) := (y : Y) \times \|\mathrm{fib}_f(y)\|_n$.

   a) Show that we can factorise f through the n-image, that is, construct a commuting diagram:

$$\begin{tikzcd} X \arrow[rr, "f"] \arrow[dr, "p"'] & & Y \\ & \mathrm{im}_n(f) \arrow[ur, "i"'] & \end{tikzcd}$$

   b) Show that p is n-connected and i is n-truncated.

   c) Show that if Z is any type, $g : X \to Z$ is n-connected, and $h : Z \to Y$ is n-truncated with $f = h \circ g$, then there is a unique equivalence

$e : \mathrm{im}_n(f) \to Z$ with $e \circ p = g$ and $h \circ e = i$:

$$\begin{tikzcd}
 & Z \arrow[dr, "h"] \arrow[dd, dotted, "e"] & \\
X \arrow[ur, "g"] \arrow[rr, "f"] \arrow[dr, "p"'] & & Y \\
 & \mathrm{im}_n(f) \arrow[ur, "i"'] &
\end{tikzcd}$$

(*Hint*) It's easier if you assume g and h are projections of dependent sums.

5. (Eckmann–Hilton). All paths will take place in a fixed type X.

   a) Given paths $p_0, p_1 : x = y$ and $q_0, q_1 : y = z$, and higher paths $r : p_0 = p_1$ and $s : q_0 = q_1$, define their **horizontal multiplication** $s * r : q_0 \cdot p_0 = q_1 \cdot p_1$ by path induction on r and p_0.

$$\begin{tikzcd}
x \arrow[r, bend left, "p_0"] \arrow[r, bend right, "p_1"'] \arrow[r, phantom, "\Downarrow r"] & y \arrow[r, bend left, "q_0"] \arrow[r, bend right, "q_1"'] \arrow[r, phantom, "\Downarrow s"] & z
\end{tikzcd}$$

   b) Given paths fitting into the following diagram

$$\begin{tikzcd}
x \arrow[r, bend left, "p_0"] \arrow[r, bend right, "p_1"'] \arrow[r, phantom, shift left=3, "r_0\Downarrow"] \arrow[r, phantom, shift right=3, "r_1\Downarrow"] & y \arrow[r, bend left, "q_0"] \arrow[r, bend right, "q_1"'] \arrow[r, phantom, shift left=3, "\Downarrow s_0"] \arrow[r, phantom, shift right=3, "\Downarrow s_1"] & z
\end{tikzcd}$$

   show we have the **interchange law**:

$$(s_1 * r_1) \cdot (s_0 * r_0) = (s_1 \cdot s_0) * (r_1 \cdot r_0)$$

   c) Deduce that for $n \geq 2$, the path composition in $\Omega^n(X, x)$ is commutative, and thus that $\pi_n(X, x)$ is abelian.

6. Show that when X is connected, that $\pi_n(X, x_0) \cong \pi_n(X, x_1)$ are merely isomorphic for any $x_0, x_1 : X$, that is, construct an element of the type

$$(x_0, x_1 : X) \to \|\pi_n(X, x_0) \simeq \pi_n(X, x_1)\|.$$

7. We define the observational equality type family of the natural numbers $E : \mathbb{N} \to \mathbb{N} \to \mathcal{U}$ by

$$\begin{aligned}
E(0, 0) &= 1 \\
E(\mathrm{succ}(n), 0) &= 0 \\
E(0, \mathrm{succ}(n)) &= 0 \\
E(\mathrm{succ}(n), \mathrm{succ}(m)) &= E(n, m)
\end{aligned}$$

a) Define a natural map $n = m \to E(n, m)$ for all $n, m : \mathbb{N}$.

b) Use the fundamental theorem of identity types to show this natural map is an equivalence.

8. Show that a type X is $n+1$-truncated if and only if the diagonal map of X,
$$\delta_X : X \to X \times X, \quad x \mapsto (x, x),$$
is n-truncated.

9. Derive the usual elimination rules for the $\exists$- and $\vee$-connectives on propositions, using the definitions as truncated Σ- and $+$-types, respectively.

10. Show that $\|X\|_0$ satisfies the universal property of the set quotient of X modulo the equivalence relation $\sim$, where $x \sim x'$ if and only if $\|x =_X x'\|$.

    a) The obvious way of defining the type of arrows over the universe $(x : X)$

    This means, in particular, that we have an automorpism $(x : X) \times (X \to Y) \simeq (x : X) \times (X \to Y)$ defined by
$$(X \xrightarrow{f} Y) \mapsto ((y : Y) \times (\mathrm{fib}_f(y)) \xrightarrow{\mathrm{fst}} Y).$$
    As a result, anytime we wish to prove a property ranging over the the slice $\mathcal{U}/Y :=$

1.2 Solutions

1.

2.

3.

4.

5.

6.

7.

8.

9.

10.

11.

Chapter 2

A menagerie of higher inductive types

2.1 Exercises

1. (Pushout-Pullback Duality) Consider a commutating square:

$$\begin{tikzcd} A \arrow[r, "g"] \arrow[d, "f"'] & C \arrow[d, "j"] \\ B \arrow[r, "i"'] & X \end{tikzcd}$$

   with $H : i \circ f \sim j \circ g$. Then the following are equivalent:

   (i) The square is a pushout square.

   (ii) the square

$$\begin{tikzcd} T^X \arrow[r, "-\circ j"] \arrow[d, "-\circ i"'] & T^C \arrow[d, "-\circ g"] \\ T^B \arrow[r, "-\circ f"'] & T^A \end{tikzcd}$$

   is a pullback square, for every type T.

2. Show that the two definitions of S^1 are equivalent, in two ways:

   (i) Show the HIT for S^1 is the same thing as $\Sigma 2$ by constructing an equivalence between them.

   (ii) By applying Exercise 2.1.

3. Show that evaluation at the loop gives pointed equivalence

$$(S^1 \to_\star X) \to_\star \Omega X$$

for any pointed type X.

Conclude, using the suspension–loop adjunction, that we have a natural equivalence

$$(S^n \to_\star X) \to_\star \Omega^n X$$

for all $n : \mathbb{N}$.

4. Show that a type X is a proposition iff $\mathrm{inl} : X \to X * X$ is an equivalence.

5. Show that $A \vee B \simeq A * B$ for propositions A, B, where $A \vee B := \|A + B\|$. (Beware the notational overload between the wedge sum and disjunction: It's not too bad in practice, because the only pointed proposition is the unit type!)

6. (i) Consider a span $A \xleftarrow{f} X \xrightarrow{g} B$. Show that the following square is pushout:

$$\begin{tikzcd} X + X \arrow[r, "{[\mathrm{id},\mathrm{id}]}"] \arrow[d, "f+g"'] \arrow[dr, phantom, "\ulcorner"] & X \arrow[d, "inr \circ g"] \\ A + B \arrow[r, "{[\mathrm{inl},\mathrm{inr}]}"'] & A \amalg_X B \end{tikzcd}$$

   (ii) Apply this to show that for every type X we get $\Sigma X \simeq \mathsf{bool} * X$.

   . (Harder) Show that $S^1 \wedge X \simeq \Sigma X$

7. (Smash Product - Pointed Map Space Adjunction)

   (i) Prove that for each pointed type B, the fucnction $(-) \wedge B$ is the left adjoint to the function $B \to_* (-)$,i.e. for every pointed types A and C, there is a pointed equivalence:

$$A \to_* (B \to_* C) \simeq_* (A \wedge B) \to_* C$$

   (ii) Use this to prove that $S^1 \wedge X \simeq \Sigma X$ for each type X.

8. Show that the smash product is equivalently the pushout:

$$\begin{tikzcd} X + Y \arrow[r] \arrow[d] & X \times Y \arrow[d] \\ 1 + 1 \arrow[r] & X \wedge Y \end{tikzcd}$$

9. (i) For all pointed types X, Y, and A, prove there is a pointed equivalence:

$$X * Y \to_* A \simeq_* X \to (Y \to_* \Omega A)$$

   (ii) For pointed types X, Y prove that

$$X * Y \simeq_* \Sigma(X \wedge Y).$$

10. The torus, T^2, is often drawn as a (filled) square with side identifications

$$\begin{tikzcd} \star \arrow[r, "\ell_1"] & \star \\ \star \arrow[u, "\ell_2"] \arrow[r, "\ell_1"'] & \star \arrow[u, "\ell_2"'] \end{tikzcd} \tag{2.1}$$

In HoTT, we can capture this as HIT with

- A point constructor $\star : T^2$
- Two paths $\ell_1, \ell_2 : \star = \star$
- A filler sq of the square of paths in (2.1).

Show this type is equivalent to $S^1 \times S^1$. (*Hint*) You may assume that eliminators compute definitionally on point constructors (as in cubical type theory), otherwise it's quite intricate!

11. We can define the unit interval type $\mathbb{I}$ to be the pushout of the constant span $1 \leftarrow 1 \rightarrow 1$. Working in Martin-Löf type theory *without* univalence (and function extensionality) but with pushouts with strict definitional computation rules, prove that

    (i) $[((x, y) : A \times A) \times (x = y)] \simeq (\mathbb{I} \rightarrow A)$

    (ii) Use the above equivalence (or, if you prefer, a variation thereof) to derive function extensionality.

12. (i) Show that if the square

$$\begin{tikzcd} A \arrow[r, "g"] \arrow[d, "f"'] & C \arrow[d, "j"] \\ B \arrow[r, "i"'] & X \end{tikzcd}$$

is a pullback, then so is

$$\begin{tikzcd} \Omega A \arrow[r, "\Omega g"] \arrow[d, "\Omega f"'] & \Omega C \arrow[d, "\Omega j"] \\ \Omega B \arrow[r, "\Omega i"'] & \Omega X \end{tikzcd}$$

    (ii) For a map $A \xrightarrow{f} B$ show that $\Omega\,(\mathrm{fib}\ f) \simeq \mathrm{fib}\,(\Omega f)$

13. Show that the square

$$\begin{tikzcd} S^1 \arrow[r] \arrow[d] & 1 \arrow[d] \\ S^1 \times S^1 \arrow[r] & S^2 \vee S^1 \end{tikzcd}$$

is a pushout.

2.2 Solutions

1.

2.

3.

4.

5.

6.

7.

8.

9.

10.

11. Write the interval as the pushout of the constant span $1 \leftarrow 1 \rightarrow 1$, with points $0, 1 : \mathbb{I}$ and path $\mathsf{seg} : 0 = 1$.

    (i) Define
    $$\Phi : \Big(\sum_{x,y:A} (x = y)\Big) \to (\mathbb{I} \to A)$$
    by interval recursion:
    $$\Phi(x, y, p) \;:=\; \mathsf{rec}_{\mathbb{I}}(x, y, p).$$
    Thus, by the strict β-rules, we get:
    $$\Phi(x, y, p)(0) \equiv x, \qquad \Phi(x, y, p)(1) \equiv y, \qquad \mathrm{ap}_{\Phi(x,y,p)}(\mathsf{seg}) \equiv p.$$
    Conversely, define
    $$\Psi : (\mathbb{I} \to A) \to \sum_{x,y:A} (x = y)$$
    by
    $$\Psi(h) \;:=\; \big(h(0),\ h(1),\ \mathrm{ap}_h(\mathsf{seg})\big).$$
    Then
    $$\Psi(\Phi(x, y, p)) \equiv (x, y, p)$$
    judgmentally, by the strict β-rules above. For the other composite, if $h : \mathbb{I} \to A$, then
    $$\Phi(\Psi(h)) = \Phi\big(h(0), h(1), \mathrm{ap}_h(\mathsf{seg})\big)$$

is the map out of $\mathbb{I}$ with the same values on $0, 1$ and the same action on seg as h; hence, by the η-rule of the pushout $\mathbb{I}$,

$$\Phi(\Psi(h)) = h.$$

Therefore

$$\Big(\sum_{x,y:A}(x = y)\Big) \simeq (\mathbb{I} \to A).$$

Note that we do not actually need strict β-rule for this part.

(ii) First we derive non-dependent function extensionality. Let $f, g : A \to B$ and assume

$$H : \prod_{a:A}(f(a) = g(a)).$$

By part (i), for each $a : A$ we obtain a map

$$\widetilde{H}(a) : \mathbb{I} \to B$$

such that

$$\widetilde{H}(a)(0) \equiv f(a), \qquad \widetilde{H}(a)(1) \equiv g(a), \qquad \mathsf{ap}_{\widetilde{H}(a)}(\mathsf{seg}) \equiv H(a).$$

Now curry these maps:

$$K : \mathbb{I} \to (A \to B), \qquad K(i)(a) := \widetilde{H}(a)(i).$$

Then, using the strict computation rules from part (i) and the judgmental η-rule for functions,

$$K(0) \equiv \lambda a.\ \widetilde{H}(a)(0) \equiv \lambda a.\ f(a) \equiv f,$$

and similarly

$$K(1) \equiv g.$$

Hence

$$\mathsf{ap}_K(\mathsf{seg}) : f = g.$$

So ordinary function extensionality holds.

To obtain dependent function extensionality, let $B : A \to \mathcal{U}$, let

$$f, g : \prod_{a:A} B(a), \qquad H : \prod_{a:A}(f(a) = g(a)).$$

Define

$$f', g' : A \to \sum_{a:A} B(a), \qquad f'(a) := (a, f(a)), \quad g'(a) := (a, g(a)).$$

For each $a : A$, using $H(a)$ we get

$$f'(a) = g'(a)$$

in the Σ-type, namely by the path $(\mathrm{refl}_a, H(a))$. By ordinary function extensionality, this yields

$$p : f' = g'.$$

Now apply ap to the map

$$s : \left(A \to \sum_{a:A} B(a)\right) \to \prod_{a:A} B(a), \qquad s(k)(a) := \pi_2(k(a)).$$

Since $s(f') = f$ and $s(g') = g$, we obtain

$$\mathrm{ap}_s(p) : f = g.$$

Thus full function extensionality follows. It seems that we can prove (ii) without assuming η-rule for $\mathbb{I}$, right? Since we only need the right to left map from (i), and not the whole equivalence.

12. (i) Assume the given square is a pullback. Then $A \simeq B \times_X C$. Since for any pointed type Y we have

$$\Omega Y \simeq 1 \times_Y 1,$$

and pullbacks commute with pullbacks (by 3-3 lemma for pullbacks), it follows that

$$\Omega A \;\simeq\; \Omega(B \times_X C) \;\simeq\; (1 \times_B 1) \times_{(1 \times_X 1)} (1 \times_C 1) \;\simeq\; \Omega B \times_{\Omega X} \Omega C.$$

(ii) The homotopy fiber of f is the pullback

$$\mathrm{fib}(f) \simeq 1 \times_B A.$$

Apply (i) to this pullback square.

13. Using the equation $S^1 \wedge X \simeq \Sigma X$ we can identify S^2 with $S^1 \wedge S^1$, which is pushout of $S^1 \vee S^1 \to S^1 \times S^1$ along the unit map. Now we have:

$$\begin{tikzcd}
S^1 \vee S^1 \arrow[r] \arrow[d] & 1 \arrow[r] \arrow[d] & S^1 \arrow[d] \\
S^1 \times S^1 \arrow[r] & S^2 \arrow[r] & S^2 \vee S^1
\end{tikzcd}$$

Since both squares are pushouts, the pasting lemma for pushout says that the outer rectangle is a pushout. We get the goal by observing that the upper square in the following diagram is a pushout:

$$\begin{tikzcd}
S^1 \arrow[r] \arrow[d] & 1 \arrow[d] \\
S^1 \vee S^1 \arrow[r] \arrow[d] & S^1 \arrow[d] \\
S^1 \times S^1 \arrow[r] & S^2 \vee S^1
\end{tikzcd}$$

Chapter 3

The Hopf fibration and the long exact sequence

3.1 Exercises

1. Construct the product on the circle directly by double circle induction.

2. Give another proof of **??** by showing that $\mathrm{ap}_\varphi : (* = *) \to (\varphi(*) = \varphi(*))$ is an equivalence.

3. Construct a retraction $\rho : \Omega S^2 \to S^1$ of $\eta : S^1 \to \Omega S^2$. (*Hint*) Use the Hopf fibration.

4. We can define a "tensor product" of $\mathbb{Z}$-torsors by setting

$$(X, t) \otimes (Y, u) := (X \times Y/\sim, s)$$

   using the set quotient by the equivalence relation generated by $(t(x), y) \sim (x, u(y))$, and setting $s[(x, y)] := [t(x), y)]$. Check that this operation is well defined and gives an H-space structure on $\mathrm{Tors}_{\mathbb{Z}}$ with the base point $(\mathbb{Z}, \mathrm{suc})$ as neutral element.

5. Investigate some other components of $\mathrm{Set}_{\circlearrowleft}$, in particular those containing $(\mathrm{Fin}\, n, \mathrm{suc})$, where $\mathrm{Fin}\, n = (k : \mathbb{N}) \times (k < n)$ and the successor operation is taken modulo n. If this component is denoted Cyc_n (for n-cycles), what are the maps $\mathrm{Cyc}_n \to \mathrm{Cyc}_m$?

Chapter 4

The Freudenthal Suspension Theorem and Degrees

4.1 Exercises

1. Deduce the dual Blakers–Massey **??** from the Join Connectivity **??** by identifying the fibers of the cogap map with the joins of the fibers of legs in the cospan.

2. Deduce the Wedge Connectivity Lemma from the dual Blakers–Massey **??**, i.e., consider the pullback square, for pointed types X, Y,

$$\begin{tikzcd} 1 \arrow[r] \arrow[d] \arrow[dr, phantom, "\lrcorner", very near start] & Y \arrow[d] \\ X \arrow[r] & X \times Y. \end{tikzcd}$$

   Show that if X is n-connected and Y is m-connected, then the wedge inclusion map $X \vee Y \to X \times Y$ is $(n+m)$-connected.

3. Show that the counit $\varepsilon : \Sigma\Omega X \to X$ of the loop–suspension adjunction is $2n$-connected if X is n-connected.

4. Show that the pushout of an embedding is an embedding, i.e., if g is (-1)-truncated, then so is inl in the pushout square:

$$\begin{tikzcd} R \arrow[r, "g"] \arrow[d, "f"'] & B \arrow[d, "\mathrm{inr}"] \\ A \arrow[r, "\mathrm{inl}"'] \arrow[ur, phantom, "\ulcorner", very near start] & Q \end{tikzcd}$$