

Edge Detection from Fourier Phase Data

Alexander Reynolds Alena Chang

Advisors: Dr. Anne Gelb, Dr. Jacqueline Davis

Arizona State University
School of Mathematical and Statistical Sciences

MCTP Summer 2015



What is Edge Detection?

Edge detection is a sub-field of signal processing, which consists of the theory, applications, algorithms, and implementations of processing or transferring information via signals. Edge detection is the study of mathematical methods which identify points in a digital image where the image brightness changes sharply. An edge is a sudden shift, or jump, of a signal.

Example: Edge Detection on an Image



Applications of Edge Detection

Primary applications:

- Magnetic Resonance Imaging (MRI)



Applications of Edge Detection

Primary applications:


- Magnetic Resonance Imaging (MRI)
- Synthetic Aperture Radar (SAR)



Applications of Edge Detection

Primary applications:


- Magnetic Resonance Imaging (MRI)
- Synthetic Aperture Radar (SAR)
- Computer/Machine Vision (e.g. Kinect)



Applications of Edge Detection

Primary applications:

- Magnetic Resonance Imaging (MRI)
- Synthetic Aperture Radar (SAR)
- Computer/Machine Vision (e.g. Kinect)
- 1-D Case: Discontinuities of a Function



What is Fourier data?

What is Fourier data?

Fourier Series

$$f(x) = \sum_{k=-\infty}^{\infty} \hat{f}_k e^{ikx}.$$


The coefficients \hat{f}_k are called the **Fourier coefficients** of f . For a piecewise-smooth function periodic on $[-\pi, \pi)$, they are defined as:

$$\hat{f}_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx, \quad k = -N, \dots, N.$$

What is the Concentration Edge Detection Method?

First, define the **jump function** of f , $[f](x)$ as the difference between the right and left hand limits of the function f at every point x ; i.e.,


$$[f](x) = f(x^+) - f(x^-).$$



What is the Concentration Edge Detection Method?

First, define the **jump function** of f , $[f](x)$ as the difference between the right and left hand limits of the function f at every point x ; i.e.,

$$[f](x) = f(x^+) - f(x^-).$$



The concentration edge detection method computes an approximation to the jump function using:

Partial Sum Approximation to the Jump Function

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x).$$

Analytical Concentration Factors

The concentration factor, $\sigma(k)$, concentrates the partial sum along the edges in the function f . There are multiple analytical solutions σ ; the three main types we are using are:

Factor	Expression	Remarks
Trigonometric	$\sigma(\eta) = \frac{\pi \sin(\pi\eta)}{\text{Si}(\pi)}$	$\text{Si}(\pi) = \int_0^\pi \frac{\sin(x)}{x} dx$
Polynomial	$\sigma(\eta) = p\pi\eta^p$	p is the order of the factor
Exponential	$\sigma(\eta) = C\eta e^{\frac{1}{\alpha\eta(\eta-1)}}$	α is the order C is a normalizing constant $C = \frac{\pi}{\int_{\frac{1}{N}}^{1-\frac{1}{N}} \exp(\frac{1}{\alpha\tau(\tau-1)}) d\tau}$

Jump Reconstruction from Fourier Data




Figure: Reconstructions of $[f](x)$ with multiple different σ .

Designed Concentration Factors

How can we design *other* concentration factors?



Designed Concentration Factors

How can we design *other* concentration factors? We want

Partial Sum Approximation of $[f](x)$

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x).$$

Designed Concentration Factors

How can we design *other* concentration factors? We want


Partial Sum Approximation of $[f](x)$

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x).$$

We can engineer σ for a function $g(x)$ where we know what $[g](x)$ is.

Designed Concentration Factors

Consider the ramp function $r(x)$ with the associated jump function $[r](x)$ on $[-\pi, \pi]$:




$$r(x) = \begin{cases} \frac{1}{2\pi}(-x - \pi) & \text{if } x < 0, \\ \frac{1}{2\pi}(-x + \pi) & \text{if } x \geq 0. \end{cases} \quad [r](x) = \begin{cases} 1 & \text{if } x = 0, \\ 0 & \text{otherwise.} \end{cases}$$



Designed Concentration Factors

Consider the ramp function $r(x)$ with the associated jump function $[r](x)$ on $[-\pi, \pi]$:



$$S_N^\sigma[r](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{r}_k e^{ikx} \approx [r](x).$$

Designed Concentration Factors

We can solve the system $S\sigma = [r](x)$ for a solution σ to

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x).$$

Designed Concentration Factors

We can solve the system $S\sigma = [r](x)$ for a solution σ to

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x).$$




Figure: Reconstruction of $[f](x)$ from σ and \hat{f}

Fourier Phase Data

What is Fourier Phase Data?



Fourier Phase Data

What is Fourier Phase Data? Recall

Fourier Series

$$f(x) = \sum_{k=-\infty}^{\infty} \hat{f}_k e^{ikx}, \text{ where } \hat{f}_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx.$$

The Fourier coefficients \hat{f}_k are complex numbers, i.e., $\hat{f}_k = \alpha_k e^{i\theta_k}$, where α_k is the **amplitude** and $e^{i\theta_k}$ is the **phase**.



Fourier Phase Data

What is Fourier Phase Data? Recall

Fourier Series

$$f(x) = \sum_{k=-\infty}^{\infty} \hat{f}_k e^{ikx}, \text{ where } \hat{f}_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx.$$

The Fourier coefficients \hat{f}_k are complex numbers, i.e., $\hat{f}_k = \alpha_k e^{i\theta_k}$, where α_k is the **amplitude** and $e^{i\theta_k}$ is the **phase**.

Let \tilde{f}_k be the Fourier phase data, i.e., $\tilde{f}_k = \frac{\hat{f}_k}{\|\hat{f}_k\|} = e^{i\theta_k}$.



Fourier Phase Data

What is Fourier Phase Data? Recall

Fourier Series

$$f(x) = \sum_{k=-\infty}^{\infty} \hat{f}_k e^{ikx}, \text{ where } \hat{f}_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-ikx} dx.$$

The Fourier coefficients \hat{f}_k are complex numbers, i.e., $\hat{f}_k = \alpha_k e^{i\theta_k}$, where α_k is the **amplitude** and $e^{i\theta_k}$ is the **phase**.

Let \tilde{f}_k be the Fourier phase data, i.e., $\tilde{f}_k = \frac{\hat{f}_k}{\|\hat{f}_k\|} = e^{i\theta_k}$.

Is it still possible to reconstruct the jumps from \tilde{f}_k ?



Jump Reconstruction from Fourier Phase Data

Is it still possible to reconstruct the jumps from \tilde{f}_k ?



Jump Reconstruction from Fourier Phase Data

Is it still possible to reconstruct the jumps from \tilde{f}_k ?

Partial Sum Approximations of $[f](x)$

$$S_N^\sigma[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \sigma_k \hat{f}_k e^{ikx} \approx [f](x)$$

$$S_N^{\tilde{\sigma}}[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \tilde{\sigma}_k \tilde{f}_k e^{ikx} \approx [f](x)$$

$$\sum_{k=-N}^N \sigma_k \hat{f}_k = \sum_{k=-N}^N \tilde{\sigma}_k \tilde{f}_k$$

Jump Reconstruction from Fourier Phase Data

Yes!



Jump Reconstruction from Fourier Phase Data

Yes! We can solve the system $S\tilde{\sigma} = [r](x)$ for a solution $\tilde{\sigma}$ to

$$S_N^{\tilde{\sigma}}[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \tilde{\sigma}_k \tilde{f}_k e^{ikx} \approx [f](x).$$

Jump Reconstruction from Fourier Phase Data

Yes! We can solve the system $S\tilde{\sigma} = [r](x)$ for a solution $\tilde{\sigma}$ to

$$S_N^{\tilde{\sigma}}[f](x) = \sum_{k=-N}^N i \operatorname{sgn}(k) \tilde{\sigma}_k \tilde{f}_k e^{ikx} \approx [f](x).$$




Figure: Reconstruction of $[f](x)$ from $\tilde{\sigma}$ and \tilde{f}

Jump Reconstruction from Fourier Phase Data




Figure: Reconstructions of multiple functions from $\tilde{\sigma}$ and \tilde{f} .

Questions that remain:

- What if the phase data is noisy?
- What if the phase data is intermittent, i.e., banded?
- What if the phase data is noisy and intermittent?
- Generalizations to two dimensions

Acknowledgements



We would like to thank the NSF for their gracious support of the program, and Dr. Kostelich for his dedication in continuing this program at ASU.

We would also like to thank Dr. Gelb and everyone in our group for their mentorship:

- Dr. Anne Gelb
- Dr. Jacqueline Davis
- Dr. Rodrigo Platte
- Denny Denker
- Shane Lubold
- Rachel Nahon

