

# Applied Data Analysis (ADA)

Summer term 2020

Erhard Cramer, Maria Kateri

Institute of Statistics  
Pontdriesch 14-16

[erhard.cramer@rwth-aachen.de](mailto:erhard.cramer@rwth-aachen.de), [maria.kateri@rwth-aachen.de](mailto:maria.kateri@rwth-aachen.de)  
[www.isw.rwth-aachen.de](http://www.isw.rwth-aachen.de)

## ➤ A preliminary note

- **Please read carefully the slides on the course concept uploaded in RWTHmoodle!**
- The lecture is split into two parts:
  - Part I: **Linear Models** (Cramer)  
Lectures from April 20 to May 18  
Tutorials: May 6, 20, June 8
  - Part II: **Generalized Linear Models** (Kateri)  
Lectures from May 25 to July 13  
Tutorials: June 17, July 1, 15
  - break June 1–5 (Pentecost week)
- Part I will be held as distance teaching course.
- The teaching concept of Part II depends on the regulations for universities due to the Corona crisis. Information will be provided in due time.

# Just to let you know who is talking to you on linear models...

## ▶ Prof. Dr. Erhard Cramer

- ▶ Institute of Statistics
- ▶ Pontdriesch 14-16
- ▶ Office: Room no. 313, third floor
- ▶ [erhard.cramer@rwth-aachen.de](mailto:erhard.cramer@rwth-aachen.de)

## ▶ In the first part, we consider...

a class of statistical models, so called **linear models**, generated by the equation

$$Y = B\beta + \varepsilon$$

with

- ▶  $Y = (Y_1, \dots, Y_n)'$  vector of observations,
- ▶  $B$  design matrix,
- ▶  $\beta$  parameter vector
- ▶  $\varepsilon$  (random) error term (not observable).



# Part I: Linear Models

## Chapter I.1

### Preliminaries

Notation, Linear Algebra & Probability

## ▶ To be discussed...

- ▶ properties of vectors & matrices, rank, trace, singular value decomposition, etc.
- ▶ Moore-Penrose general inverse
- ▶ Image, kernel, orthogonal projectors
- ▶ Random vectors, expectations, covariance matrix
- ▶ selected probability distributions on the real line connected to the normal distribution
- ▶ non-central  $\chi^2$ - and F-distribution

# Part I: Linear Models

## Chapter I.1

### Preliminaries

Linear Algebra

# Notation & basic definitions

## 1.1.1 Notation (vectors and matrices)

➤  $\mathbb{R}^p$ :  $p$ -dimensional Euclidean space

➤  $\mathbb{R}^{p \times q}$ : set of all  $(p \times q)$ -matrices

➤ vectors are written in bold italics:  $\mathbf{x} = (x_i)_{1 \leq i \leq p} = \begin{pmatrix} x_1 \\ \vdots \\ x_p \end{pmatrix}$

➤ random vectors are written in capital bold italics:  $\mathbf{X} = (X_i)_{1 \leq i \leq p} = \begin{pmatrix} X_1 \\ \vdots \\ X_p \end{pmatrix}$

➤ matrices are written in capitals:  $A = (a_{ij})_{1 \leq i \leq p, 1 \leq j \leq q} = \begin{pmatrix} a_{11} & \cdots & a_{1q} \\ \vdots & \cdots & \vdots \\ a_{p1} & \cdots & a_{pq} \end{pmatrix}.$

# Notation & basic definitions

## 1.1.2 Notation (special vectors and matrices)

- $A = \text{diag}(a_1, \dots, a_p)$ : diagonal matrix with diagonal elements  $a_1, \dots, a_p$
- $\mathbb{1}_p \in \mathbb{R}^p$ : vector of ones,  $\mathbf{0} \in \mathbb{R}^p$  zero vector
- $e_{1,p}, \dots, e_{p,p}$ : standard basis of  $\mathbb{R}^p$
- $I_p = \text{diag}(1, \dots, 1)$ :  $p$ -dimensional identity matrix
- $\mathbb{1}_{p \times p} = \mathbb{1}_p \mathbb{1}_p'$ : matrix of ones
- $E_p = I_p - \frac{1}{p} \mathbb{1}_{p \times p}$ : ortho-projection matrix
- $\|x\| = \sqrt{\sum_{i=1}^p x_i^2}$  denotes the (Euclidean) norm of a vector  $x \in \mathbb{R}^p$ .
- $\text{rank}(A)$  denotes the rank of a matrix  $A$ .
- $\det(A)$  denotes the determinant of a squared matrix  $A$ .
- $\text{trace}(A)$  denotes the trace of a squared matrix  $A \in \mathbb{R}^{p \times p}$ , i.e.,  $\text{trace}(A) = \sum_{i=1}^p a_{ii}$
- The transpose of a matrix  $A$  is denoted by  $A'$ .
- The inverse of a matrix  $A \in \mathbb{R}^{p \times p}$  is denoted by  $A^{-1}$  (provided it exists), i.e.,  $AA^{-1} = A^{-1}A = I_p$ .



# Notation & basic definitions

## ➤ I.1.3 Definition

- A matrix  $A \in \mathbb{R}^{p \times p}$  is called symmetric if  $A = A'$ .
- A matrix  $A \in \mathbb{R}^{p \times p}$  is called an orthogonal matrix if  $AA' = A'A = I_p$ .
- A matrix  $A \in \mathbb{R}^{p \times p}$  is called positive (non-negative) definite if  $A = A'$  and

$$\mathbf{x}'A\mathbf{x} > (\geq) 0 \quad \forall \mathbf{x} \in \mathbb{R}^p \setminus \{\mathbf{0}\}.$$

For short, we write  $A > 0$  or  $A \geq 0$ , respectively.

- $\mathbb{R}_{>0}^{p \times p}$ : set of all positive definite  $(p \times p)$ -matrices
- $\mathbb{R}_{\geq 0}^{p \times p}$ : set of all non-negative definite  $(p \times p)$ -matrices

# Some linear algebra

## ► I.1.4 Lemma

Let  $A, C \in \mathbb{R}^{p \times p}$  with  $\det(AC) \neq 0$  and  $B \in \mathbb{R}^{k \times p}$ ,  $1 \leq k \leq p$ . Then:

- ①  $(C')^{-1} = (C^{-1})'$
- ②  $(AC)' = C'A'$
- ③  $(AC)^{-1} = C^{-1}A^{-1}$
- ④  $\text{rank}(B') = \text{rank}(B)$
- ⑤  $\text{rank}(BC) = \text{rank}(B)$
- ⑥ For  $D \in \mathbb{R}^{p \times k}$ , we have  $\text{rank}(BD) = \text{rank}(DB)$ .
- ⑦  $\text{rank}(BB') = \text{rank}(B'B) = \text{rank}(B)$
- ⑧  $\det(AC) = \det(CA) = \det(A) \cdot \det(C)$  for all  $A, C \in \mathbb{R}^{p \times p}$
- ⑨  $\text{trace}(A + C) = \text{trace}(A) + \text{trace}(C)$
- ⑩  $\text{trace}(BD) = \text{trace}(DB)$  for all  $D \in \mathbb{R}^{p \times k}$

# Singular Value Decomposition (SVD)

## ► I.1.5 Theorem

Let  $\Sigma \in \mathbb{R}_{\geq 0}^{p \times p}$ . Then, the **singular value decomposition** (eigen decomposition) of  $\Sigma$  is given by

$$\Sigma = V \Lambda V',$$

where  $\lambda_1 \geq \dots \geq \lambda_p \geq 0$  denote the eigenvalues and  $\mathbf{v}_1, \dots, \mathbf{v}_p$  the corresponding (orthonormal) eigenvectors of  $\Sigma$ . Further,  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$  and  $V = [\mathbf{v}_1 \mid \dots \mid \mathbf{v}_p]$  with  $V'V = VV' = I_p$ .

Furthermore, with the definitions  $\Lambda^{1/2} = \text{diag}(\sqrt{\lambda_1}, \dots, \sqrt{\lambda_p})$  and  $\Sigma^{1/2} = V \Lambda^{1/2} V'$ , we have

►  $\Sigma = \Sigma^{1/2} \Sigma^{1/2}$  and  $(\Sigma^{1/2})' = \Sigma^{1/2}$ .

►  $\Sigma^{1/2}$  is non-negative definite.

👉  $\Sigma^{1/2}$  is called the **root of  $\Sigma$** .

► Notice that, for a regular matrix  $\Sigma$ ,  $(\Sigma^{1/2})^{-1} = \Sigma^{-1/2}$  where  $\Sigma^{-1/2} = V \Lambda^{-1/2} V'$  and  $\Lambda^{-1/2} = \text{diag}(\sqrt{\lambda_1^{-1}}, \dots, \sqrt{\lambda_p^{-1}})$ .

# Moore-Penrose (general) inverse of a matrix

## 1.1.6 Theorem

The Moore-Penrose (general) inverse of a matrix  $A$  is denoted by  $A^+$ , i.e.,  $A^+$  is the unique matrix satisfying the four equations:

$$AA^+A = A, \quad A^+AA^+ = A^+, \quad (AA^+)' = AA^+, \quad (A^+A)' = A^+A.$$

It has the following properties:

- $(A^+)^+ = A$
- $(A^+)' = (A')^+$
- $A^+ = A^+(A^+)'A'$
- $A = AA'(A^+)'$
- $A' = A'AA^+$
- $A' = A^+AA'$
- $(AA')^+ = (A')^+A^+$
- $A^+ = (A'A)^+A'$
- $A^+ = A'(AA')^+$
- If  $A \in \mathbb{R}^{p \times q}$  exhibits the SVD  $A = U\Lambda V'$ , then  $A^+$  has the SVD  $A^+ = V\Lambda^+U'$  where  $\Lambda^+$  is the Moore-Penrose inverse of the matrix  $\Lambda$ .
- If  $A \in \mathbb{R}^{p \times p}$  is a regular matrix then  $A^+ = A^{-1}$ .

# Image, Kernel, Orthogonal Projectors

## 1.1.7 Definition

- For a matrix  $A \in \mathbb{R}^{p \times q}$ , let
  - $\text{Ker}(A) = \{\mathbf{x} \in \mathbb{R}^q \mid A\mathbf{x} = \mathbf{0}\}$  be the kernel (null space) of  $A$ .
  - $\text{Im}(A) = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^q\}$  be the image of  $A$ .
- For a linear subspace  $\mathcal{A} \subseteq \mathbb{R}^p$ ,  $\mathcal{A}^\perp = \{\mathbf{y} \in \mathbb{R}^p \mid \mathbf{x}'\mathbf{y} = 0 \text{ for all } \mathbf{x} \in \mathcal{A}\}$  denotes the corresponding orthogonal space.
- Let  $\mathcal{A}, \mathcal{B} \subseteq \mathbb{R}^p$  be linear subspaces with  $\mathcal{A} \cap \mathcal{B} = \{\mathbf{0}\}$ . Then,  $\mathcal{A} \oplus \mathcal{B} = \{\mathbf{x} + \mathbf{y} \mid \mathbf{x} \in \mathcal{A}, \mathbf{y} \in \mathcal{B}\}$  is called the direct sum of  $\mathcal{A}, \mathcal{B}$ .

Notice that  $\text{Ker}(A) \subseteq \mathbb{R}^q$  and  $\text{Im}(A) \subseteq \mathbb{R}^p$  are linear subspaces.

## 1.1.8 Definition

A matrix  $Q \in \mathbb{R}^{p \times p}$  is called

- idempotent if  $Q^2 = Q$
- orthogonal projector on a linear subspace  $\mathcal{A} \subseteq \mathbb{R}^p$  if
  - for any  $\mathbf{x} \in \mathcal{A}$ :  $Q\mathbf{x} = \mathbf{x}$
  - for any  $\mathbf{y} \in \mathcal{A}^\perp$ :  $Q\mathbf{y} = \mathbf{0}$

# Properties of Orthogonal Projectors and Moore-Penrose inverse

## ▶ I.1.9 Lemma

- ▶ Orthogonal projectors on a linear subspace  $\{0\} \neq \mathcal{A} \subseteq \mathbb{R}^p$  are unique.
- ▶  $Q \in \mathbb{R}^{p \times p}$  is an orthogonal projector (on  $\text{Im}(Q)$ ) iff  $Q^2 = Q$  and  $Q' = Q$ .

## ▶ I.1.10 Theorem

Let  $A \in \mathbb{R}^{p \times q}$  with Moore-Penrose inverse  $A^+$  and define  $P_1 = I_q - A^+A$ ,  $P_2 = I_p - AA^+$ . Then:

- ▶  $P_1$  and  $P_2$  are orthogonal projectors, respectively, that is,  $P_i^2 = P_i$ ,  $P_i' = P_i$ ,  $i = 1, 2$ .
- ▶  $Q = AA^+ = A(A'A)^+A'$  is the (unique) orthogonal projector on  $\text{Im}(A)$ .
- ▶  $A^+A = A'(A'A)^+A$  is the (unique) orthogonal projector on  $\text{Im}(A')$ .
- ▶  $\text{Ker}(A) = \text{Im}(P_1)$ ,  $\text{Im}(A) = \text{Ker}(P_2)$ .
- ▶  $\text{Ker}(A^+) = \text{Im}(P_2)$ ,  $\text{Im}(A^+) = \text{Ker}(P_1)$
- ▶  $\text{Im}(A) = \text{Ker}(A^+)^{\perp}$ ,  $\text{Im}(A^+) = \text{Ker}(A)^{\perp}$ ,
- ▶  $\text{Ker}(A) \oplus \text{Im}(A^+) = \mathbb{R}^q$ ,  $\text{Ker}(A^+) \oplus \text{Im}(A) = \mathbb{R}^p$

# Part I: Linear Models

## Chapter I.1

### Preliminaries

#### Probability

# Expectations of random vectors and random matrices

## 1.1.11 Definition (expectation of random vectors and random matrices)

- ① The **expectation of a random vector**  $\mathbf{X} = (X_1, \dots, X_p)'$  is defined by the vector of means, that is,

$$E\mathbf{X} = \begin{pmatrix} EX_1 \\ \vdots \\ EX_p \end{pmatrix};$$

subsequently, we use the notation  $\boldsymbol{\mu} = E\mathbf{X}$ ;

- ② The **expectation of a random matrix**  $\mathcal{X} = (X_{ij})_{1 \leq i \leq p, 1 \leq j \leq q}$  is defined by the matrix of means, that is,

$$E\mathcal{X} = \begin{pmatrix} EX_{11} & \cdots & EX_{1q} \\ \vdots & \ddots & \vdots \\ EX_{p1} & \cdots & EX_{pq} \end{pmatrix}.$$



In the following, all expectations are supposed to exist.

### ► I.1.12 Lemma

- ① Let  $\mathbf{X} = (X_1, \dots, X_p)'$  be a  $p$ -dimensional random vector and  $A \in \mathbb{R}^{k \times p}$ ,  $\mathbf{b} \in \mathbb{R}^k$ . Then:

$$E(A\mathbf{X} + \mathbf{b}) = AE(\mathbf{X}) + \mathbf{b}.$$

- ② Let  $Z_1, \dots, Z_n$  be  $p$ -dimensional random vectors and  $A_1, \dots, A_n \in \mathbb{R}^{k \times p}$ . Then:

$$E\left(\sum_{j=1}^n A_j Z_j\right) = \sum_{j=1}^n A_j E(Z_j) \in \mathbb{R}^k.$$

### ➤ I.1.13 Definition (variance-covariance matrix)

Let  $\mathbf{X} = (X_1, \dots, X_p)'$ ,  $\mathbf{Y} = (Y_1, \dots, Y_q)'$  be random vectors. Then, the **covariance matrix** of  $\mathbf{X}$  and  $\mathbf{Y}$  is defined by

$$\text{Cov}(\mathbf{X}, \mathbf{Y}) = \begin{pmatrix} \text{Cov}(X_1, Y_1) & \cdots & \text{Cov}(X_1, Y_q) \\ \vdots & \ddots & \vdots \\ \text{Cov}(X_p, Y_1) & \cdots & \text{Cov}(X_p, Y_q) \end{pmatrix}.$$

The **variance-covariance matrix** of  $\mathbf{X}$  is defined by  $\Sigma = \text{Cov}(\mathbf{X}) = \text{Cov}(\mathbf{X}, \mathbf{X})$ .

### ➤ I.1.14 Remark

Defining the random matrix  $\mathcal{C}_{\mathbf{X}, \mathbf{Y}} = (\mathbf{X} - E(\mathbf{X}))(\mathbf{Y} - E(\mathbf{Y}))'$ , we get

- ①  $\text{Cov}(\mathbf{X}, \mathbf{Y}) = E\mathcal{C}_{\mathbf{X}, \mathbf{Y}}$
- ②  $\text{Cov}(\mathbf{X}) = \text{Cov}(\mathbf{X}, \mathbf{X}) = E\mathcal{C}_{\mathbf{X}, \mathbf{X}}$
- ③  $\text{Cov}(\mathbf{X}, \mathbf{Y}) = E(\mathbf{X}\mathbf{Y}') - E\mathbf{X} \cdot E\mathbf{Y}'$
- ④  $\text{Cov}(\mathbf{X}) = E(\mathbf{X}\mathbf{X}') - E\mathbf{X} \cdot E\mathbf{X}'$

Covariance matrices are always non-negative definite, that is,  $\text{Cov}(\mathbf{X}) \geq 0$ .

### > I.1.15 Notation (block matrix)

A matrix  $A \in \mathbb{R}^{(p+q) \times (k+r)}$  can be written as a **block matrix**

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad \text{with } A_{11} \in \mathbb{R}^{p \times k}, A_{12} \in \mathbb{R}^{p \times r}, A_{21} \in \mathbb{R}^{q \times k}, A_{22} \in \mathbb{R}^{q \times r}.$$

### > I.1.16 Lemma

With the notation from Definition I.1.13, we get for  $A \in \mathbb{R}^{k \times p}$ ,  $B \in \mathbb{R}^{r \times q}$ ,  $\mathbf{b} \in \mathbb{R}^k$ ,  $\mathbf{c} \in \mathbb{R}^r$ :

- ①  $\text{Cov}(AX + \mathbf{b}, BY + \mathbf{c}) = A \text{Cov}(X, Y) B'$ ,
- ②  $\text{Cov}(AX + \mathbf{b}) = A \text{Cov}(X) A'$ ,
- ③  $\text{Cov} \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{bmatrix} \text{Cov}(X) & \text{Cov}(X, Y) \\ \text{Cov}(Y, X) & \text{Cov}(Y) \end{bmatrix}$ ,
- ④  $\text{Cov}(X, Y) = \text{Cov}(Y, X)'$ .

Using Lemma I.1.16, we can write with  $\Sigma_{XY} = \text{Cov}(X, Y)$ :

$$\Sigma_{\begin{pmatrix} X \\ Y \end{pmatrix}} = \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma_{YX} & \Sigma_{YY} \end{bmatrix} \stackrel{(*)}{=} \begin{bmatrix} \Sigma_{XX} & \Sigma_{XY} \\ \Sigma'_{XY} & \Sigma_{YY} \end{bmatrix}.$$

# Probability distributions on $\mathbb{R}$

## ► I.1.17 Remark (density functions of distributions on $\mathbb{R}$ )

- Normal distribution  $N(\mu, \sigma^2)$ :

$$f(x) = \varphi_{\mu, \sigma^2}(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\}, \quad x \in \mathbb{R}$$

- $\chi^2$ -distribution  $\chi^2(p)$  with  $p \in \mathbb{N}$  degrees of freedom:

$$f(x) = \frac{1}{2^{p/2}\Gamma(p/2)} x^{p/2-1} e^{-x/2}, \quad x > 0$$

- t-distribution  $t(p)$  with  $p \in \mathbb{N}$  degrees of freedom:

$$f(x) = \frac{\Gamma(\frac{n+1}{2})}{\sqrt{p\pi}\Gamma(\frac{p}{2})} \left(1 + \frac{x^2}{p}\right)^{-(p+1)/2}, \quad x \in \mathbb{R}$$

- F-distribution  $F(p, q)$  with  $p \in \mathbb{N}$  numerator and  $q \in \mathbb{N}$  denominator degrees of freedom:

$$f(x) = \frac{\Gamma(\frac{p+q}{2})}{\Gamma(\frac{p}{2})\Gamma(\frac{q}{2})} \left(\frac{p}{q}\right)^{n/2} \frac{x^{p/2-1}}{(1 + \frac{p}{q}x)^{\frac{p+q}{2}}}, \quad x > 0$$

# Connections of probability distributions

## ➤ I.1.18 Notation

The notation  $X_1, \dots, X_k \stackrel{\text{iid}}{\sim} P$  means that the random variables  $X_1, \dots, X_k$  are independent and identically distributed (iid) with  $X_1 \sim P$ .

The same notation is used for samples of random vectors.

## ➤ I.1.19 Proposition

- ① Let  $X \sim N(0, 1)$  and  $\mu \in \mathbb{R}, \sigma > 0$ . Then,  $\mu + \sigma X \sim N(\mu, \sigma^2)$ .
- ② Let  $X_1, \dots, X_p \stackrel{\text{iid}}{\sim} N(0, 1)$ . Then,  $\sum_{i=1}^p X_i^2 \sim \chi^2(p)$ .
- ③ Let  $X \sim \chi^2(p)$  and  $Z \sim \chi^2(q)$  be independent random variables. Then,  $X + Z \sim \chi^2(p + q)$ .
- ④ Let  $X \sim N(0, 1)$  and  $Z \sim \chi^2(p)$  be independent random variables. Then,  $\frac{X}{\sqrt{\frac{1}{p}Z}} \sim t(p)$ .
- ⑤ Let  $X \sim \chi^2(p)$  and  $Z \sim \chi^2(q)$  be independent random variables. Then,  $\frac{\frac{1}{p}X}{\frac{1}{q}Z} \sim F(p, q)$ .

## non-central $\chi^2$ - and F-distribution

### ► I.1.20 Remark

- Given independent random variables  $X_1, \dots, X_p$  with  $X_i \sim N(\mu_i, 1)$ ,  $\mu_i \in \mathbb{R}$ ,  $1 \leq i \leq p$ , the distribution of

$$\sum_{i=1}^p X_i^2$$

is called non-central  $\chi^2$ -distribution  $\chi^2(p, \delta)$  with  $p \in \mathbb{N}$  degrees of freedom and non-centrality parameter  $\delta = \frac{1}{2} \sum_{i=1}^p \mu_i^2 \geq 0$ .

Clearly,  $\chi^2(p) = \chi^2(p, 0)$ .

- Let  $X \sim \chi^2(p, \delta)$  and  $Z \sim \chi^2(q)$  be independent random variables. Then,

$$\frac{\frac{1}{p}X}{\frac{1}{q}Z} \sim F(p, q, \delta),$$

that is, the ratio has a non-central F-distribution  $F(p, q, \delta)$  with  $p \in \mathbb{N}$  numerator and  $q \in \mathbb{N}$  denominator degrees of freedom and non-centrality parameter  $\delta \geq 0$ .

Clearly,  $F(p, q) = F(p, q, 0)$ .