

# EIGEN-6C4

## The latest combined global gravity field model including GOCE data up to degree and order 2190 of GFZ Potsdam and GRGS Toulouse

Ch. Förste<sup>1</sup>, S.L. Bruinsma<sup>2</sup>, O. Abrikosov<sup>1</sup>, J.-M. Lemoine<sup>2</sup>, T. Schaller<sup>3</sup>, H.-J. Götze<sup>3</sup>, J. Ebbing<sup>3</sup>, J.C. Marty<sup>2</sup>, F. Flechtner<sup>1</sup>, G. Balmino<sup>2</sup> and R. Biancale<sup>2</sup>

<sup>1</sup>GFZ Potsdam, Dept. Geodesy and Remote Sensing, Telegrafenberg, D-14473 Potsdam, Germany (e-mail:  
[foer@gfz-potsdam.de](mailto:foer@gfz-potsdam.de))

<sup>2</sup>CNES/GRGS, 18, avenue Edouard Belin, F-31055 Toulouse, France (e-mail: [sean.bruinsma@cnes.fr](mailto:sean.bruinsma@cnes.fr))

<sup>3</sup>Department of Geosciences, Kiel University, D-24118 Kiel, Germany (e-mail: [jebbing@geophysik.uni-kiel.de](mailto:jebbing@geophysik.uni-kiel.de))



# Introduction

- GFZ Potsdam and GRGS/CNES Toulouse have a long-time close cooperation in the field of global gravity field determination which presently focuses among others on:
    - GOCE gravity field determination (“Direct Approach”): behalf of ESA and in the framework of the GOCE High Level Processing Facility” (GOCE-HPF), the latest release is **GO\_CONS\_GCF\_2\_DIR\_R5** (Bruinsma et al. 2014)
    - Computation of high resolution global gravity field models incl. GOCE SGG data:  
**EIGEN\*-6C** (2011, Shako et al. 2013, max. d/o 1420)  
**EIGEN-6C2** (2012, Förste et al. 2012, max. d/o 1949)  
**EIGEN-6C3** (2013, Förste et al. 2013, max d/o 1949), was taken as basis for the new Canadian Height Reference System: **Canadian Gravimetric Geoid CGG2013**
  - Subject of this presentation: **EIGEN-6C4** (max d/o 2190), the latest release of the EIGEN-6C-Series, containing the complete SGG data of the GOCE-Mission
- \* EIGEN = “European Improved Gravity model of the Earth by New techniques”

# Data used for EIGEN-6C4

## LAGEOS-1/2 SLR data:

1985 – 2010 of GRGS release 2 normal equations to degree/order 30

## GRACE GPS-SST and K-band range-rate data:

Feb 2003 – Dec 2012 of GRGS release 3 normal equations to degree 175

## GOCE data:

- SGG data ( $T_{xx}$ ,  $T_{yy}$ ,  $T_{zz}$ ,  $T_{xz}$ ) from 01 November 2009 – 20 October 2013
- weighting per measurement (based on RMS of residual), cos-latitude weighting
- individual normal equations for each SGG component (4) up to degree/order 300
- application of a (120 – 8) s band-pass filter for all four SGG components

## Surface data:

DTU 2'x2' global gravity anomaly grid (Anderson, 2010)

= altimetry over the oceans, EGM2008 (Pavlis et al. 2012) over continents

Block diagonal solution for degrees 371-2190

The combination of the different satellite and surface parts is done by a band-limited combination of normal equations, as a function of their resolution and accuracy (c.f. Förste et al. 2008).

# Combination scheme of the normal equations for EIGEN-6C4

Accumulation of a full normal matrix up to d/o 370:

~200.000 parameters, ~ 250 GByte


contribution to the solution EIGEN-6C4:



eliminated beforehand (block matrix reduction):




Separate block diagonal solution:




# EIGEN-6C4 – Spectral behaviour

# Spectral evaluation – difference degree amplitudes to EGM2008 (1)



## Spectral evaluation – difference degree amplitudes to EGM2008 (2)



# EIGEN-6C4 – Evaluation

# GPS/Levelling test with EIGEN-6C4

**Comparison with geoid heights determined point-wise  
by GPS positioning and levelling:**

- Root mean square (cm) about mean of GPS-Levelling minus model-derived geoid heights (number of points in brackets).

## GPS/Leveling fit in cm up to d/o 2190

	EGM2008	EIGEN-6C4
<b>Europe (1234)</b>	<b>20.7</b>	<b>20.9</b>
<b>Germany (675)</b>	<b>4.3</b>	<b>4.8</b>
<b>Canada (1930)</b>	<b>12.6</b>	<b>12.4</b>
<b>USA (6169)</b>	<b>24.6</b>	<b>24.5</b>
<b>Australia (201)</b>	<b>21.5</b>	<b>21.1</b>
<b>Japan (816)</b>	<b>8.2</b>	<b>7.8</b>
<b>Brasil (672)</b>	<b>36.6</b>	<b>30.6</b>

### Sources/References for the used GPS/Lev data:

- **USA:** (Milbert, 1998)
- **Canada:** (M. Véronneau, personal communication 2003, Natural Resources Canada)
- **Europe/Germany:** (Ihde et al., 2002)
- **Australia:** (G. Johnston, Geoscience Australia and W. Featherstone, Curtin University of Technology, personal communication 2007)
- **Japan:** (Tokuro Kodama, Geospatial Information Authority of Japan, personal communication 2013)
- **Brazil:** Denizar Blitzkow and Ana Cristina Oliveira Cancoro de Matos, Centro de Estudos de Geodesia (CENEVIDEO), personal communication, the data belongs to the Brazilian Institute of Geography and Statistics (IBGE)

# GOCE and CHAMP Orbit adjustment tests

- Observations: Kinematic GOCE (Bock et al. 2011) and CHAMP (Prange et al. 2010) orbit positions
- Dynamic orbit computation
- **GOCE: 60 arcs** (01.11. – 31.12.2009), Arclength = **1.25 days**
- **CHAMP: 75 arcs** (01.02. – 01.05.2008), Arclength = **1.0 day**
- Parametrization for GOCE (this is similar for CHAMP):
  - Accelerometer **biases**: 2/rev for cross track / radial / along track
  - Accelerometer **scaling factor**: along track fixed (set to 1.0), 1/arc for cross track / radial

Rms values [cm] of the orbit fit residuals (mean values for the 75/60 arcs)

Gravity Field Model	GOCE Max d/o 180 x 180	CHAMP Max d/o 150 x 150
EGM2008 (Pavlis et al. 2012)	<b>2.79</b>	<b>2.68</b>
GIF48 (Ries et al. 2011)	<b>1.64</b>	<b>2.52</b>
GOCO03S (Mayer-Gürr et al. 2012)	<b>1.64</b>	<b>2.55</b>
GO_CONS_GCF_2_DIR_R3	<b>1.60</b>	<b>2.50</b>
GO_CONS_GCF_2_DIR_R4 (Bruinsma et al. 2013)	<b>1.58</b>	<b>2.50</b>
GO_CONS_GCF_2_TIM_R4	<b>1.52</b>	<b>8.84</b>
GO_CONS_GCF_2_TIM_R5 (Brockmann et al. 2014)	<b>1.52</b>	<b>7.18</b>
GO_CONS_GCF_2_DIR_R5 (Bruinsma et al. 2014)	<b>1.51</b>	<b>2.50</b>
EIGEN-6C3stat	<b>1.55</b>	<b>2.50</b>
EIGEN-6C4	<b>1.50</b>	<b>2.50</b>

- EIGEN-6C4 gives the best orbit fits for GOCE
- The best CHAMP orbit fits for EIGEN-6C4, EIGEN-6C3 and the three GOCE models of the Direct Approach
- Worse CHAMP orbit fit results for the timewise models due to the GOCE polar gaps
  - the TIM-models are not good for orbit computations in general

# EIGEN-6C4

## Comparison with GEOHALO Airborne Gravimetry

HALO – the German High Altitude and LOng Range Research Aircraft (HALO).




### GEOHALO –

Multidisciplinary geoscientific airborne mission on the HALO aircraft over Italy  
June 2012



GFZ's airborne gravimeter on HALO:  
CHEKAN - AM

# The GEOHALO flight tracks over Italy (June 2012)




GEOHALO flight characteristics

Flight altitude: 3500 m


Speed: 120 m/s (425 km/h)

Application of a low-pass filter:  
cut-off period 260s → ~15 km spatial resolution along track




Fit of the track „R01-R03“ to EGM2008

Diff. to model: Min/Max/RMS: -6.80/5.75/2.13 mgal



## Fit of the track „R01-R03“ to EIGEN-6C4

Diff. to model: Min/Max/RMS: -5.55/6.15/1.78 mgal



## Statistics (mGal) of all GEOHALO airborne gravimetry tracks:

Fit of the measured gravity at flight altitude

to EGM2008 and EIGEN-6C4

Profile	#obs	Track	Min	Max	RMS	Min	Max	RMS
		length[km]	-----	eigen-6c4	-----	-----	egm2008	-----
G04-G05	3001	360	-4.54	3.33	<b>1.54</b>	-3.00	3.98	1.32
G06-G07	2101	250	-7.32	3.09	1.83	-7.58	3.01	1.97
G08-G09	1081	130	-6.47	4.13	1.63	-6.53	4.10	1.59
G10-G11	1921	230	-2.35	3.99	1.14	-3.20	3.06	1.06
G12-G13	3421	410	-4.31	8.76	1.98	-5.09	8.15	1.98
G14-G15	1981	240	-6.71	5.98	2.65	-7.22	5.52	2.66
G15-Ven	2101	250	-6.08	5.76	2.03	-7.17	5.83	2.11
<b>R01-R03</b>	<b>8401</b>	<b>1010</b>	<b>-5.55</b>	<b>6.15</b>	<b>1.78</b>	<b>-6.80</b>	<b>5.75</b>	<b>2.13</b>
R05-R06	3901	468	-12.61	8.85	2.82	-14.78	7.63	2.72
R08-R09	3961	470	-6.78	7.61	2.69	-7.02	7.08	2.63
B01-B03	4201	504	-9.96	13.72	3.24	-11.43	11.09	3.39
B09-B10	5101	612	-7.17	11.82	2.82	-6.24	11.80	2.67
B12-B14a	5161	619	-11.81	11.32	2.96	-12.99	9.68	3.09
B12-B14b	1201	144	-8.23	8.94	3.71	-7.93	9.76	3.81
H07-H06	6661	800	-13.52	9.68	2.73	-14.84	8.38	2.77
H03-H02	4681	562	-9.82	8.66	2.48	-9.24	8.81	2.57
<b>Overall RMS (58876 points/7065 km): 2.50 2.55</b>								

## 'GEOHALO flight characteristics'

Flight altitude: 3500 m

Speed: 120 m/s (425 km/h)

Application of a low-pass filter:

cut-off period 260s → ~15 km spatial resolution along track

# EIGEN-6C4

## Application in 3D density modelling

# Application of EIGEN-6C4 in density modelling for the Earth's crust and mantle

## 3D-Density modelling over South America

### Example: Profile at 22.2° South over South America

#### Input Gravity Data:


- Terrestrial Gravimetry (black dots, Götze et al. 1988 - 2006)
- Satellite Altimetry (Anderson & Knudsen 1998))

#### Constraining Data:


- Magnetotelluric
- Seismic (refraction & reflexion)
- Earthquake hypocentres
- Heat flow data
- Geological information

#### Modelling Software:



- IGMAS+ (Schmidt & Götze Univ. Kiel together with Transinsight Dresden)




## Previous density model from terrestrial data (Schaller, 2013; Diploma thesis)




## Previous density model – Gravity calculated for EIGEN-6C4 data



## Modified density model fitted to EIGEN-6C4



## Comparison: Previous and modified density models




### Modifications:


- 1 Depth of the boundary between upper and lower oceanic mantle lithosphere
- 2 Depth of the boundary between oceanic mantle lithosphere and asthenosphere

- 3 Strong inhomogeneity in mantle lithosphere of the old model
- 4 Geometry of the upward bulge of the downgoing plate
- 5 Smoothing of the boundary between upper and middle continental crust
- 6 Geometry of the asthenospheric wedge (on some model sections the size changed significantly)

## Comparison: Previous and modified density models



→ The verification of these model modifications based on EIGEN-6C4 is subject for further investigations, mainly by using seismological data



### Modifications:

- 1 Depth of the boundary between upper and lower oceanic mantle lithosphere
- 2 Depth of the boundary between oceanic mantle lithosphere and asthenosphere

- 3 Strong inhomogeneity in mantle lithosphere of the old model
- 4 Geometry of the upward bulge of the downgoing plate
- 5 Smoothing of the boundary between upper and middle continental crust
- 6 Geometry of the asthenospheric wedge (on some model sections the size changed significantly)

## Conclusion/Summary

- **EIGEN-6C4** is a new Global Combined Gravity Field model to degree/order 2190
- This model has been inferred from the combination of LAGEOS, GRACE, GOCE and DTU ground data
- EIGEN-6C4 contains the complete SGG data of the GOCE mission.
- On the continents for wavelengths beyond spher. harm. degree 235, EIGEN-6C4 is basically a reconstruction of EGM2008
- The final version of EIGEN-6C4 is downloadable at the ICGEM data base at GFZ Potsdam

<http://icgem.gfz-potsdam.de>

# Thank you for your attention!


## Main characteristics/differences of EIGEN-6C, EIGEN-6C2, EIGEN-6C3stat and EIGEN-6C4

	EIGEN-6C (2011)	EIGEN-6C2 (2012)	EIGEN-6C3stat (2013)	EIGEN-6C4 (2014)
Max d/o	1420	1949	1949	2190
LAGEOS	GRGS 200301 - 200906	GRGS, 1985 - 2010	GRGS, 1985 - 2010	GRGS, 1985 - 2010
GRACE	GRGS RL02 200301 - 200906	GRGS RL02 200303 - 201012	GRGS RL02 (deg. 2 – 100) 200302 – 201101 + GFZ RL05 (deg. 55 – 180) 200310 - 201209	GRGS RL03 10 years 2003 – 2012
Max d/o GRACE	130	130	180	130
GOCE SGG data	200 days $T_{xx}$ $T_{yy}$ $T_{zz}$ out of 20091101 – 20100630	350 days $T_{xx}$ $T_{yy}$ $T_{zz}$ out of 20091101 – 20110419	nominal orbit altitude: 837 days $T_{xx}$ $T_{yy}$ $T_{zz}$ $T_{xz}$ out of 20091101 – 20120801 + lower orbit phases: 225 days $T_{xx}$ $T_{yy}$ $T_{zz}$ out of 20120901 – 20130524	nominal orbit altitude: 837 days $T_{xx}$ $T_{yy}$ $T_{zz}$ $T_{xz}$ out of 20091101 – 20120801 + lower orbit phases: 422 days $T_{xx}$ $T_{yy}$ $T_{zz}$ $T_{xz}$ out of 20120801 – 20131020
Max d/o GOCE	210	210	235	235
Terrestrial data	DTU Global gravity anomalies	DTU Global gravity anomalies  DTU Ocean geoid + EGM2008 geoid grid	DTU Global gravity anomalies  DTU Ocean geoid + EGM2008 geoid grid	DTU Global gravity anomalies  DTU Ocean geoid + EGM2008 geoid grid


# EGM-DIR-5 compared with EGM2008: spatial

Satellite-only minus EGM2008, Max degree=260

$\sigma = 18.9 \text{ cm}$




GOCE-DIR-R5 vs. EGM2008 max d/o 260  
 $\zeta, 0.75^\circ \times 0.75^\circ$



# EIGEN-6C4 compared to EGM2008: spatial


EGM2008 minus EIGEN-6C4, max. degree = 260 (as for the GOCE-only model)



# EIGEN-6C4 compared to EGM2008: spatial

EGM2008 minus EIGEN-6C4, max. degree = 2190 (full resolution)

$\sigma = 12.4 \text{ cm} \rightarrow \text{the same as for max degree 260}$



## References:

- Andersen, O. B. and Knudsen, P. (1998), Global marine gravity field from the ERS-1 and Geosat geodetic mission Altimetry J. Geophys. Res., 103/C4, pp 8129 – 8137
- Andersen O. B. P. Knudsen and P. Berry. (2009): DNSC08 mean sea surface and mean dynamic topography models, Journal of Geophysical Research, Vol. 114, c11001 12 pp., 2009, doi:10.1029/2008JC005179
- Andersen, O. B.,(2010): The DTU10 Gravity field and Mean sea surface (2010), Second international symposium of the gravity field of the Earth (IGFS2), Fairbanks, Alaska
- Bock H, Jäggi A, Meyer U, Visser P, van den Ijssel J, van Helleputte T, Heinze M and Hugentobler U (2011), GPS-derived orbits for the GOCE satellite, J Geod (2011) 85:807–818, DOI 10.1007/s00190-011-0484-9
- Bruinsma, S. L., J. M. Lemoine, R. Biancale, and N. Vales (2010), CNES/GRGS 10-day gravity field models (release 2) and their evaluation, Adv. Space Res., 45(4), 587–601, doi:10.1016/j.asr.2009.10.012.
- Bruinsma, S.-L., Förste,C., Abrikosov, O., Marty, J.-C., Rio, M.-H., Mulet, S., Bonvalot, S. (2013): The new ESA satellite-only gravity field model via the direct approach, Geoph. Res. Lett., DOI 10.1002/grl.50716
- Sean L. Bruinsma, Christoph Förste, Oleg Abrikosov, Jean-Michel Lemoine, Jean-Charles Marty, Sandrine Mulet, Marie-Helene Rio, and Sylvain Bonvalot (2014), ESA's satellite-only gravity field model via the direct approach based on all GOCE data, Geoph. Res. Lett., DOI 10.1002/2014GL062045
- Dahle, C., F. Flechtner, C. Gruber, R. König, G. Michalak, and K. H. Neumayer (2012), GFZ GRACE level-2 processing standards document for level-2 product release 0005, Sci. Tech. Rep., Data, 12/02, Potsdam, Germany
- Förste, C.; Schmidt, R.; Stubenvoll, R.; Flechtner, F.; Meyer, U.; König, R.; Neumayer, H.; Biancale, R.; Lemoine, J.-M.; Bruinsma, S.; Loyer, S.; Barthelmes, F.; Esselborn, S. (2008): The GeoForschungsZentrum Potsdam / Groupe de Recherche de Géodésie Spatiale satellite-only and combined gravity field models: EIGEN-GL04S1 and EIGEN-GL04C. Journal of Geodesy, 82, 6, 331-346, DOI 10.1007/s00190-007-0183-8
- Förste C, S. L. Bruinsma, R. Shako, O. Abrikosov, F. Flechtner, J.-C. Marty, J.-M. Lemoine, C. Dahle, H. Neumeyer, F. Barthelmes, R. Biancale, G. Balmino and R. König (2012), A new release of EIGEN-6: The latest combined global gravity field model including LAGEOS, GRACE and GOCE data from the collaboration of GFZ Potsdam and GRGS Toulouse, Geophysical Research Abstracts Vol. 14, EGU2012-2821-2, EGU General Assembly 2012
- Ch. Förste, F. Flechtner, Ch. Dahle, O. Abrikosov, H. Neumayer, Franz Barthelmes, R. König, S.L. Bruinsma, J.C. Marty, J.-M. Lemoine, R. Biancale
- EIGEN-6C3 - The latest Combined Global Gravity Field Model incl. GOCE data up to d/o 1949 of GFZ Potsdam and GRGS Toulouse, Abstracts, AGU 2013 Fall Meeting (San Francisco, USA 2013)
- Götze H.-J. and MIGRA-Group (1996), Group updates Gravity Database for central Andes, EOS Trans. Am. Geophys. Un., 1996, VOLUME 77, number 19, pages 181--181, DOI = 10.1029/96EO00121
- Mayer-Gürr, T. and the GOCO consortium (2012), The new combined satellite only model GOCO03s, paper presented at International Symposium on Gravity, Geoid and Height Systems GGHS 2012, Venice, Italy
- Prange, L., A. Jäggi, R. Dach, H. Bock, G. Beutler, L. Mervart; 2010: AIUB-CHAMP02S: The influence of GNSS model changes on gravity field recovery using spaceborne GPS. Advances in Space Research, vol.45(2), pp. 215-224, DOI:10.1016/j.asr.2009.09.020
- Ries, J.C., S. Bettadpur, S. Poole, and T. Richter (2011): Mean Background Gravity Fields for GRACE Processing, GRACE Science Team Meeting, Austin, TX, 8-10 August 2011
- Pavlis, N.K., Holmes, S.A., Kenyon, S.C., and Factor, J.K. (2012), The development and evaluation of the Earth Gravitational Model 2008 (EGM2008), Journal of Geoph. Res., Vol. 117, B04406, doi:10.1029/2011JB008916, 2012
- Schmidt, S. and Götze, H.-J. Bouguer and Isostatic Maps of the Central Andes (2006), in: The Andes, pages 559—562, Springer, doi=10.1007/978-3-540-48684-8\_28
- Shako R, Förste C, Abrikosov O, Bruinsma SL, Marty J - C, Lemoine J - M, Flechtner F, Neumayer KH and Dahle C. (2013), EIGEN-6C: A High-Resolution Global Gravity Combination Model Including GOCE Data, in F. Flechtner et al. (eds.), Observation of the System Earth from Space – CHAMP, GRACE, GOCE and future missions, Advanced Technologies in Earth Sciences, DOI: 10.1007/978-3-642-32135-1\_20, Springer-Verlag Berlin Heidelberg 2013