

# Paths of analysis\*

BMK3

Synthia

October 10, 2022

## 1 Analysis parameters

**Analysis type:** Automatic Retrosynthesis

**Rules:** none selected

**Filters:** Exclude Diastereoselective reactions, Tunnels, FGI, FGI with protections

**Max. paths returned:** 50

**Max. iterations:** 2000

**Commercial:**

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

**Published:**

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

**My Stockroom:**

1. Max. molecular weight - 1000 g/mol

**Reaction scoring formula:**  $TUNNEL\_COEF * FGI\_COEF * STEP * 20 + 1000 * (CONFLICT + NON\_SELECTIVITY + FILTERS + PROTECT)$

**Chemical scoring formula:**  $SMALLER^3, SMALLER^{1.5}$

**Min. search width:** 400

**Max. reactions per product:** 60

---

\*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

**Strategies:** none selected

**FGI Coeff:** 0

**Tunnels Coeff:** 0

**JSON Parameters:** {}

## 2 Paths

5 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

### 2.1 Path 1

**Score:** 56.25





Figure 1: Outline of path 1

#### 2.1.1 Synthesis of imides from anhydrides



**Substrates:**

1. Aniline - *available at Sigma-Aldrich*
2. 3a,4,7,7a-Tetrahydroisobenzofuran-1,3-dione - *available at Sigma-Aldrich*

**Products:**

1. 2-phenyl-3a,4,7,7a-tetrahydro-isoindole-1,3-dione - *ChemBridge Corporation*


**Typical conditions:** AcOH

**Protections:** none

**Reference:** [10.1080/00397910802474966](#) and [10.1021/ja9024676](#) (SI) and [10.1002/ejoc.201402202](#)

**Retrosynthesis ID:** 8178

### 2.1.2 Heck Reaction



**Substrates:**

1. 2-Benzyloxy-1-bromonaphthalene - [available at Sigma-Aldrich](#)
2. 2-phenyl-3a,4,7,7a-tetrahydro-isoindole-1,3-dione - [ChemBridge Corporation](#)

**Products:**

1. O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1ccccc1


**Typical conditions:** Pd (cat). Ligand e.g. TXPTS. Base. Temp

**Protections:** none

**Reference:** [10.1016/j.tetlet.2013.01.077](#) or [10.1039/C3GC40493E](#)  
[10.1021/ol0360288](#) or [10.1021/ol702755g](#) or [10.1055/s-0033-1340319](#) or [10.1016/j.tet.2004.10.049](#)

**Retrosynthesis ID:** 9170

### 2.1.3 Benzylic oxidation



**Substrates:**

1. O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1ccccc1

**Products:**

1. O=C(Oc1ccc2ccccc2c1C1=CCC2C(=O)N(c3ccccc3)C(=O)C2C1)c1ccccc1

**Typical conditions:** Oxidant eg. O<sub>2</sub> or K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> or HIO<sub>4</sub>.solvent

**Protections:** none

**Reference:** [10.1039/B404823G](#) and [10.1055/s-0036-1588429](#)  
and [10.1016/j.tetlet.2010.09.021](#) and [10.1002/chem.201604750](#) and  
[10.1016/j.apcata.2014.01.042](#) and [10.1039/c3nj00045a](#) and [10.1021/jacs.6b08305](#)

**Retrosynthesis ID:** 31019416

## 2.2 Path 2

**Score:** 64.06





Figure 2: Outline of path 2

### 2.2.1 Heck Reaction



#### Substrates:

1. Cyclohex-4-ene-1,2-dicarboxylic acid - *available at Sigma-Aldrich*
2. 2-Benzyloxy-1-bromonaphthalene - *available at Sigma-Aldrich*

#### Products:

1. O=C(O)C1CC=C(c2c(OCc3ccccc3)ccc3ccccc23)CC1C(=O)O


**Typical conditions:** Pd (cat). Ligand e.g. TXPTS. Base. Temp

**Protections:** none

**Reference:** [10.1016/j.tetlet.2013.01.077](https://doi.org/10.1016/j.tetlet.2013.01.077) or [10.1039/C3GC40493E](https://doi.org/10.1039/C3GC40493E)  
[10.1021/ol0360288](https://doi.org/10.1021/ol0360288) or [10.1021/ol702755g](https://doi.org/10.1021/ol702755g) or [10.1055/s-0033-1340319](https://doi.org/10.1055/s-0033-1340319) or  
[10.1016/j.tet.2004.10.049](https://doi.org/10.1016/j.tet.2004.10.049)

**Retrosynthesis ID:** 9170

### 2.2.2 Synthesis of cyclic anhydrides



#### Substrates:

1. O=C(O)C1CC=C(c2c(OCc3ccccc3)ccc3ccccc23)CC1C(=O)O

**Products:**

1. O=C1OC(=O)C2CC(c3c(OCc4ccccc4)ccc4ccccc34)=CCC12


**Typical conditions:** MgCl<sub>2</sub>.Boc<sub>2</sub>O.THF.40C

**Protections:** none

**Reference:** DOI: [10.1021/cs501237p](https://doi.org/10.1021/cs501237p)

**Retrosynthesis ID:** 7263

### 2.2.3 Synthesis of imides from anhydrides



**Substrates:**

1. Aniline - *available at Sigma-Aldrich*
2. O=C1OC(=O)C2CC(c3c(OCc4ccccc4)ccc4ccccc34)=CCC12

**Products:**

1. O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1ccccc1


**Typical conditions:** AcOH

**Protections:** none

**Reference:** [10.1080/00397910802474966](https://doi.org/10.1080/00397910802474966) and [10.1021/ja9024676](https://doi.org/10.1021/ja9024676) (SI) and [10.1002/ejoc.201402202](https://doi.org/10.1002/ejoc.201402202)

**Retrosynthesis ID:** 8178

### 2.2.4 Benzylic oxidation



**Substrates:**

1. O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1ccccc1

**Products:**

1. O=C(Oc1ccc2ccccc2c1C1=CCC2C(=O)N(c3ccccc3)C(=O)C2C1)c1ccccc1

**Typical conditions:** Oxidant eg. O<sub>2</sub> or K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> or HIO<sub>4</sub>.solvent

**Protections:** none

**Reference:** [10.1039/B404823G](#) and [10.1055/s-0036-1588429](#)  
and [10.1016/j.tetlet.2010.09.021](#) and [10.1002/chem.201604750](#) and  
[10.1016/j.apcata.2014.01.042](#) and [10.1039/c3nj00045a](#) and [10.1021/jacs.6b08305](#)

**Retrosynthesis ID:** 31019416

### 2.3 Path 3

**Score:** 64.06





Figure 3: Outline of path 3

### 2.3.1 Heck Reaction



#### Substrates:

- 2-(1,3-dioxo-1,3,3a,4,7,7a-hexahydro-isoindol-2-yl)-benzoic acid - *available at Sigma-Aldrich*
- 2-Benzyloxy-1-bromonaphthalene - *available at Sigma-Aldrich*

#### Products:

- O=C(O)c1ccccc1N1C(=O)C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C1=O


**Typical conditions:** Pd (cat). Ligand e.g. TXPTS. Base. Temp

**Protections:** none

**Reference:** [10.1016/j.tetlet.2013.01.077](#) or [10.1039/C3GC40493E](#)  
[10.1021/ol0360288](#) or [10.1021/ol702755g](#) or [10.1055/s-0033-1340319](#) or  
[10.1016/j.tet.2004.10.049](#)

**Retrosynthesis ID:** 9170

### 2.3.2 Schmidt Reaction



#### Substrates:

- hydrazoic acid
- O=C(O)c1ccccc1N1C(=O)C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C1=O



**Products:**

1. Nc1cccc1N1C(=O)C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C1=O


**Typical conditions:** azide.H+.40C

**Protections:** none

**Reference:** [10.1039/B505080D](#)

**Retrosynthesis ID:** 10953

**2.3.3 Hydredediazoniatio**



**Substrates:**

1. Nc1cccc1N1C(=O)C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C1=O

**Products:**

1. O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1cccc1


**Typical conditions:** 1) HCl.NaNO<sub>2</sub> 2) H<sub>3</sub>PO<sub>2</sub>

**Protections:** none

**Reference:** [10.1016/j.bmcl.2013.10.058](#) and [10.1021/jm0004906](#) and [10.1002/ejoc.200600030](#) and [10.1016/j.tet.2016.02.011](#)

**Retrosynthesis ID:** 9999756

### 2.3.4 Benzylic oxidation



**Substrates:**

- O=C1C2CC=C(c3c(OCc4ccccc4)ccc4ccccc34)CC2C(=O)N1c1ccccc1

**Products:**

- O=C(Oc1ccc2ccccc2c1C1=CCC2C(=O)N(c3ccccc3)C(=O)C2C1)c1ccccc1

**Typical conditions:** Oxidant eg. O<sub>2</sub> or K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> or HIO<sub>4</sub>.solvent

**Protections:** none

**Reference:** [10.1039/B404823G](#) and [10.1055/s-0036-1588429](#)  
and [10.1016/j.tetlet.2010.09.021](#) and [10.1002/chem.201604750](#) and  
[10.1016/j.apcata.2014.01.042](#) and [10.1039/c3nj00045a](#) and [10.1021/jacs.6b08305](#)

**Retrosynthesis ID:** 31019416

### 2.4 Path 4

**Score:** 70.00





Figure 4: Outline of path 4

### 2.4.1 Hydrodediazonation



#### Substrates:

1. 2-(2-aminophenyl)-2,3,3a,4,7,7a-hexahydro-1H-isoindole-1,3-dione - *available at Sigma-Aldrich*

#### Products:

1. 2-phenyl-3a,4,7,7a-tetrahydro-isoindole-1,3-dione - *ChemBridge Corporation*


**Typical conditions:** 1) HCl.NaNO<sub>2</sub> 2) H<sub>3</sub>PO<sub>2</sub>

**Protections:** none

**Reference:** [10.1016/j.bmcl.2013.10.058](https://doi.org/10.1016/j.bmcl.2013.10.058) and [10.1021/jm0004906](https://doi.org/10.1021/jm0004906) and [10.1002/ejoc.200600030](https://doi.org/10.1002/ejoc.200600030) and [10.1016/j.tet.2016.02.011](https://doi.org/10.1016/j.tet.2016.02.011)

**Retrosynthesis ID:** 9999756

### 2.4.2 Reaction of acyl chlorides with alcohols and phenols



#### Substrates:

1. Benzoyl chloride - *available at Sigma-Aldrich*
2. 1-Bromo-2-naphthol - *available at Sigma-Aldrich*

#### Products:

1. benzoic acid-(1-bromo-[2]naphthyl ester)


**Typical conditions:** base.DCM

**Protections:** none

**Reference:** [10.1016/j.bmcl.2012.03.021](#) AND [10.1021/ja026266i](#) (SI, hydroperoxides) AND [10.1016/j.tetasy.2004.07.044](#) AND [10.1021/jm1006929](#) (SI) AND [10.1016/j.tet.2011.05.017](#) AND [10.1016/j.tetasy.2012.09.002](#) AND [10.1021/ol016268s](#) (SI) AND [10.1021/jo801116n](#) AND [10.1021/jo00279a041](#) AND WO2013/64518 A1, 2013 (page 102)

**Retrosynthesis ID:** 28549

### 2.4.3 Heck Reaction



**Substrates:**

- benzoic acid-(1-bromo-[2]naphthyl ester)
- 2-phenyl-3a,4,7,7a-tetrahydro-isoindole-1,3-dione - [ChemBridgeCorporation](#)

**Products:**

- O=C(Oc1ccc2ccccc2c1C1=CCC2C(=O)N(c3ccccc3)C(=O)C2C1)c1ccccc1

**Typical conditions:** Pd (cat). Ligand e.g. TXPTS. Base. Temp

**Protections:** none

**Reference:** [10.1016/j.tetlet.2013.01.077](#) or [10.1039/C3GC40493E](#)  
[10.1021/ol0360288](#) or [10.1021/ol702755g](#) or [10.1055/s-0033-1340319](#) or  
[10.1016/j.tet.2004.10.049](#)

**Retrosynthesis ID:** 9170


## 2.5 Path 5

**Score:** 70.00



## Retrosynthesis ID: 31015963

## 2.5.2 Reaction of acyl chlorides with alcohols and phenols



**Substrates:**

1. 1-Iodo-2-naphthol - *available at Sigma-Aldrich*
2. Benzoyl chloride - *available at Sigma-Aldrich*

**Products:**

1. O=C(Oc1ccc2ccccc2c1I)c1ccccc1


**Typical conditions:** base.DCM

**Protections:** none

**Reference:** [10.1016/j.bmcl.2012.03.021](#) AND [10.1021/ja026266i](#) (SI, hydroperoxides) AND [10.1016/j.tetasy.2004.07.044](#) AND [10.1021/jm1006929](#) (SI) AND [10.1016/j.tet.2011.05.017](#) AND [10.1016/j.tetasy.2012.09.002](#) AND [10.1021/ol016268s](#) (SI) AND [10.1021/jo801116n](#) AND [10.1021/jo00279a041](#) AND WO2013/64518 A1, 2013 (page 102)

**Retrosynthesis ID:** 28549

## 2.5.3 Heck Reaction



**Substrates:**

1. O=C(Oc1ccc2ccccc2c1I)c1ccccc1

2. 2-phenyl-3a,4,7,7a-tetrahydro-isoindole-1,3-dione - *ChemBridge Corporation*

**Products:**

1. O=C(Oc1ccc2ccccc2c1C1=CCC2C(=O)N(c3ccccc3)C(=O)C2C1)c1ccccc1

**Typical conditions:** Pd (cat). Ligand e.g. TXPTS. Base. Temp

**Protections:** none

**Reference:** *10.1016/j.tetlet.2013.01.077* or *10.1002/9780470716076* or *10.1021/op050106k* or *10.1021/ol0360288* or *10.1021/ol702755g* or *10.1055/s-0033-1340319* or *10.1016/j.tet.2004.10.049*

**Retrosynthesis ID:** 9186