

Paths of analysis*

Y3A

Synthia

October 10, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: Exclude Diastereoselective reactions, Tunnels, FGI, FGI with protections

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 100000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

Strategies: none selected

FGI Coeff: 0

Tunnels Coeff: 0

JSON Parameters: {}

2 Paths

5 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 1000100.08





Figure 1: Outline of path 1

2.1.1 Alkylation of ketones



Substrates:

1. Allyl bromide - *available at Sigma-Aldrich*
2. p-chlor-benzoyl-dithioessigsaeure-methylester

Products:

1. a-allyl-a-(p-chlorobenzoyl)dithioacetate


Typical conditions: LDA or other base.THF.-78C

Protections: none

Reference: DOI: [10.1021/jo1019738](https://doi.org/10.1021/jo1019738) OR DOI: [10.1021/jm00114a016](https://doi.org/10.1021/jm00114a016)

Retrosynthesis ID: 1866

2.1.2 Hydroxymethylation of aryl chlorides with acetoxymethyltrifluoroborate



Substrates:

1. α -allyl- α -(p-chlorobenzoyl)dithioacetate
2. potassium (acetoxymethyl)trifluoroborate - *available at Sigma-Aldrich*

Products:

1. C=CCC(C(=O)c1ccc(CO)cc1)C(=S)SC


Typical conditions: Pd(dba)₃.RuPhos.Na₂CO₃.dioxane/H₂O.reflux

Protections: none

Reference: [10.1021/ol300149b](#)

Retrosynthesis ID: 10033559

2.1.3 Synthesis of nitriles from alcohols



Substrates:

1. C=CCC(C(=O)c1ccc(CO)cc1)C(=S)SC

Products:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC


Typical conditions: 1) TEMPO.DCM.rt. 2) NH₃.I₂ or NH₃.tBuOCl

Protections: none

Reference: [10.1055/s-0033-1338489](#)

Retrosynthesis ID: 50205

2.1.4 Reduction of ketones with NaBH₄



Substrates:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC

Products:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1


Typical conditions: NaBH₄.EtOH.0-20 C

Protections: none

Reference: [10.1016/j.ejmech.2020.112360](#) p. 3, 8 and [10.1016/j.ejmech.2010.10.012](#) p. 434, 436

Retrosynthesis ID: 50432

2.1.5 Condensation of ketones with dithioesters



Substrates:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1
2. Methyl p-tolyl ketone - [available at Sigma-Aldrich](#)

Products:

1. C=CCC(C(=S)CC(=O)c1ccc(C)cc1)C(O)c1ccc(C#N)cc1

Typical conditions: NaH.DMF

Protections:

Functional group SMARTS	Classification	Protecting groups
[#6][CH]([#6])[OH]	alcohols	Methoxymethyl Ether (MOM)
		2-Methoxyethoxymethyl Ether (MEM)
		Tetrahydropyranyl Ether (THP)
		Benzyl Ether (PMB)
		t-Butyldimethylsilyl Ether (TB-DMS)
		Methyl Ether

Reference: [10.1021/jo400599e](#) and [10.1002/ejoc.201301667](#)

Retrosynthesis ID: 9996413

2.2 Path 2

Score: 1000115.31





Figure 2: Outline of path 2

2.2.1 Synthesis of O-substituted N-substituted hydroxamic acids



Substrates:

1. 1-methyl-1,1-dithiooxalaldehyde
2. n-methoxymethylamine - *available at Sigma-Aldrich*

Products:

1. CON(C)C(=O)C(=S)SC


Typical conditions: DCC.DMAP or CDI.TEA.DCM

Protections: none

Reference: Patent: WO2007/67333A2, 2007 & [10.1016/j.bmcl.2008.09.100](#)

Retrosynthesis ID: 1152

2.2.2 Synthesis of ketones from Weinreb amides



Substrates:

1. Allyl bromide - *available at Sigma-Aldrich*
2. CON(C)C(=O)C(=S)SC

Products:

1. C=CCC(=O)C(=S)SC


Typical conditions: 1. RmgBr.THF 2. TFA.DCM

Protections: none

Reference: [10.1021/jm051185t](#) and [10.1021/ol101021v](#) (supporting info)

Retrosynthesis ID: 6837

2.2.3 Olefination of ketones followed by hydrolysis



Substrates:

1. triphenylphosphonium methoxymethylide
2. C=CCC(=O)C(=S)SC

Products:

1. C=CCC(C=O)C(=S)SC


Typical conditions: KHMDS.THF hydrolysis: pTsOH.water.acetone

Protections: none

Reference: [10.1002/anie.201811403](#) and [10.1002/anie.201809130](#) and [10.1002/anie.201705809](#) and [10.1002/anie.201409038](#) and [10.1021/ol3028994](#) (SI)

Retrosynthesis ID: 31014861

2.2.4 Grignard-Type Reaction



Substrates:

1. C=CCC(C=O)C(=S)SC
2. C7H4BrMgN

Products:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1


Typical conditions: Mg or Li.ether

Protections: none

Reference: [10.1055/s-0030-1260809](#) or [10.1021/jm061429p](#) or [10.1021/jo0621423](#) or [10.1021/ja00373a036](#) or [10.1016/S0040-4020\(01\)00457-4](#)

Retrosynthesis ID: 25123

2.2.5 Condensation of ketones with dithioesters



Substrates:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1
2. Methyl p-tolyl ketone - [available at Sigma-Aldrich](#)

Products:

1. C=CCC(C(=S)CC(=O)c1ccc(C)cc1)C(O)c1ccc(C#N)cc1

Typical conditions: NaH.DMF

Protections:

Functional group SMARTS	Classification	Protecting groups
[#6][CH]([#6])[OH]	alcohols	Methoxymethyl Ether (MOM)
		2-Methoxyethoxymethyl Ether (MEM)
		Tetrahydropyranyl Ether (THP)
		Benzyl Ether (PMB)
		t-Butyldimethylsilyl Ether (TB-DMS)
		Methyl Ether

Reference: [10.1021/jo400599e](#) and [10.1002/ejoc.201301667](#)

Retrosynthesis ID: 9996413

2.3 Path 3

Score: 1000139.14





Figure 3: Outline of path 3

2.3.1 Hydroxymethylation of aryl chlorides with acetoxymethyltrifluoroborate



Substrates:

1. p-chlor-benzoyl-dithioessigsaeure-methylester
2. potassium (acetoxymethyl)trifluoroborate - *available at Sigma-Aldrich*

Products:

1. CSC(=S)CC(=O)c1ccc(CO)cc1


Typical conditions: Pd(dba)₂.RuPhos.Na₂CO₃.dioxane/H₂O.reflux

Protections: none

Reference: [10.1021/ol300149b](#)

Retrosynthesis ID: 10033559

2.3.2 Aldol Condensation



Substrates:

1. Acrolein
2. CSC(=S)CC(=O)c1ccc(CO)cc1

Products:

1. C=CC=C(C(=O)c1ccc(CO)cc1)C(=S)SC


Typical conditions: NaOEt.base

Protections: none

Reference: [10.1080/00397911.2016.1206938](#)

Retrosynthesis ID: 10049

2.3.3 Synthesis of nitriles from alcohols



Substrates:

1. C=CC=C(C(=O)c1ccc(CO)cc1)C(=S)SC

Products:

1. C=CC=C(C(=O)c1ccc(C#N)cc1)C(=S)SC


Typical conditions: 1) TEMPO.DCM.rt. 2) NH₃.I₂ or NH₃.tBuOCl

Protections: none

Reference: [10.1055/s-0033-1338489](https://doi.org/10.1055/s-0033-1338489)

Retrosynthesis ID: 50205

2.3.4 Reduction of enones to saturated alcohols



Substrates:

1. C=CC=C(C(=O)c1ccc(C#N)cc1)C(=S)SC

Products:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1


Typical conditions: NaBH₄.transition.metal.salt.(eg.Pd(OAc)₂.or.CeCl₃)

Protections: none

Reference: [10.1080/00397910902788117](#) AND [10.1021/jo00235a009](#)
AND [10.1016/0040-4020\(95\)00125-R](#) AND [10.1021/ja01327a041](#) AND
[10.1021/jo00302a056](#) AND [10.1002/adsc.200900628](#)

Retrosynthesis ID: 15304

2.3.5 Condensation of ketones with dithioesters



Substrates:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1

2. Methyl p-tolyl ketone - *available at Sigma-Aldrich*

Products:

1. C=CCC(C(=S)CC(=O)c1ccc(C)cc1)C(O)c1ccc(C#N)cc1

Typical conditions: NaH.DMF

Protections:

Functional group SMARTS	Classification	Protecting groups
[#6][CH]([#6])[OH]	alcohols	Methoxymethyl Ether (MOM)
		2-Methoxyethoxymethyl Ether (MEM)
		Tetrahydropyranyl Ether (THP)
		Benzyl Ether (PMB)
		t-Butyldimethylsilyl Ether (TB-DMS)
		Methyl Ether

Reference: [10.1021/jo400599e](#) and [10.1002/ejoc.201301667](#)

Retrosynthesis ID: 9996413

2.4 Path 4

Score: 1000139.14





Figure 4: Outline of path 4

2.4.1 Hydroxymethylation of aryl chlorides with acetoxymethyltrifluoroborate



Substrates:

1. p-chlor-benzoyl-dithioessigsaeure-methylester
2. potassium (acetoxymethyl)trifluoroborate - *available at Sigma-Aldrich*

Products:

1. CSC(=S)CC(=O)c1ccc(CO)cc1


Typical conditions: Pd(dba)₂.RuPhos.Na₂CO₃.dioxane/H₂O.reflux

Protections: none

Reference: [10.1021/ol300149b](https://doi.org/10.1021/ol300149b)

Retrosynthesis ID: 10033559

2.4.2 Conversion of Alcohols into Nitriles



Substrates:

1. CSC(=S)CC(=O)c1ccc(CO)cc1

Products:

1. CSC(=S)CC(=O)c1ccc(C#N)cc1


Typical conditions: I₂.RT

Protections: none

Reference: DOI: [10.1021/jo0625352](https://doi.org/10.1021/jo0625352)

Retrosynthesis ID: 10973

2.4.3 Alkylation of ketones



Substrates:

1. Allyl bromide - *available at Sigma-Aldrich*
2. CSC(=S)CC(=O)c1ccc(C#N)cc1

Products:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC


Typical conditions: LDA or other base.THF.-78C

Protections: none

Reference: DOI: [10.1021/jo1019738](https://doi.org/10.1021/jo1019738) OR DOI: [10.1021/jm00114a016](https://doi.org/10.1021/jm00114a016)

Retrosynthesis ID: 1866

2.4.4 Reduction of ketones with NaBH4



Substrates:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC

Products:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1


Typical conditions: NaBH4.EtOH.0-20 C

Protections: none

Reference: [10.1016/j.ejmech.2020.112360](https://doi.org/10.1016/j.ejmech.2020.112360) p. 3, 8 and [10.1016/j.ejmech.2010.10.012](https://doi.org/10.1016/j.ejmech.2010.10.012) p. 434, 436

Retrosynthesis ID: 50432

2.4.5 Condensation of ketones with dithioesters



Substrates:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1
2. Methyl p-tolyl ketone - *available at Sigma-Aldrich*

Products:

1. C=CCC(C(=S)CC(=O)c1ccc(C)cc1)C(O)c1ccc(C#N)cc1

Typical conditions: NaH.DMF

Protections:

Functional group SMARTS	Classification	Protecting groups
[#6][CH]([#6])[OH]	alcohols	Methoxymethyl Ether (MOM)
		2-Methoxyethoxymethyl Ether (MEM)
		Tetrahydropyranyl Ether (THP)
		Benzyl Ether (PMB)
		t-Butyldimethylsilyl Ether (TB-DMS)
		Methyl Ether

Reference: [10.1021/jo400599e](#) and [10.1002/ejoc.201301667](#)

Retrosynthesis ID: 9996413

2.5 Path 5

Score: 1000151.35





Figure 5: Outline of path 5

2.5.1 Alkylation of ketones



Substrates:

1. Allyl bromide - *available at Sigma-Aldrich*
2. p-chlor-benzoyl-dithioessigsaeure-methylester

Products:

1. a-allyl-a-(p-chlorobenzoyl)dithioacetate


Typical conditions: LDA or other base.THF.-78C

Protections: none

Reference: DOI: [10.1021/jo1019738](https://doi.org/10.1021/jo1019738) OR DOI: [10.1021/jm00114a016](https://doi.org/10.1021/jm00114a016)

Retrosynthesis ID: 1866

2.5.2 Nucleophilic aromatic substitution



Substrates:

1. a-allyl-a-(p-chlorobenzoyl)dithioacetate

Products:

1. C=CCC(C(=O)c1ccc(N)cc1)C(=S)SC


Typical conditions: solvent. Heating or pressure

Protections: none

Reference: [10.1021/jm00040a009](#) or [10.1111/bph.12233](#) or [10.1246/cl.1987.1187](#)

Retrosynthesis ID: 5003

2.5.3 Preparation of diazonium salts



Substrates:

1. C=CCC(C(=O)c1ccc(N)cc1)C(=S)SC

Products:

1. C=CCC(C(=O)c1ccc([N+]#N)cc1)C(=S)SC


Typical conditions: HCl.NaNO₂.OC


Protections: none

Reference: WO2006134111A1 p.32 and [10.1016/j.jfluchem.2012.11.007](#) and [10.3987/COM-97-7918](#)

Retrosynthesis ID: 11170

2.5.4 Sandmeyer Reaction





Substrates:

1. C=CCC(C(=O)c1ccc([N+](#N)cc1)C(=S)SC
2. Cupricin - *available at Sigma-Aldrich*

Products:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC


Typical conditions: Cu salt.heat

Protections: none

Reference: [10.1021/cr60126a003](https://doi.org/10.1021/cr60126a003) and [10.1021/acs.accounts.7b00566](https://doi.org/10.1021/acs.accounts.7b00566)

Retrosynthesis ID: 11500

2.5.5 Reduction of ketones with NaBH4



Substrates:

1. C=CCC(C(=O)c1ccc(C#N)cc1)C(=S)SC

Products:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1


Typical conditions: NaBH4.EtOH.0-20 C

Protections: none

Reference: [10.1016/j.ejmech.2020.112360](https://doi.org/10.1016/j.ejmech.2020.112360) p. 3, 8 and [10.1016/j.ejmech.2010.10.012](https://doi.org/10.1016/j.ejmech.2010.10.012) p. 434, 436

Retrosynthesis ID: 50432

2.5.6 Condensation of ketones with dithioesters



Substrates:

1. C=CCC(C(=S)SC)C(O)c1ccc(C#N)cc1
2. Methyl p-tolyl ketone - *available at Sigma-Aldrich*

Products:

1. C=CCC(C(=S)CC(=O)c1ccc(C)cc1)C(O)c1ccc(C#N)cc1

Typical conditions: NaH.DMF

Protections:

Functional group SMARTS	Classification	Protecting groups
[#6][CH]([#6])[OH]	alcohols	Methoxymethyl Ether (MOM)
		2-Methoxyethoxymethyl Ether (MEM)
		Tetrahydropyranyl Ether (THP)
		Benzyl Ether (PMB)
		t-Butyldimethylsilyl Ether (TB-DMS)
		Methyl Ether

Reference: [10.1021/jo400599e](#) and [10.1002/ejoc.201301667](#)

Retrosynthesis ID: 9996413