

# Paths of analysis\*

Y6

Synthia

October 10, 2022

## 1 Analysis parameters

**Analysis type:** Automatic Retrosynthesis

**Rules:** none selected

**Filters:** Exclude Diastereoselective reactions, Tunnels, FGI, FGI with protections

**Max. paths returned:** 50

**Max. iterations:** 2000

**Commercial:**

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

**Published:**

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

**My Stockroom:**

1. Max. molecular weight - 1000 g/mol

**Reaction scoring formula:**  $\text{TUNNEL\_COEF} * \text{FGI\_COEF} * \text{STEP} * 20 + 1000 * (\text{CONFLICT} + \text{NON\_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

**Chemical scoring formula:**  $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

**Min. search width:** 400

**Max. reactions per product:** 60

---

\*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

**Strategies:** none selected

**FGI Coeff:** 0

**Tunnels Coeff:** 0

**JSON Parameters:** {}

## 2 Paths

5 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

### 2.1 Path 1

**Score:** 2250084.06





Figure 1: Outline of path 1

#### 2.1.1 Aldehyde Formation



**Substrates:**

1. 2-[2-(methoxycarbonyl)ethyl]-2-phenyl-1,3-dioxolane

**Products:**

1. 3-(2-phenyl-[1,3]dioxolan-2-yl)-propionaldehyde


**Typical conditions:** DIBAL.solvent e.g. DCM

**Protections:** none

**Reference:** [10.1039/C39940000483](#) and [10.1039/C3CC47867J](#) and [10.1021/jo00222a054](#) and [10.1021/ja9934908](#) and [10.1021/jo902426z](#)

**Retrosynthesis ID:** 28551

### 2.1.2 Tebbe Olefination



**Substrates:**

1. 3-(2-phenyl-[1,3]dioxolan-2-yl)-propionaldehyde

**Products:**

1. C=CCCC1(c2ccccc2)OCCO1


**Typical conditions:** Cp<sub>2</sub>TiCl<sub>2</sub>.AlMe<sub>3</sub>.toluene

**Protections:** none

**Reference:** [10.1016/j.tet.2007.03.015](#) and [10.1002/9780470638859.conrr617](#)

**Retrosynthesis ID:** 11714

### 2.1.3 Allylic oxidation to alcohol



**Substrates:**

1. C=CCCC1(c2ccccc2)OCCO1

**Products:**

1. C=CC(O)CC1(c2ccccc2)OCCO1


**Typical conditions:** ArCOOOH or t-BuOOOH

**Protections:** none

**Reference:** DOI: [10.1021/ja00458a072](https://doi.org/10.1021/ja00458a072) AND [10.1016/j.tetlet.2013.03.046](https://doi.org/10.1016/j.tetlet.2013.03.046) AND [10.1039/b612423b](https://doi.org/10.1039/b612423b)

**Retrosynthesis ID:** 7603

### 2.1.4 Swern Oxidation



**Substrates:**

1. C=CC(O)CC1(c2ccccc2)OCCO1

**Products:**

1. C=CC(=O)CC1(c2ccccc2)OCCO1


**Typical conditions:** oxalyl chloride.DMSO.DCM.NMe<sub>3</sub>.-40C

**Protections:** none

**Reference:** [10.1055/s-1990-27036](#)

**Retrosynthesis ID:** 11163

### 2.1.5 Alkenylation-Aldol reaction of enones and enoate esters



**Substrates:**

1. C=CC(=O)CC1(c2ccccc2)OCCO1
2. a,a,a-Trifluoro-p-tolualdehyde - *available at Sigma-Aldrich*
3. Bromoethylene - *available at Sigma-Aldrich*

**Products:**

1. C=CCC(C(=O)CC1(c2ccccc2)OCCO1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** 1.RCuLi.2.RCHO


**Protections:** none

**Reference:** [10.1016/S0040-4039\(01\)80891-1](#) AND [10.1016/S0040-4020\(01\)82115-3](#) AND [10.1021/jo2010186](#) AND [10.1021/jo101439h](#) AND [10.1021/ja906241w](#)

**Retrosynthesis ID:** 20547

### 2.1.6 Hydrolysis of ketals





**Substrates:**

1. C=CCC(C(=O)CC1(c2ccccc2)OCCO1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** H<sub>2</sub>O.HCl

**Protections:** none

**Reference:** [10.1021/jo0159035](#) and [10.1021/jo00194a003](#) and

**Retrosynthesis ID:** 31013139

### 2.1.7 Keto-enol Tautomerism



**Substrates:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** solvent

**Protections:** none

**Reference:** [10.1021/ja01065a003](#) AND [10.1021/jo8012385](#)

**Retrosynthesis ID:** 7781

### 2.1.8 Synthesis of Thioketones using Lawesson's Reagent



#### Substrates:

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1
2. 4-methoxyphenyl-dithiophosphonate

#### Products:

1. C=CCC(C(=S)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Typical conditions:** Lawesson's Reagent.neat.microwave

**Protections:** none

**Reference:** DOI: [10.1021/ol990629a](https://doi.org/10.1021/ol990629a)

**Retrosynthesis ID:** 10798

### 2.2 Path 2

Score: 2250084.06





Figure 2: Outline of path 2

### 2.2.1 Alkenylation-Aldol reaction of enones and enoate esters



#### Substrates:

1. a,a,a-Trifluoro-p-tolualdehyde - *available at Sigma-Aldrich*
2. Bromoethylene - *available at Sigma-Aldrich*
3. 3-Buten-2-one - *available at Sigma-Aldrich*

#### Products:

1. C=CCC(C(C)=O)C(O)c1ccc(C(F)(F)F)cc1


Typical conditions: 1.RCuLi.2.RCHO

Protections: none

Reference: [10.1016/S0040-4039\(01\)80891-1](#) AND [10.1016/S0040-4020\(01\)82115-3](#) AND [10.1021/jo2010186](#) AND [10.1021/jo101439h](#) AND [10.1021/ja906241w](#)

Retrosynthesis ID: 20547

### 2.2.2 Condensation of methyl ketones with esters



#### Substrates:

1. C=CCC(C(C)=O)C(O)c1ccc(C(F)(F)F)cc1
2. Methyl benzoate - *available at Sigma-Aldrich*

#### Products:



1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** NaOMe.MeOH

**Protections:** none

**Reference:** [10.1016/j.tetlet.2007.10.010](#) and [10.1016/j.tetlet.2013.09.025](#) and [10.1016/j.ejmech.2013.10.072](#) and [10.1002/ange.19921040631](#)

**Retrosynthesis ID:** 4792

### 2.2.3 Keto-enol Tautomerism



**Substrates:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** solvent


**Protections:** none

**Reference:** [10.1021/ja01065a003](#) AND [10.1021/jo8012385](#)

**Retrosynthesis ID:** 7781

### 2.2.4 Synthesis of Thioketones using Lawesson's Reagent





**Substrates:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1
2. 4-methoxyphenyl-dithiophosphonsaeureanhydrid

**Products:**

1. C=CCC(C(=S)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Typical conditions:** Lawesson's Reagent.neat.microwave

**Protections:** none

**Reference:** DOI: [10.1021/ol990629a](https://doi.org/10.1021/ol990629a)

**Retrosynthesis ID:** 10798

## 2.3 Path 3

Score: 2250115.31





Figure 3: Outline of path 3

### 2.3.1 Aldol-like condensation with nitriles



**Substrates:**

1. a,a,a-Trifluoro-p-tolualdehyde - *available at Sigma-Aldrich*
2. 4-Pentenitrile - *available at Sigma-Aldrich*

**Products:**

1. C=CCC(C#N)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** LDA.THF.cooling

**Protections:** none

**Reference:** [10.1039/B800634B](#) and [10.1002/anie.201302613](#) and [10.1021/jm701319c](#) and [10.1016/S0040-4020\(98\)00122-7](#) and [10.1021/jo025872t](#)

**Retrosynthesis ID:** 23727

### 2.3.2 Blaise Reaction



**Substrates:**

1. Phenacyl bromide - *available at Sigma-Aldrich*
2. C=CCC(C#N)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** Zn.TMSCl.THF then HCl

**Protections:** none

**Reference:** [10.1002/ejoc.201403402](#)

**Retrosynthesis ID:** 10000153

### 2.3.3 Keto-enol Tautomerism



**Substrates:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** solvent

**Protections:** none

**Reference:** [10.1021/ja01065a003](#) AND [10.1021/jo8012385](#)

**Retrosynthesis ID:** 7781

### 2.3.4 Synthesis of Thioketones using Lawesson's Reagent



**Substrates:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1
2. 4-methoxyphenyl-dithiophosphonsaeureanhydrid

**Products:**

1. C=CCC(C(=S)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Typical conditions:** Lawesson's Reagent.neat.microwave

**Protections:** none

**Reference:** DOI: [10.1021/ol990629a](https://doi.org/10.1021/ol990629a)

**Retrosynthesis ID:** 10798

## 2.4 Path 4

Score: 2250115.31





Figure 4: Outline of path 4

### 2.4.1 Condensation of esters with aldehydes



**Substrates:**

1. a,a,a-Trifluoro-p-tolualdehyde - *available at Sigma-Aldrich*
2. Methyl 4-pentenoate - *available at Sigma-Aldrich*

**Products:**

1. C=CCC(C(=O)OC)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** LDA.THF

**Protections:** none

**Reference:** [10.1016/j.bmcl.2005.02.066](https://doi.org/10.1016/j.bmcl.2005.02.066) and [10.3762/bjoc.9.175](https://doi.org/10.3762/bjoc.9.175) and [10.1021/ol1016178](https://doi.org/10.1021/ol1016178)

**Retrosynthesis ID:** 4788

#### 2.4.2 Condensation of methyl ketones with esters



**Substrates:**

1. C=CCC(C(=O)OC)C(O)c1ccc(C(F)(F)F)cc1
2. Acetophenone - [available at Sigma-Aldrich](#)

**Products:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** NaOMe.MeOH

**Protections:** none

**Reference:** [10.1016/j.tetlet.2007.10.010](https://doi.org/10.1016/j.tetlet.2007.10.010) and [10.1016/j.tetlet.2013.09.025](https://doi.org/10.1016/j.tetlet.2013.09.025) and [10.1016/j.ejmech.2013.10.072](https://doi.org/10.1016/j.ejmech.2013.10.072) and [10.1002/ange.19921040631](https://doi.org/10.1002/ange.19921040631)

**Retrosynthesis ID:** 4792

#### 2.4.3 Keto-enol Tautomerism



**Substrates:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** solvent

**Protections:** none

**Reference:** [10.1021/ja01065a003](#) AND [10.1021/jo8012385](#)

**Retrosynthesis ID:** 7781

**2.4.4 Synthesis of Thioketones using Lawesson's Reagent**



**Substrates:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1
2. 4-methoxyphenyl-dithiophosphonane

**Products:**

1. C=CCC(C(=S)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Typical conditions:** Lawesson's Reagent.neat.microwave

**Protections:** none

**Reference:** DOI: [10.1021/ol990629a](#)

**Retrosynthesis ID:** 10798

**2.5 Path 5**

**Score:** 2250125.08





Figure 5: Outline of path 5

### 2.5.1 Condensation of esters with aldehydes



#### Substrates:

1. a,a,a-Trifluoro-p-tolualdehyde - *available at Sigma-Aldrich*
2. Methyl 4-pentenoate - *available at Sigma-Aldrich*

#### Products:

1. C=CCCC(=O)OC(O)c1ccc(C(F)(F)F)cc1


Typical conditions: LDA.THF

Protections: none

Reference: [10.1016/j.bmcl.2005.02.066](https://doi.org/10.1016/j.bmcl.2005.02.066) and [10.3762/bjoc.9.175](https://doi.org/10.3762/bjoc.9.175) and [10.1021/ol1016178](https://doi.org/10.1021/ol1016178)

Retrosynthesis ID: 4788

### 2.5.2 Acid catalyzed transesterification



#### Substrates:



1. C=CCC(C(=O)OC)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC1C(=O)OC1c1ccc(C(F)(F)F)cc1


**Typical conditions:** H+

**Protections:** none

**Reference:** [10.1021/cr00020a004](#)

**Retrosynthesis ID:** 50438

**2.5.3 Ring opening of lactones with enolates**



**Substrates:**

1. C=CCC1C(=O)OC1c1ccc(C(F)(F)F)cc1

2. Acetophenone - [available at Sigma-Aldrich](#)

**Products:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** LiHMDS.THF

**Protections:** none

**Reference:** [10.1021/ol801493w](#) and [10.1021/ol403423r](#) and [10.1021/ja061938g](#) and [10.1021/ja036521e](#)

**Retrosynthesis ID:** 24105

### 2.5.4 Keto-enol Tautomerism



**Substrates:**

1. C=CCC(C(=O)CC(=O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Products:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1


**Typical conditions:** solvent

**Protections:** none

**Reference:** [10.1021/ja01065a003](#) AND [10.1021/jo8012385](#)

**Retrosynthesis ID:** 7781

### 2.5.5 Thionation of Carbonyl Compounds using PSCl3



**Substrates:**

1. C=CCC(C(=O)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1
2. Phosphorus thiochloride - [available at Sigma-Aldrich](#)

**Products:**

1. C=CCC(C(=S)/C=C(\O)c1ccccc1)C(O)c1ccc(C(F)(F)F)cc1

**Typical conditions:** NEt<sub>3</sub>.H<sub>2</sub>O.microwave.70-100C

**Protections:** none

**Reference:** DOI: [10.1021/jo7022069](https://doi.org/10.1021/jo7022069)

**Retrosynthesis ID:** 11555