

Paths of analysis*

OP_Exp

Synthia

February 13, 2024

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: Expert-Coded Rules

Published Reactions: none selected

Filters: Exclude Diastereoselective reactions, Tunnels, FGI, FGI with protections

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Shorter paths: no

Pathway linearity: COMBO

Protecting groups: BALANCED

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 100000 * (\text{FILTERS} + \text{CONFLICT} + \text{NON_SELECTIVITY}) + 40 * \text{PROTECT}$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

FGI Coeff: 1

Tunnels Coeff: 1

JSON Parameters: {}

2 Paths

1 path found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 136.30





Figure 1: Outline of path 1

2.1.1 Synthesis of lactones from hydroxyketones and Bestmann ylide



Substrates:

1. Methyl (triphenylphosphoranylidene)acetate - *available at Sigma-Aldrich*
2. 2-Allyl-2-hydroxycyclohexanone

Products:

1. C=CCC12CCCCC1=CC(=O)O2


Typical conditions: toluene.heat

Protections: none

Reference: [10.1021/ol503150c](#) and [10.1002/anie.200704959](#) and [10.1016/S0040-4039\(02\)02535-2](#) and [10.1016/j.tetasy.2007.07.027](#) and [10.1016/S0040-4020\(97\)10428-8](#) and [10.1002/anie.201407788](#) and [10.1002/hlca.201300203](#) and reagent prep: [10.15227/orgsyn.082.0140](#)

Retrosynthesis ID: 10030991

2.1.2 Addition of silanes to Michael acceptors followed by oxidation



Substrates:

1. DMPSCl - *available at Sigma-Aldrich*
2. C=CCC12CCCCC1=CC(=O)O2

Products:

1. C=CCC12CCCCC1(O)CC(=O)O2


Typical conditions: 1.nBuLi.2.CuCN.3.electrophile.4.H2O2

Protections: none

Reference: [10.1021/ja058370g](#) AND (Oxidation) [10.1021/jo9905672](#) or [10.1021/ol300832f](#)

Retrosynthesis ID: 20296

2.1.3 Deoxygenation of alcohols with silanes



Substrates:

1. C=CCC12CCCCC1(O)CC(=O)O2

Products:

1. C=CCC12CCCCC1CC(=O)O2


Typical conditions: Et₃SiH.Lewis.or.Bronsted.Acid

Protections: none

Reference: [10.1021/jo0158534](#) AND [10.1021/ol3020144](#)

Retrosynthesis ID: 8163

2.1.4 Alkylation of Esters



Substrates:

1. Allyl bromide - *available at Sigma-Aldrich*

2. C=CCC12CCCCC1CC(=O)O2

Products:

1. C=CCC1C(=O)OC2(CC=C)CCCCC12

Typical conditions: base e.g. BuLi.THF

Protections: none

Reference: [10.1021/ja065404r](#) and [10.1016/S0040-4020\(01\)88337-X](#) and [10.1021/ja058303m](#) and [10.1021/acs.orglett.9b03078](#) and [10.1016/S0040-4020\(01\)80336-7](#)

Retrosynthesis ID: 31017104