

Paths of analysis*

O1

Synthia

October 10, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: Exclude Diastereoselective reactions, Tunnels, FGI, FGI with protections

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

Strategies: none selected

FGI Coeff: 0

Tunnels Coeff: 0

JSON Parameters: {}

2 Paths

5 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 45.00





Figure 1: Outline of path 1

2.1.1 Suzuki coupling of alkyl-9-BBNs with vinyl bromides



Substrates:

- 1,2-Dibromoethene - *available at Sigma-Aldrich*
- 9-(3,3-pentamethyleneallyl)-9-borabicyclo[3.3.1]nonane

Products:

1. Br/C=C/CC=C1CCCCC1


Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/ja00183a048](#) and [10.1039/b707338k](#) and [10.1016/j.tet.2015.05.039](#) and [10.1021/jo991064z](#) and [10.1021/ol060290+](#) and [10.1246/bcsj.65.2863](#)

Retrosynthesis ID: 25174

2.1.2 Suzuki coupling of arylboronic acids with vinyl Bromides



Substrates:

- (4-Acetoxyphenyl)boronic acid - *available at Sigma-Aldrich*
- Br/C=C/CC=C1CCCCC1

Products:

- CC(=O)Oc1ccc(/C=C/CC=C2CCCCC2)cc1

Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/cr00039a007](https://doi.org/10.1021/cr00039a007) and [10.1007/3418_2012_32](https://doi.org/10.1007/3418_2012_32) and [10.1021/cr0505268](https://doi.org/10.1021/cr0505268) and [10.1016/j.jfluchem.2016.01.018](https://doi.org/10.1016/j.jfluchem.2016.01.018) and [10.1039/C3CS60197H](https://doi.org/10.1039/C3CS60197H)

Retrosynthesis ID: 24862

2.2 Path 2

Score: 45.00





Figure 2: Outline of path 2

2.2.1 Takai olefination



Substrates:

1. Iodoform - *available at Sigma-Aldrich*
2. 3-oxo-propionsaeure-methylester

Products:

1. methyl 4-iodo-3(e)-butenoate


Typical conditions: CrCl₂.THF

Protections: none

Reference: [10.1021/ja00283a046](https://doi.org/10.1021/ja00283a046) and [10.1021/ja00237a081](https://doi.org/10.1021/ja00237a081)

Retrosynthesis ID: 10497

2.2.2 Esters reduction with LAH



Substrates:

1. methyl 4-iodo-3(e)-butenoate

Products:

1. (e)-4-iodo-3-buten-1-ol


Typical conditions: LiAlH₄.THF.0-20 C

Protections: none

Reference: [10.1016/j.ejmech.2019.112011](https://doi.org/10.1016/j.ejmech.2019.112011) p. 5, 10 and [10.1016/j.ejmech.2020.112910](https://doi.org/10.1016/j.ejmech.2020.112910) p. 3, 7

Retrosynthesis ID: 9910006

2.2.3 Appel Reaction



Substrates:

1. (e)-4-iodo-3-buten-1-ol

Products:

1. C₄H₆BrI


Typical conditions: PPh₃.CBr₄

Protections: none

Reference: [10.1021/ja800574m](#) and [10.1016/j.tet.2012.05.010](#) and [10.1016/j.tet.2004.09.021](#) (experimental)

Retrosynthesis ID: 9990037

2.2.4 HWE/Wittig Olefination



Substrates:

1. Cyclohexanone - [available at Sigma-Aldrich](#)
2. C₄H₆BrI

Products:

1. I/C=C/CC=C1CCCCC1


Typical conditions: 1.PPh₃ or trialkylphosphite.2.base.aldehyde

Protections: none

Reference: [10.1002/anie.200705005](#) and [10.1021/ol052106a](#) and [10.1021/jo00075a064](#) and [10.1021/ol3027297](#)

Retrosynthesis ID: 24425

2.2.5 Suzuki coupling of arylboronic pinacol esters with vinyl iodides



Substrates:

1. 4-Acetoxyphenylboronic acid pinacol ester - *available at Sigma-Aldrich*
2. I/C=C/CC=C1CCCCC1

Products:

1. CC(=O)Oc1ccc(/C=C/CC=C2CCCCC2)cc1

Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/cr00039a007](#) and [10.1007/3418_2012_32](#) and [10.1021/cr0505268](#) and [10.1016/j.jfluchem.2016.01.018](#) and [10.1039/C3CS60197H](#)

Retrosynthesis ID: 5045

2.3 Path 3

Score: 45.00





Figure 3: Outline of path 3

2.3.1 Corey-Seebach



Substrates:

- 1,3-Dithiane - *available at Sigma-Aldrich*
- (2-bromo-ethylidene)-cyclohexane

Products:

- 3-cyclohexylidenepropanal


Typical conditions: 1.BuLi.TMEDA.2.TCCA

Protections: none

Reference: [10.1039/P19860000183](#) AND [10.1016/S0040-4020\(01\)85646-5](#) AND [10.1039/c5ob00638d](#) deprotection: [10.1016/j.tetlet.2006.06.131](#)

Retrosynthesis ID: 15272

2.3.2 Wittig-Schlosser olefination



2.4 Path 4

Score: 45.00





Figure 4: Outline of path 4

2.4.1 Shapiro reaction followed by alkyl bromide addition



Substrates:

1. Hexahydrobenzaldehyde - *available at Sigma-Aldrich*
2. (e)-3-bromo-1-tributylstannylpropene

Products:

1. CCCC[Sn](/C=C/CC=C1CCCCC1)(CCCC)CCCC


Typical conditions: 1.TsNH₂NH₂.2.Mes₂Mg.LiCl.THF.heating then alkyl bromide.cooling

Protections: none

Reference: [10.1016/S0040-4039\(00\)75263-4](#) and [10.1021/ol300652k](#) and [10.1021/jo015699l](#)

Retrosynthesis ID: 9990458

2.4.2 Vinylation of aryl chlorides with stannanes



Substrates:

1. 4-CHLOROPHENYL ACETATE - *available at Sigma-Aldrich*
2. CCCC[Sn](/C=C/CC=C1CCCCC1)(CCCC)CCCC

Products:

1. CC(=O)Oc1ccc(/C=C/CC=C2CCCCC2)cc1

Typical conditions: [Pd].catalyst.phosphine.CsF

Protections: none

Reference: US2004/167128 p.97 and [10.3184/174751913X13635315066265](#) and [10.1021/ol0495927](#) and [10.1002/\(SICI\)1521-3773\(19990816\)38:16<2411::AID-ANIE2411>3.0.CO;2-T](#) and [10.1021/ol0495927](#) and [10.1021/ja020012f](#)

Retrosynthesis ID: 32849

2.5 Path 5

Score: 45.00





Figure 5: Outline of path 5

2.5.1 Takai olefination



Substrates:

1. 4-[(tert-Butyldimethylsilyl)oxy]benzaldehyde - *available at Sigma-Aldrich*
2. Iodoform - *available at Sigma-Aldrich*

Products:

1. CC(C)(C)[Si](C)(C)Oc1ccc(/C=C/I)cc1


Typical conditions: CrCl₂.THF

Protections: none

Reference: [10.1021/ja00283a046](#) and [10.1021/ja00237a081](#)

Retrosynthesis ID: 10497

2.5.2 Deprotection of TBS aryl ethers



Substrates:

1. CC(C)(C)[Si](C)(C)Oc1ccc(/C=C/I)cc1

Products:

1. C8H7IO


Typical conditions: TBAF.THF

Protections: none

Reference: [10.1016/j.tet.2013.01.017](#) and [10.1016/j.tet.2004.04.042](#)

Retrosynthesis ID: 31011409

2.5.3 Suzuki coupling of alkyl-9-BBNs with vinyl iodides



Substrates:

1. 9-(3,3-pentamethyleneallyl)-9-borabicyclo3.3.1nonane
2. C₈H₇IO

Products:

1. Oc1ccc(/C=C/CC=C2CCCCC2)cc1


Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/jo015995y](#) and [10.1016/j.tetlet.2010.11.139](#) And [10.1021/ol0600741](#) and [10.1055/s-2002-32602](#) and [10.1002/anie.200501760](#)

Retrosynthesis ID: 25168

2.5.4 Cu(OTf)₂ catalyzed acylation of phenols



Substrates:

1. Oc1ccc(/C=C/CC=C2CCCCC2)cc1
2. Acetic anhydride - *available at Sigma-Aldrich*

Products:

1. CC(=O)Oc1ccc(/C=C/CC=C2CCCCC2)cc1

Typical conditions: Cu(II).triflate.DCM.RT

Protections: none

Reference: DOI: [10.1016/S0040-4020\(01\)01229-7](https://doi.org/10.1016/S0040-4020(01)01229-7)

Retrosynthesis ID: 11601