Paths of analysis*

Synthia

October 10, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: Exclude Diastereoselecitve reactions, Tunnels, FGI, FGI with protec-

tions

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol

2. Max. price - 1500 \$/g

Published:

- 1. Max. molecular weight 1000 g/mol
- 2. Popularity 5

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: TUNNEL_COEF*FGI_COEF*STEP*20+1000 000*(CONFLICT+NON SELECTIVITY+FILTERS+PROTECT)

Chemical scoring formula: SMALLER^ 3,SMALLER^ 1.5

Min. search width: 400

Max. reactions per product: 60

^{*}The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

Strategies: none selected

FGI Coeff: 0

Tunnels Coeff: 0

JSON Parameters: {}

2 Paths

5 paths found. Paths are sorted by score. Reactions are sorted in appearance order for each path.

2.1 Path 1

Score: 2250125.08





Figure 1: Outline of path 1

2.1.1 Appel Reaction



Substrates:

1. 2-phenylselanyl-ethanol

Products:

1. 2-bromaethylphenylselenid

Typical conditions: PPh3.CBr4

Protections: none

Reference: 10.1021/ja800574m and 10.1016/j.tet.2012.05.010 and

10.1016/j.tet.2004.09.021 (experimental)

Retrosynthesis ID: 9990037

2.1.2 Conjugated addition of cuprate-aldol sequence

Substrates:

1. 3-Buten-2-one - available at Sigma-Aldrich

2. 2-bromaethylphenylselenid

3. 3-Cyanobenzaldehyde - available at Sigma-Aldrich

Products:

1. CC(=O)C(CCC[Se]c1ccccc1)C(O)c1cccc(C#N)c1

Typical conditions: 1.RCuLi.2.RCHO

Protections: none

2.1.3 Condensation of methyl ketones with esters

Substrates:

- 1. Methyl p-toluate available at Sigma-Aldrich
- 2. CC(=O)C(CCC[Se]c1ccccc1)C(O)c1cccc(C#N)c1

Products:

1. Cc1ccc(C(=O)CC(=O)C(CCC[Se]c2cccc2)C(O)c2cccc(C#N)c2)cc1

Typical conditions: NaOMe.MeOH

Protections: none

Reference: 10.1016/j.tetlet.2007.10.010 and 10.1016/j.tetlet.2013.09.025 and 10.1016/j.ejmech.2013.10.072 and 10.1002/ange.19921040631

Retrosynthesis ID: 4792

2.1.4 Selenoxide Elimination

Substrates:

 $1. \ \ Cc1ccc(C(=O)CC(=O)C(CCC[Se]c2cccc2)C(O)c2cccc(C\#N)c2)cc1$

Products:

1. C=CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C#N)c1

Typical conditions: 1) O3 or H2O2 or NaIO4. low temperature. 2) pyridine

or Et3N

Protections: none

Reference: DOI: 10.1021/ja00852a019 or DOI: 10.1021/ja00258a056 or DOI: 10.1039/B716256A or DOI: 10.1055/s-1998-1970 or DOI: 10.1016/S0040-4039(00)76646-9

Retrosynthesis ID: 8381

2.1.5 Keto-enol Tautomerism

Substrates:

 $1. \ C{=}CCC(C({=}O)CC({=}O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Products:

1. $C=CCC(C(=O)/C=C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: solvent

Protections: none

Reference: 10.1021/ja01065a003 AND 10.1021/jo8012385

2.1.6 Synthesis of Thioketones using Lawesson's Reagent

Substrates:

- $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$
- 2. 4-methoxyphenyl-dithiophosphonsaeureanhydrid

Products:

 $1. \ C{=}CCC(C({=}S)/C{=}C(\backslash O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: Lawesson's Reagent.neat.microwave

Protections: none

Reference: DOI: 10.1021/ol990629a

Retrosynthesis ID: 10798

2.2 Path 2

Score: 2250164.14




Figure 2: Outline of path 2

2.2.1 Condensation of esters with aldehydes

Substrates:

1. Methyl 4-pentenoate - available at Sigma-Aldrich

2. 3-Cyanobenzaldehyde - available at Sigma-Aldrich

Products:

1. C=CCC(C(=O)OC)C(O)c1cccc(C#N)c1

Typical conditions: LDA.THF

Protections: none

Reference: 10.1016/j.bmcl.2005.02.066 and 10.3762/bjoc.9.175 and

10.1021/ol1016178

Retrosynthesis ID: 4788

2.2.2 Acid catalyzed transesterification

Substrates:

1. C=CCC(C(=O)OC)C(O)c1cccc(C#N)c1

Products:

1. C=CCC1C(=O)OC1c1cccc(C#N)c1

Typical conditions: H+

Protections: none

Reference: 10.1021/cr00020a004

Retrosynthesis ID: 50438

2.2.3 Ring opening of lactones with enolates

Substrates:

1. Methyl p-tolyl ketone - available at Sigma-Aldrich

2. C=CCC1C(=O)OC1c1cccc(C#N)c1

Products:

 $1. \ C = CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: LiHMDS.THF

Protections: none

Reference: 10.1021/ol801493w and 10.1021/ol403423r and 10.1021/ja061938g

and 10.1021/ja036521e

2.2.4 Keto-enol Tautomerism

Substrates:

1. C=CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C#N)c1

Products:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: solvent

Protections: none

Reference: 10.1021/ja01065a003 AND 10.1021/jo8012385

Retrosynthesis ID: 7781

2.2.5 Synthesis of Thioketones using Lawesson's Reagent

Substrates:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

 $2.\ \, 4\hbox{-methoxyphenyl-} dithiophosphons a eurean hydrid$

Products:

 $1. \ C{=}CCC(C({=}S)/C{=}C(\backslash O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: Lawesson's Reagent.neat.microwave

Protections: none

Reference: DOI: 10.1021/ol990629a

Retrosynthesis ID: 10798

2.3 Path 3

Score: 2250164.14




Figure 3: Outline of path 3

2.3.1 Aldol Condensation

Substrates:

- 1. 3-Cyanobenzaldehyde available at Sigma-Aldrich
- 2. 1-diazo-hex-5-en-2-one

Products:

1. C=CCC(=Cc1cccc(C#N)c1)C(=O)C=[N+]=[N-]

Typical conditions: NaOEt.base

Protections: none

Reference: 10.1080/00397911.2016.1206938

Retrosynthesis ID: 10049

2.3.2 Homologation of aldehydes to ketones with diazoalkanes

Substrates:

1. p-Tolualdehyde - available at Sigma-Aldrich

2. C=CCC(=Cc1cccc(C#N)c1)C(=O)C=[N+]=[N-]

Products:

 $1. \ C = CCC(=Cc1cccc(C\#N)c1)C(=O)CC(=O)c1ccc(C)cc1$

Typical conditions: Lewis.acid

Protections: none

Reference: 10.1021/jo00275a006 AND 10.1016/j.tet.2014.05.107 AND

10.1016/j.tet.2014.11.059 AND 10.1021/ol9010932

2.3.3 Addition of silanes to Michael acceptors followed by oxidation

Substrates:

- $1. \ \ DMPSCl- \quad \ \ available \ at \ Sigma-Aldrich$
- $2. \ C = CCC(=Cc1cccc(C\#N)c1)C(=O)CC(=O)c1ccc(C)cc1$

Products:

 $1. \ C = CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: 1.nBuLi.2.CuCN.3.electrophile.4.H2O2

Protections: none

Reference: 10.1021/ja058370g AND (Oxidation) 10.1021/jo9905672 or 10.1021/ol300832f

Retrosynthesis ID: 20301

2.3.4 Keto-enol Tautomerism

Substrates:

 $1. \ C = CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Products:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: solvent

Protections: none

Reference: 10.1021/ja01065a003 AND 10.1021/jo8012385

Retrosynthesis ID: 7781

2.3.5Synthesis of Thioketones using Lawesson's Reagent

Substrates:

- 1. $C=CCC(C(=O)/C=C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$
- 2. 4-methoxyphenyl-dithiophosphonsaeureanhydrid

Products:

 $1. \ C = CCC(C(=S)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: Lawesson's Reagent.neat.microwave

Protections: none

Reference: DOI: 10.1021/ol990629a

Retrosynthesis ID: 10798

Path 4 2.4

Score: 2250164.14




Figure 4: Outline of path 4

2.4.1 Aldol Condensation

Substrates:

1. Allylacetone - available at Sigma-Aldrich

2. 3-Cyanobenzaldehyde - available at Sigma-Aldrich

Products:

1. C=CCC(=Cc1cccc(C#N)c1)C(C)=O

 ${\bf Typical\ conditions:}\ {\bf NaOEt.base}$

 ${\bf Protections:}\ {\rm none}$

Reference: 10.1080/00397911.2016.1206938

2.4.2 Addition of silanes to Michael acceptors followed by oxidation

Substrates:

- 1. DMPSCl available at Sigma-Aldrich
- 2. C=CCC(=Cc1cccc(C#N)c1)C(C)=O

Products:

1. C=CCC(C(C)=O)C(O)c1cccc(C#N)c1


Typical conditions: 1.nBuLi.2.CuCN.3.electrophile.4.H2O2

Protections: none

Reference: 10.1021/ja058370g AND (Oxidation) 10.1021/jo9905672 or 10.1021/ol300832f

Retrosynthesis ID: 20301

2.4.3 Condensation of methyl ketones with esters



Substrates:

- 1. C=CCC(C(C)=O)C(O)c1cccc(C#N)c1
- 2. Methyl p-toluate available at Sigma-Aldrich

Products:

 $1. \ C = CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

 ${\bf Typical\ conditions:}\ {\rm NaOMe.MeOH}$

Protections: none

Reference: 10.1016/j.tetlet.2007.10.010 and 10.1016/j.tetlet.2013.09.025 and

10.1016/j.ejmech.2013.10.072 and 10.1002/ange.19921040631

Retrosynthesis ID: 4792

2.4.4 Keto-enol Tautomerism

Substrates:

 $1. \ C = CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Products:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: solvent

Protections: none

Reference: 10.1021/ja01065a003 AND 10.1021/jo8012385

2.4.5 Synthesis of Thioketones using Lawesson's Reagent

Substrates:

- $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$
- 2. 4-methoxyphenyl-dithiophosphonsaeureanhydrid

Products:

 $1. \ C{=}CCC(C({=}S)/C{=}C(\backslash O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: Lawesson's Reagent.neat.microwave

Protections: none

Reference: DOI: 10.1021/ol990629a

Retrosynthesis ID: 10798

2.5 Path 5

Score: 2250164.14




Figure 5: Outline of path 5

2.5.1 Alkylation of vinyl esters

Substrates:

1. (E)-Methyl 3-(3-cyanophenyl)acrylate - available at Sigma-Aldrich

2. Allyl iodide - available at Sigma-Aldrich

Products:

1. $C=CC/C(=C \cdot c1cccc(C#N)c1)C(=O)OC$

Typical conditions: LDA.THF

Protections: none

Reference: DOI: 10.1039/C39870001410

Retrosynthesis ID: 886

2.5.2 Condensation of methyl ketones with esters

Substrates:

1. $C=CC/C(=C \cdot c1cccc(C#N)c1)C(=O)OC$

2. Methyl p-tolyl ketone - available at Sigma-Aldrich

Products:

 $1. \ C = CC/C(=C \setminus c1cccc(C\#N)c1)C(=O)CC(=O)c1ccc(C)cc1$

Typical conditions: NaOMe.MeOH

Protections: none

Reference: 10.1016/j.tetlet.2007.10.010 and 10.1016/j.tetlet.2013.09.025 and

10.1016/j.ejmech.2013.10.072 and 10.1002/ange.19921040631

Retrosynthesis ID: 4792

2.5.3 Addition of silanes to Michael acceptors followed by oxidation

Substrates:

1. DMPSCl - available at Sigma-Aldrich

 $2. \ C=CC/C(=C \setminus c1cccc(C\#N)c1)C(=O)CC(=O)c1ccc(C)cc1$

Products:

1. C=CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C#N)c1

Typical conditions: 1.nBuLi.2.CuCN.3.electrophile.4.H2O2

Protections: none

 $\textbf{Reference:} \hspace{0.5cm} 10.1021/ja058370g \hspace{0.5cm} \text{AND} \hspace{0.5cm} \text{(Oxidation)} \hspace{0.5cm} 10.1021/jo9905672 \hspace{0.5cm} \text{or} \hspace{0.5cm} 10.1021/ja0905672 \hspace{0.$

10.1021/ol300832f

2.5.4 Keto-enol Tautomerism

Substrates:

1. C=CCC(C(=O)CC(=O)c1ccc(C)cc1)C(O)c1cccc(C#N)c1

Products:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

Typical conditions: solvent

Protections: none

Reference: 10.1021/ja01065a003 AND 10.1021/jo8012385

Retrosynthesis ID: 7781

2.5.5 Synthesis of Thioketones using Lawesson's Reagent

Substrates:

 $1. \ C = CCC(C(=O)/C = C(\setminus O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

 $2. \ \ 4\text{-methoxyphenyl-dithiophosphonsaeureanhydrid}$

Products:

 $1. \ C{=}CCC(C({=}S)/C{=}C(\backslash O)c1ccc(C)cc1)C(O)c1cccc(C\#N)c1$

 ${\bf Typical\ conditions:}\ Lawesson's\ Reagent.neat.microwave$

Protections: none

Reference: DOI: 10.1021/ol990629a