# Paths of analysis\*

# Synthia

October 10, 2022

# 1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: Exclude Diastereoselecitve reactions, Tunnels, FGI, FGI with protec-

tions

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol

2. Max. price - 1500 \$/g

#### Published:

- 1. Max. molecular weight 1000 g/mol
- 2. Popularity 5

## My Stockroom:

1. Max. molecular weight - 1000 g/mol

**Reaction scoring formula:** TUNNEL\_COEF\*FGI\_COEF\*STEP\*20+1000 000\*(CONFLICT+NON SELECTIVITY+FILTERS+PROTECT)

Chemical scoring formula: SMALLER^ 3,SMALLER^ 1.5

Min. search width: 400

Max. reactions per product: 60

<sup>\*</sup>The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection

with the rights afforded in the license agreement and for no other purpose.

Strategies: none selected

FGI Coeff: 0

Tunnels Coeff: 0

JSON Parameters: {}

# 2 Paths

5 paths found. Paths are sorted by score. Reactions are sorted in appearance order for each path.

## 2.1 Path 1

Score: 45.00




Figure 1: Outline of path 1

# 2.1.1 Ring-opening of epoxides or thiiranes with azides

$$+ N = N^{+} NH$$

$$+ S$$

$$?$$

#### Substrates:

- 1. hydrazoic acid
- 2. 7-Oxabicyclo[4.1.0]heptan-2-one available at Sigma-Aldrich

#### **Products:**

1. [N-]=[N+]=NC1C(=O)CCCC1O

 $\textbf{Typical conditions:}\ \ NaN3.NH4Cl.MeOH.H2O.65\ C$ 


Protections: none

**Reference:** 10.1021/jm400529f p. 4361, 4367 and 10.1021/ja003713q p. 1590,

1594

Retrosynthesis ID: 858

# 2.1.2 Dehydration of Beta Hydroxy Carbonyl Compounds



## Substrates:

1. [N-]=[N+]=NC1C(=O)CCCC1O

#### **Products:**

1. 2-azidocyclohex-2-enone

Typical conditions: TsOH

Protections: none

**Reference:** DOI:10.1002/anie.201204977 AND 10.1021/ol0627770

Retrosynthesis ID: 7731

# 2.2 Path 2

Score: 76.25




Figure 2: Outline of path 2

# 2.2.1 Addition of silanes to Michael acceptors followed by oxidation

### Substrates:

1. DMPSCl - available at Sigma-Aldrich

2. 2-chloro-cyclohex-2-enone

## **Products:**

 $1. \ \, \hbox{$2$-chloro-$3-hydroxy-cyclohexanone}$ 

 $\textbf{Typical conditions:}\ 1. nBuLi. 2. CuCN. 3. electrophile. 4. H2O2$ 

Protections: none

**Reference:** 10.1021/ja058370g AND (Oxidation) 10.1021/jo9905672 or

10.1021/ol300832f

Retrosynthesis ID: 20295

# 2.2.2 Nucleophilic substitution with azides



#### Substrates:

1. Potassium azide - available at Sigma-Aldrich

2. 2-chloro-3-hydroxy-cyclohexanone

## **Products:**

1. [N-]=[N+]=NC1C(=O)CCCC1O

 ${\bf Typical\ conditions:\ DMF.} heat$ 


Protections: none

**Reference:** 10.1016/j.tet.2013.11.027 and 10.1021/jo015632y and 10.3987/COM-

06-S(K)18

Retrosynthesis ID: 31011248

# 2.2.3 Dehydration of Beta Hydroxy Carbonyl Compounds



## Substrates:

1. [N-]=[N+]=NC1C(=O)CCCC1O

## **Products:**

1. 2-azidocyclohex-2-enone

Typical conditions: TsOH

Protections: none

Reference: DOI:10.1002/anie.201204977 AND 10.1021/ol0627770


# 2.3 Path 3

Score: 76.25



Figure 3: Outline of path 3

# 2.3.1 Oxidation of Chiral Alcohols



# Substrates:

1. C6H10O2

## **Products:**

 $1. \ \mathrm{O}{=}\mathrm{C1}\mathrm{CCC}[\mathrm{C@H}]\mathrm{2OC12}$ 

Typical conditions: CAN.NaBrO3.CH3CN or other oxidant e.g. TPAP or NaOCl

Protections: none

**Reference:** DOI: 10.1016/s0040-4039(00)86883-5 or 10.1021/ja00054a005 or

10.1016/S0040-4039(00)85677-4

# 2.3.2 Ring-opening of epoxides or thiiranes with azides

$$+ N = N^{\pm}NH$$

$$+ N = N^{\pm}NH$$

$$? + ?$$

## Substrates:

- $1. \ \mathrm{O}{=}\mathrm{C1}\mathrm{CCC}[\mathrm{C@H}]\mathrm{2OC12}$
- 2. hydrazoic acid

# Products:

1. [N-]=[N+]=NC1C(=O)CCC[C@H]1O

Typical conditions: NaN3.NH4Cl.MeOH.H2O.65 C


Protections: none

**Reference:** 10.1021/jm400529f p. 4361, 4367 and 10.1021/ja003713q p. 1590,

1594

Retrosynthesis ID: 858

# 2.3.3 Dehydration of beta-ketoalcohols



#### Substrates:

1. [N-]=[N+]=NC1C(=O)CCC[C@H]1O

## **Products:**

1. 2-azidocyclohex-2-enone

 $\textbf{Typical conditions:} \ 1. MsCl. NEt 3$ 

Protections: none

**Reference:** 10.1021/ol301090v and 10.1021/ja00521a062 and

10.1002/ejoc.201201636 (SI)

Retrosynthesis ID: 20813

# 2.4 Path 4

Score: 84.06





Figure 4: Outline of path 4

# 2.4.1 Pinacol Coupling Reaction



## Substrates:

- 1. 7-Oxabicyclo[4.1.0]heptan-2-one available at Sigma-Aldrich
- 2. Ethanal available at Sigma-Aldrich

# Products:

1. CC(O)C1(O)CCCC2OC21

 $\textbf{Typical conditions:} \ \, \text{Mg.NH4Cl.H2O or Mg.SmI2.TMSCl.THF.HMPA}$ 

Protections: none

**Reference:** 10.1021/jo982497p p. 3234, 3236 and 10.1021/ol0506258 p. 2366, SI

p. S12

Retrosynthesis ID: 10205

# 2.4.2 Ring-opening of epoxides or thiiranes with azides

#### Substrates:

1. hydrazoic acid

2. CC(O)C1(O)CCCC2OC21

## **Products:**

1. CC(O)C1(O)CCCC(O)C1N=[N+]=[N-]

Typical conditions: NaN3.NH4Cl.MeOH.H2O.65  $\,\mathrm{C}$ 

Protections: none

**Reference:** 10.1021/jm400529f p. 4361, 4367 and 10.1021/ja003713q p. 1590,

1594

Retrosynthesis ID: 858

# 2.4.3 Cleavage of 1,2-diols with NaIO4



#### Substrates:

1. CC(O)C1(O)CCCC(O)C1N=[N+]=[N-]

#### **Products:**

1. [N-]=[N+]=NC1C(=O)CCCC1O

Typical conditions: NaIO4.solvent


Protections: none

**Reference:** 10.1039/C5OB00238A and 10.1002/chem.201301371 and

10.1021/ol052106a

Retrosynthesis ID: 31017508

# 2.4.4 Dehydration of Beta Hydroxy Carbonyl Compounds



#### Substrates:

1. [N-]=[N+]=NC1C(=O)CCCC1O

#### **Products:**

1. 2-azidocyclohex-2-enone

Typical conditions: TsOH

Protections: none

**Reference:** DOI:10.1002/anie.201204977 AND 10.1021/ol0627770

# 2.5 Path 5

Score: 84.06




Figure 5: Outline of path 5

# 2.5.1 Addition of alcohols or phenols to Michael acceptors

# Substrates:

- 1. methoxymethanol
- 2. 2-Bromocyclohex-2-en-1-one available at Sigma-Aldrich

# Products:

1. COCOC1CCCC(=O)C1Br

Typical conditions: cat.Na.DMF

Protections: none

 $\textbf{Reference:} \ 10.1016/S0957\text{-}\cancel{4}166(97)00\cancel{4}79\text{-}5 \ \text{AND} \ 10.1016/S00\cancel{4}0\text{-}\cancel{4}020(98)00817\text{-}$ 

 $5~{\rm AND}~10.1021/np970346w~{\rm AND}~10.1021/ol049820x$ 

# 2.5.2 Hydrolysis of acetals

#### Substrates:

 $1. \ COCOC1CCCC(=O)C1Br$ 

## **Products:**

1. 2-bromo-3-hydroxycyclohexanone


 $\textbf{Typical conditions:} \ \ \textbf{HCl.THF} \ \ \textbf{ot} \ \ \textbf{TFA.DCM}$ 

Protections: none

**Reference:** 10.1002/ejoc.201301615 and 10.3762/bjoc.9.289 and 10.1080/00397919808004478 and 10.1016/S0040-4039(99)02200-5 and 10.1055/s-0030-1258583 and 10.1016/j.tet.2011.04.072

Retrosynthesis ID: 31013131

# 2.5.3 Nucleophilic substitution with azides



Substrates:

- 1. 2-bromo-3-hydroxycyclohexanone
- 2. Potassium azide available at Sigma-Aldrich

#### **Products:**

1. [N-]=[N+]=NC1C(=O)CCCC1O

Typical conditions: DMF.heat


Protections: none

**Reference:** 10.1021/ol049369+ and 10.1016/S0040-4039(00)61343-6 and

10.1016/j.bmcl.2005.03.055

Retrosynthesis ID: 31011250

# 2.5.4 Dehydration of Beta Hydroxy Carbonyl Compounds



## Substrates:

1. 
$$[N-]=[N+]=NC1C(=O)CCCC1O$$

## **Products:**

1. 2-azidocyclohex-2-enone

Typical conditions: TsOH

Protections: none

**Reference:** DOI:10.1002/anie.201204977 AND 10.1021/ol0627770