

Paths of analysis*

L11_DIA

Synthia

October 11, 2022

1 Analysis parameters

Analysis type: Automatic Retrosynthesis

Rules: none selected

Filters: Tunnels, FGI, FGI with protections

Max. paths returned: 50

Max. iterations: 2000

Commercial:

1. Max. molecular weight - 1000 g/mol
2. Max. price - 1500 \$/g

Published:

1. Max. molecular weight - 1000 g/mol
2. Popularity - 5

My Stockroom:

1. Max. molecular weight - 1000 g/mol

Reaction scoring formula: $\text{TUNNEL_COEF} * \text{FGI_COEF} * \text{STEP} * 20 + 1000000 * (\text{CONFLICT} + \text{NON_SELECTIVITY} + \text{FILTERS} + \text{PROTECT})$

Chemical scoring formula: $\text{SMALLER}^3, \text{SMALLER}^{1.5}$

Min. search width: 400

Max. reactions per product: 60

Strategies: none selected

*The results stated herein were generated using the proprietary platform owned and maintained by Grzybowski Scientific Inventions, Inc., a subsidiary of Merck KGaA, Darmstadt Germany. The results are provided on an as is basis, and shall be used solely in connection with the rights afforded in the license agreement and for no other purpose.

FGI Coeff: 0

Tunnels Coeff: 0

JSON Parameters: {}

2 Paths

2 paths found. *Paths are sorted by score. Reactions are sorted in appearance order for each path.*

2.1 Path 1

Score: 281.47





Figure 1: Outline of path 1

2.1.1 Oxidation of primary alcohols with DMP



1. 4-hydroxy-2-butenyl-benzoat

Products:

1. O=CC=CCOC(=O)c1ccccc1


Typical conditions: DMP.DCM.0-25 C

Protections: none

Reference: [10.1016/j.bmc.2020.115469](https://doi.org/10.1016/j.bmc.2020.115469) p. 3, 9 and [10.1021/acs.jmedchem.8b01878](https://doi.org/10.1021/acs.jmedchem.8b01878) SI p. S43

Retrosynthesis ID: 50426

2.1.2 Hydrolysis of benzoates



Substrates:

1. O=CC=CCOC(=O)c1ccccc1

Products:

1. 4-hydroxy-but-2-enal


Typical conditions: LiOH/K₂CO₃/NH₃.MeOH.H₂O.THF

Protections: none

Reference: [10.1021/jm0502788](https://doi.org/10.1021/jm0502788) and [10.1016/j.tetlet.2008.09.165](https://doi.org/10.1016/j.tetlet.2008.09.165) and [10.1021/jm034098e](https://doi.org/10.1021/jm034098e) and [10.1021/jo049277y](https://doi.org/10.1021/jo049277y) and [10.1055/s-0033-1338657](https://doi.org/10.1055/s-0033-1338657)

Retrosynthesis ID: 25136

2.1.3 Suzuki coupling of vinyl bromides with alkenyl boronic acids



Substrates:

1. (E)-2-(methoxycarbonyl)ethyl-1-enylboronic acid
2. 1-bromo-1,2-dichloro-ethene

Products:

1. COC(=O)C=CC(Cl)=CCl


Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/cr00039a007](#) and [10.1007/3418_2012_32](#) and [10.1021/cr0505268](#) and [10.1016/j.jfluchem.2016.01.018](#) and [10.1039/C3CS60197H](#)

Retrosynthesis ID: 24937

2.1.4 Acid catalyzed transesterification



Substrates:

1. 4-hydroxybut-2-enal
2. COC(=O)C=CC(Cl)=CCl

Products:

1. O=CC=CCOC(=O)C=CC(Cl)=CCl


Typical conditions: H⁺

Protections: none

Reference: [10.1021/cr00020a004](https://doi.org/10.1021/cr00020a004)

Retrosynthesis ID: 50438

2.1.5 Diels-Alder



Substrates:

1. O=CC=CCOC(=O)C=CC(Cl)=CCl

Products:

1. O=CC1C(Cl)C(Cl)=CC2C(=O)OCC21


Typical conditions: Lewis acid or chiral Lewis acid. Solvent.

Protections: none

Reference: DOI: [10.1002/1521-3773\(20020517\)41:10<1668::AID-ANIE1668>3.0.CO;2-Z](https://doi.org/10.1002/1521-3773(20020517)41:10<1668::AID-ANIE1668>3.0.CO;2-Z) AND [10.1021/ja062508t](https://doi.org/10.1021/ja062508t)

Retrosynthesis ID: 18116

2.1.6 Takai olefination



Substrates:

- 1,1-Diiodoethane - *available at Sigma-Aldrich*
- O=CC1C(Cl)C(Cl)=CC2C(=O)OCC21

Products:

- C/C=C/C1C(Cl)C(Cl)=CC2C(=O)OCC21


Typical conditions: CrCl₂.THF.DMF

Protections: none

Reference: [10.1021/ja00283a046](#) and [10.1021/ja00237a081](#)

Retrosynthesis ID: 10942

2.1.7 Synthesis of organotrifluoroborate salts in mild conditions



Substrates:

- 2-(p-Toluenesulfonylamino)phenylboronic acid pinacol ester - *available at Sigma-Aldrich*

Products:

- Cc1ccc(S(=O)(=O)Nc2ccccc2[B-](F)(F)F)cc1


Typical conditions: KF.tartaric acid.CH₃CN.THF.H₂O.r.t

Protections: none

Reference: DOI: [10.1002/anie.201203930](#)

Retrosynthesis ID: 6072

2.1.8 Alkylation of sulfonamides with alkyl chlorides



Substrates:

1. Cc1ccc(S(=O)(=O)Nc2ccccc2[B-](F)(F)F)cc1
2. C/C=C/C1C(Cl)C(Cl)=CC2C(=O)OCC21

Products:

1. C/C=C/C1C2COC(=O)C2C=C(Cl)C1N(c1ccccc1[B-](F)(F)F)S(=O)(=O)c1ccc(C)cc1


Typical conditions: LDA

Protections: none

Reference: [10.1002/1099-0690\(200101\)2001:2<323::AID-EJOC323>3.0.CO;2-A](#)

Retrosynthesis ID: 7727

2.1.9 Claisen Condensation



Substrates:

1. C/C=C/C1C2COC(=O)C2C=C(Cl)C1N(c1ccccc1[B-](F)(F)F)S(=O)(=O)c1ccc(C)cc1
2. Methyl acetate - *available at Sigma-Aldrich*

Products:

1. C/C=C/C1C(N(c2ccccc2[B-](F)(F)F)S(=O)(=O)c2ccc(C)cc2)C(Cl)=CC2(C(C)=O)C(=O)OCC12


Typical conditions: Base.Solvent

Protections: none

Reference: [10.1021/cr020703u](#) and [10.1021/cr60088a002](#)

Retrosynthesis ID: 5015

2.1.10 Suzuki Coupling of aryltrifluoroborates with alkenyl chlorides



Substrates:

1. C/C=C/C1C2C(=CC3(C(C)=O)C(=O)OCC13)c1ccccc1N2S(=O)(=O)c1ccc(C)cc1

Products:

1. C/C=C/C1C2C(=CC3(C(C)=O)C(=O)OCC13)c1ccccc1N2S(=O)(=O)c1ccc(C)cc1

Typical conditions: PdCl₂(dppf).K₂CO₃.toluene.reflux

Protections: none

Reference: [10.1002/chem.200900425](#) and [10.1016/j.tetlet.2014.10.078](#)

Retrosynthesis ID: 10034105

2.2 Path 2

Score: 322.94





Figure 2: Outline of path 2

2.2.1 Appel Reaction



Substrates:

1. 2,3-Dibromoallyl alcohol - *available at Sigma-Aldrich*

Products:

1. 1,2,3-tribrom-propen


Typical conditions: PPh₃.CBr₄

Protections: none

Reference: [10.1021/ja800574m](#) and [10.1016/j.tet.2012.05.010](#) and [10.1016/j.tet.2004.09.021](#) (experimental)

Retrosynthesis ID: 9990037

2.2.2 Alkylation of 1,3 activated CH acids



Substrates:

1. 3-Oxobutanenitrile - *available at Sigma-Aldrich*
2. 1,2,3-tribrom-propen

Products:

1. CC(=O)C(C#N)CC(Br)=CBr


Typical conditions: base e.g. NaH.DMF

Protections: none

Reference: [10.1002/ejic.201403224](#) and [10.1016/j.bmcl.2005.11.008](#) and [10.1021/ja058303m](#) and [10.1021/acs.orglett.9b03078](#) and [10.1016/S0040-4020\(01\)80336-7](#)

Retrosynthesis ID: 31017076

2.2.3 Enol esters and ethers synthesis



Substrates:

1. TMSCl - *available at Sigma-Aldrich*
2. CC(=O)C(C#N)CC(Br)=CBr

Products:

1. CC(O[Si](C)(C)C)=C(C#N)CC(Br)=CBr


Typical conditions: 1. Et3N.Electrophile

Protections: none

Reference: [10.1016/S0040-4020\(03\)00977-3](#) AND [10.1021/ja00056a002](#)

Retrosynthesis ID: 7799

2.2.4 Dehydrogenation of silyl enol ethers



Substrates:

1. CC(O[Si](C)(C)C)=C(C#N)CC(Br)=CBr

Products:

1. CC(=O)C(C#N)=CC(Br)=CBr


Typical conditions: Pd(OAc)₂.Cu(OAc)₂.O₂.MeCN

Protections: none

Reference: [10.1271/bbb.60.405](#) and [10.1039/C3CC46778C](#) and US2015284405 p.40 and [10.1016/S0040-4039\(01\)81518-5](#) and US2010204477 p. 15-16 and [10.1016/0040-4039\(95\)00694-8](#) and [10.1021/jo00089a034](#) and [10.1016/S0040-4020\(01\)90587-3](#) and [10.1080/00397919008052802](#) and [10.1021/ja00218a060](#)

Retrosynthesis ID: 9999877

2.2.5 Acid hydrolysis of nitriles to carboxylic acids



Substrates:

1. CC(=O)C(C#N)=CC(Br)=CBr

Products:

1. CC(=O)C(=CC(Br)=CBr)C(=O)O


Typical conditions: Hcl.heating.H2O

Protections: none

Reference: [10.1021/jm301796k](#) and [10.1016/j.bmcl.2007.06.054](#) and [10.1021/jm801532e](#)

Retrosynthesis ID: 16027

2.2.6 Opening of epoxides with carboxylic acids



Substrates:

1. ethynyl-oxirane
2. CC(=O)C(=CC(Br)=CBr)C(=O)O

Products:

1. C#CC(O)COC(=O)C(=CC(Br)=CBr)C(C)=O


Typical conditions: RCOOH.catalyst


Protections: none

Reference: [10.1021/ol051051+](#) AND [10.1016/j.tet.2005.05.050](#)
and US2011/86912 A1 (P.13) and [10.1055/s-2003-42416](#) and [10.5012/bkcs.2013.34.8.2286](#)

Retrosynthesis ID: 15151

2.2.7 Meyer-Schuster Rearrangement





Substrates:

1. C#CC(O)COC(=O)C(=CC(Br)=CBr)C(C)=O

Products:

1. CC(=O)C(=CC(Br)=CBr)C(=O)OCC=CC=O


Typical conditions: H+

Protections: none

Reference: [10.1021/cr60273a001](https://doi.org/10.1021/cr60273a001)

Retrosynthesis ID: 10143

2.2.8 Diels-Alder



Substrates:

1. CC(=O)C(=CC(Br)=CBr)C(=O)OCC=CC=O

Products:

1. CC(=O)C12C=C(Br)C(Br)C(C=O)C1COC2=O


Typical conditions: Lewis acid or chiral Lewis acid. Solvent.

Protections: none

Reference: DOI: [10.1002/1521-3773\(20020517\)41:10<1668::AID-ANIE1668>3.0.CO;2-Z](https://doi.org/10.1002/1521-3773(20020517)41:10<1668::AID-ANIE1668>3.0.CO;2-Z) AND [10.1021/ja062508t](https://doi.org/10.1021/ja062508t)

Retrosynthesis ID: 18116

2.2.9 Takai olefination



Substrates:

- 1,1-Diiodoethane - *available at Sigma-Aldrich*
- CC(=O)C12C=C(Br)C(Br)C(C=O)C1COC2=O

Products:

- C/C=C/C1C(Br)C(Br)=CC2(C(C)=O)C(=O)OCC12


Typical conditions: CrCl2.THF.DMF

Protections: none

Reference: [10.1021/ja00283a046](#) and [10.1021/ja00237a081](#)

Retrosynthesis ID: 10942

2.2.10 Alkylation of amines with alkyl bromides



Substrates:

- 2-(p-Toluenesulfonylamino)phenylboronic acid pinacol ester - *available at Sigma-Aldrich*
- C/C=C/C1C(Br)C(Br)=CC2(C(C)=O)C(=O)OCC12

Products:

1. C/C=C/C1C(N(c2ccccc2B2OC(C)(C)C(C)(C)O2)S(=O)(=O)c2ccc(C)cc2)C(Br)=CC2(C(C)=O)C(=O


Typical conditions: K₂CO₃ or other base

Protections: none

Reference: [10.1016/j.tetlet.2007.09.110](#)

Retrosynthesis ID: 7668

2.2.11 Suzuki coupling of arylboronic pinacol esters with vinyl Bromides



Substrates:

1. C/C=C/C1C(N(c2ccccc2B2OC(C)(C)C(C)(C)O2)S(=O)(=O)c2ccc(C)cc2)C(Br)=CC2(C(C)=O)C(=O

Products:

1. C/C=C/C1C2C(=CC3(C(C)=O)C(=O)OCC13)c1ccccc1N2S(=O)(=O)c1ccc(C)cc1

Typical conditions: Pd catalyst.base.solvent

Protections: none

Reference: [10.1021/cr00039a007](#) and [10.1007/3418_2012_32](#) and [10.1021/cr0505268](#) and [10.1016/j.jfluchem.2016.01.018](#) and [10.1039/C3CS60197H](#)

Retrosynthesis ID: 10695