CLINICAL NOTE · BASELINE, TREATMENT SELECTION & PROGRESS TRACKING

Myopia Management Program

Track axial length and cycloplegic refraction over time, document risk, treatment choice and informed consent, and judge efficacy at each 6-monthly review.

Completed by: Clinician | Use as: Note template | Time: 15–25 min

Patient name	Date of birth	Patient ID / MRN	Date of service	Clinician

01 Risk profile

Age at myopia onset

Myopic parents ○ 0 ○ 1 ○ 2

Outdoor time h/day

Near work / screens h/day

Other risk factors

- ☐ Rapid progression (≥ 0.50 D/yr)
- ☐ Siblings myopic
- ☐ Near esophoria / high lag
- ☐ Early onset (< 8 y)
- ☐ Intensive schooling
- ☐ Other: _____

02 Baseline measurements

Baseline	Cyclo SE	Sphere / cyl / axis	Axial length	AL / CR ratio	K mean	Pupil (photopic)	Device
OD	D		mm		D	mm	
OS	D		mm		D	mm	

Near phoria Δ

Accommodative lag (MEM) D

Binocular VA

Axial length percentile for age / sex

03 Treatment options discussed

Option	Discussed	Selected	Details / parameters
Low-dose atropine (0.01% / 0.025% / 0.05%)	☐	☐	Strength, nightly, both eyes
Defocus spectacle lenses (DIMS, HAL / H.A.L.T.)	☐	☐	Brand, full-time wear
Dual-focus / EDOF soft contact lenses	☐	☐	Brand, power, days/week
Orthokeratology	☐	☐	Design, target
Combination therapy	☐	☐	
Single-vision correction only / monitor	☐	☐	Reason

Informed consent

Several options are used off-label in some jurisdictions. Record that benefits, limitations (slows rather than stops progression), costs, side-effects and wear requirements were explained, and that the parent/guardian consented.

Consent obtained ○ Verbal — documented ○ Written — on file

Discussed with

Annual cost quoted

04 Progress log

every 6 months

Date	Age	Treatment / adherence	Cyclo SE OD	Cyclo SE OS	AL OD (mm)	AL OS (mm)	ΔAL / yr OD · OS	Side-effects	Decision

Judge efficacy primarily on **axial elongation** (refraction alone can be masked by ortho-k or atropine). Consider escalation or combination if progression remains above the expected treated rate for age.

05 Today's plan

Assessment

Plan (continue / modify / escalate / stop)

Advice given

- ☐ ≥ 2 h outdoor daylight per day
- ☐ 20-20-20 breaks
- ☐ Reading distance ≥ 30 cm
- ☐ Adherence & side-effect check
- ☐ Lens / drop supply

Next review

Axial length due

Clinician signature

×

Clinician name & credentials

Date / time signed