LOGO

[Practice name]
[Street address] · [City, State / Region, Postcode]
[Phone] · [Email] · [Website]

PROCEDURE CONSENT · COVERS A COURSE OF INJECTIONS

Intravitreal Injection Informed Consent

Injections of medicine into the eye treat conditions that cause swelling or abnormal blood vessels in the retina. Most patients need a series of treatments.

Completed by: Patient + clinician | Use as: Consent form (e-signature) | Time: 5–10 min

Patient full name	Date of birth	Patient ID (office use)	Date completed

01 Treatment

Condition

- ☐ Wet (neovascular) AMD
- ☐ Geographic atrophy
- ☐ Diabetic macular oedema
- ☐ Retinal vein occlusion
- ☐ Myopic CNV
- ☐ Other: _____

Eye(s)

○ Right ○ Left ○ Both (same day permitted)

Planned course / duration of consent

Medicine(s) consented

- ☐ Aflibercept
- ☐ Bevacizumab (off-label)
- ☐ Avacincaptad pegol
- ☐ Faricimab
- ☐ Brolucizumab
- ☐ Dexamethasone implant
- ☐ Ranibizumab (or biosimilar)
- ☐ Pegcetacoplan
- ☐ Triamcinolone
- ☐ Other: _____

Off-label use

Bevacizumab is approved for cancer treatment and is widely used off-label in the eye after preparation by a compounding pharmacy. I understand this and have been offered the approved alternatives.

Initials

02 What happens

After numbing drops and antiseptic cleaning, a small device holds the eyelids open and the medicine is injected through the white of the eye with a very fine needle. It usually feels like brief pressure. Eye pressure and vision are checked afterwards.

Benefits & alternatives

Injections aim to stabilise and often improve vision; they do not cure the condition and may need to continue long term. Stopping or delaying treatment may cause permanent vision loss. Alternatives may include observation, laser, photodynamic therapy, steroid treatment or a different medicine.

Initials

03 Risks

Common and temporary

Initials

- Red patch on the white of the eye (subconjunctival haemorrhage)
- Grittiness, watering or mild ache for 1–2 days
- Floaters or small bubbles in vision
- Temporary rise in eye pressure

Rare but serious Initials

- Infection inside the eye (endophthalmitis) — needs urgent treatment
- Retinal tear or detachment
- Injury to the lens causing cataract
- Bleeding inside the eye
- Severe inflammation, including inflammation of retinal blood vessels (reported with some medicines)
- Long-term rise in eye pressure
- Possible effects elsewhere in the body (e.g. stroke, heart attack) — risk unclear
- For geographic atrophy treatments: increased chance of developing wet AMD

After every injection

Call the **same day** if you develop **increasing pain, worsening vision, increasing redness, light sensitivity or discharge**, or a **curtain or shadow** in your vision. Phone: Practice / after-hours phone

Pregnant, planning pregnancy or breastfeeding? ○ No ○ Yes — discussed with clinician

Allergy to iodine / povidone? ○ No ○ Yes

04 Consent

I consent to intravitreal injections of the medicine(s) ticked above into the eye(s) named, as a course of treatment, until I withdraw consent, my treatment changes, or 12 months — whichever is sooner. I may withdraw at any time.

Patient signature × | **Patient printed name** | **Date**

Parent / guardian / legal representative signature (if applicable) × | **Printed name** | **Relationship to patient** | **Date**

Clinician statement. I have explained the nature, purpose, benefits, risks and alternatives of the procedure, answered the patient's questions, and believe the patient understands and consents.

Surgeon / clinician signature × | **Name & credentials** | **Date**

Interpreter name & ID (if used) | **Language** | **Witness signature** × | **Witness name**