CLINICAL NOTE · INFANTS, PRESCHOOL & SCHOOL-AGE CHILDREN

Pediatric Eye Examination

Age-appropriate acuity, binocular and cycloplegic assessment with amblyopia, strabismus and myopia-risk documentation. Pair with the parent history (EYE-02).

Completed by: Clinician | Use as: Note template | Time: 20–40 min

Patient name	Date of birth	Patient ID / MRN	Date of service	Clinician

01 Visit

Age (years / months) _____ Accompanied by _____ Informant _____

Cooperation: ○ Excellent ○ Good ○ Fair ○ Poor

Reason for visit / parent concerns _____

Interval history (since EYE-02 / last visit) _____

Prematurity / ROP history: ○ None ○ Premature — no ROP ○ ROP (stage / treatment): _____

Wears glasses: ○ No ○ Yes — full time ○ Yes — part time

02 Visual acuity

Method

☐ Fix & follow ☐ CSM (central, steady, maintained) ☐ Preferential looking (Teller / Cardiff) ☐ LEA symbols

☐ HOTV ☐ Snellen / ETDRS ☐ Crowded optotypes

Acuity	Distance SC	Distance CC	Near	Method / test distance	Fixation preference
OD					
OS					
OU					

03 Binocular vision & motility

Deviation	Distance SC	Distance CC	Near SC	Near CC
Cover test	Δ	Δ	Δ	Δ
Hirschberg / Krimsky				

Stereo (test & result) _____ arc sec

Worth 4-dot (distance / near) _____


NPC _____ cm

Bruckner red reflex: ○ Symmetric ○ Asymmetric


Head posture: ○ Normal ○ Tilt ○ Turn ○ Chin up / down

Nystagmus: ○ None ○ Present: _____

Versions / ductions (-4 to +4) — OD OD



Versions / ductions (-4 to +4) — OS OS



04 Cycloplegic refraction

Agent

☐ Cyclopentolate 1% ☐ Cyclopentolate 0.5% (< 1 y) ☐ Tropicamide 1% ☐ Atropine 1% (home)

Drops × / time instilled

Time refracted

Retinoscopy	Sphere	Cylinder	Axis	Spherical equivalent	VA
OD					
OS					

Spectacle Rx prescribed	Sphere	Cylinder	Axis	Prism	Base	Add
OD						
OS						

Lens notes

☐ Full-time wear ☐ Polycarbonate / Trivex ☐ Flexible frame / strap

☐ Bifocal / add for accommodative ET ☐ Myopia-control lens (see EYE-14)

05 Ocular health

Anterior segment	OD	OS
Lids / lashes	☐ WNL	☐ WNL
Conjunctiva / sclera	☐ WNL	☐ WNL
Cornea	☐ WNL	☐ WNL
Anterior chamber	☐ WNL	☐ WNL
Iris / pupils	☐ WNL	☐ WNL
Lens	☐ WNL	☐ WNL

Posterior segment	OD	OS
Red reflex	☐ WNL	☐ WNL
Optic nerve (C/D)	☐ WNL	☐ WNL
Macula / foveal reflex	☐ WNL	☐ WNL
Vessels	☐ WNL	☐ WNL
Periphery	☐ WNL	☐ WNL

Dilated

○ Yes ○ No

IOP (if measured) OD / OS

mmHg

Method

06 Assessment

Amblyopia

☐ None ☐ Refractive (anisometropic / isoametropic) ☐ Strabismic

☐ Deprivation ☐ Mixed ☐ Amblyopia suspect

Severity (amblyopic eye BCVA)

○ Mild (20/25–20/30) ○ Moderate (20/40–20/80) ○ Severe (20/100–20/400)

Inter-ocular difference

lines

- ☐ **H53.02-** Refractive amblyopia
- ☐ **H53.03-** Strabismic amblyopia
- ☐ **H53.04-** Amblyopia suspect
- ☐ **H50.0-** Esotropia
- ☐ **H50.1-** Exotropia
- ☐ **H52.1-** Myopia
- ☐ **H52.0-** Hypermetropia
- ☐ **H52.2-** Astigmatism
- ☐ **H52.31** Anisometropia
- ☐ **Z01.020/1** Exam after failed vision screening

#	Diagnosis / assessment	ICD-10	Eye	Status / severity	Plan
1					
2					
3					

07 Plan

Treatment

- ☐ Glasses only — re-check acuity in 6–12 weeks
- ☐ Patching
- ☐ Atropine penalisation
- ☐ Dichoptic / binocular digital therapy
- ☐ Vision therapy
- ☐ Myopia management (EYE-14)
- ☐ Refer: paediatric ophthalmology / strabismus surgery
- ☐ Observe

Patching: hours / day & eye

Atropine: strength, eye, frequency

Expected review

Parent education & school

- ☐ Diagnosis explained
- ☐ Wear-time expectations
- ☐ Seat near front of class
- ☐ School report / letter (EYE-34)
- ☐ Outdoor time 2 h/day advised
- ☐ Screen-break habits

Return to clinic

☐ 1 day ☐ 1 week ☐ 2–4 weeks ☐ 3 months ☐ 6 months ☐ 12 months ☐ PRN

Specific RTC date / interval

Clinician signature

×

Clinician name & credentials

Date / time signed