CLINICAL NOTE · MACULAR DISEASE ASSESSMENT + INJECTION PROCEDURE

Retina: AMD & Intravitreal Injection Note

Classify AMD / macular disease per eye, record OCT activity, decide treat-and-extend interval and document the injection with time-out, drug lot and post-injection checks.

Completed by: Clinician + technician | Use as: Note template | Time: 10–15 min

Patient name	Date of birth	Patient ID / MRN	Date of service	Clinician

01 History

Indication nAMD, GA, DME, RVO-ME, myopic CNV

Injection no. (this eye)

Last agent / date / interval

Symptoms since last visit

☐ Stable ☐ Worse vision ☐ Distortion ☐ New scotoma ☐ Amsler change ☐ Pain / redness after last injection ☐ Floaters

AREDS2 supplement

○ Taking ○ Not taking ○ N/A

Smoking

○ Never ○ Former ○ Current

Anticoagulants / allergies (iodine)

02 Assessment

Function	BCVA (Snellen / letters)	IOP pre (mmHg)	Amsler	Lens status
OD				
OS				

OCT	CST (µm)	IRF	SRF	PED / sub-RPE	SHRM / haem	GA / atrophy	vs last visit
OD		○ Y ○ N	○ Y ○ N				○ Better ○ Same ○ Worse
OS		○ Y ○ N	○ Y ○ N				○ Better ○ Same ○ Worse

AMD stage	Classification
OD	○ No AMD ○ Early ○ Intermediate ○ GA non-central ○ GA central ○ nAMD active ○ nAMD inactive ○ Disciform scar
OS	○ No AMD ○ Early ○ Intermediate ○ GA non-central ○ GA central ○ nAMD active ○ nAMD inactive ○ Disciform scar

AREDS: **early** medium drusen (63–125 µm) · **intermediate** large drusen (> 125 µm) or pigment changes · **advanced** GA or neovascular. ICD-10: dry AMD H35.31[eye][stage 1 early · 2 intermediate · 3 GA without subfoveal involvement · 4 GA with subfoveal involvement]; wet AMD H35.32[eye][1 active CNV · 2 inactive CNV · 3 inactive scar].

FA / ICG / OCTA (if done)

Fundus findings

Decision

○ Inject today ○ Observe ○ Hold — reason: _____

Treat-and-extend

○ Extend ○ Maintain ○ Shorten

Next interval

weeks

03 Injection procedure

Pre-procedure time-out (all confirmed)

☐ Two patient identifiers
☐ Consent signed (EYE-27)
☐ Eye marked & verified
☐ Drug & dose verified
☐ Allergies reviewed
☐ No active lid / ocular infection

Drug	Agent & dose	Lot no.	Expiry	HCPCS & units	Wasted (JW/JZ)
OD					
OS					

- ☐ J0178 Aflibercept 1 mg
- ☐ J2777 Faricimab 0.1 mg
- ☐ J2778 Ranibizumab 0.1 mg
- ☐ J9035 Bevacizumab 10 mg (off-label)
- ☐ J7312 Dexamethasone implant 0.1 mg
- ☐ 67028 Intravitreal injection

Anaesthesia

☐ Topical drops ☐ Lidocaine gel
☐ Subconjunctival lidocaine

Antisepsis

☐ Povidone-iodine 5% conjunctiva ☐ Povidone-iodine lids / lashes
☐ Chlorhexidine (iodine allergy)

Speculum / quadrant e.g. inferotemporal

Distance from limbus mm

Needle gauge G

Injected by

Time

Post-injection	Vision post	Optic nerve perfusion	IOP post (time)	Complications
OD	○ CF ○ HM ○ Other	○ Perfused ○ Not		
OS	○ CF ○ HM ○ Other	○ Perfused ○ Not		

Post-injection warning signs

Give written instructions: mild grittiness and a subconjunctival haemorrhage are common. Call the **same day** for **increasing pain, worsening vision, increasing redness or light sensitivity** (possible endophthalmitis) or a curtain / shadow (possible detachment).

04 Plan

Assessment & plan

Fellow eye plan / home monitoring

Next visit (weeks) & eye

Next imaging

Next agent

○ Same ○ Switch: _____

Clinician signature

×

Clinician name & credentials

Date / time signed