PROCEDURE CONSENT · SLT, IRIDOTOMY, YAG CAPSULOTOMY, RETINAL LASER

Laser Eye Procedure Informed Consent

Tick the procedure you are having. Read the general information and the section for your procedure, then initial and sign.

Completed by: Patient + clinician | Use as: Consent form (e-signature) | Time: 5 min

Patient full name	Date of birth	Patient ID (office use)	Date completed

01 Procedure

Eye
○ Right (OD) ○ Left (OS) ○ Both

Clinician

Date

Procedure
○ Selective laser trabeculoplasty (SLT)
○ Laser peripheral iridotomy (LPI)
○ YAG laser capsulotomy
○ Pan-retinal photocoagulation (PRP)
○ Focal / grid macular laser
○ Laser retinopexy (retinal tear)

General information

Initials

Laser treatment is performed in the clinic while you sit at a microscope. Numbing drops and a special contact lens are usually used. You may see bright flashes. Most people return to normal activities the same or next day. All laser procedures can cause a temporary rise in eye pressure, inflammation, blurred vision and light sensitivity, and may need to be repeated.

02 Procedure-specific information

read your section

SLT — to lower eye pressure in glaucoma

Initials

Treats the eye's drainage channels to improve fluid outflow. Effect develops over 4–8 weeks, may not work for everyone, and often wears off over years; it can usually be repeated. You may still need drops. Risks: pressure spike, inflammation, redness, rarely corneal swelling.

LPI — to open or protect a narrow drainage angle

Initials

Makes a tiny opening in the iris to reduce the risk of angle-closure glaucoma. Risks: bleeding from the iris, pressure spike, inflammation, glare, ghost images or a line of light in vision, need for a second treatment; rarely cataract progression.

YAG capsulotomy — to clear a cloudy capsule after cataract surgery

Initials

Opens the clouded membrane behind the lens implant to restore clear vision. Risks: floaters, pressure rise, macular swelling, small marks on the implant, and rarely retinal tear or detachment or implant movement.

Retinal laser — PRP, focal / grid, retinopexy

Initials

Seals leaking vessels, reduces abnormal vessel growth or welds the retina around a tear. Aims to prevent further loss rather than improve vision. Risks: discomfort, reduced side or night vision (PRP), blind spots, worsening of macular swelling, and rarely accidental burn of the central vision; several sessions may be needed.

03 Warning signs

Call **immediately** for severe eye pain, headache with nausea, sudden worsening of vision, or a new shower of floaters, flashes or a curtain over your vision.

04 Consent

I have read the information for my procedure, had my questions answered, and consent to the laser procedure ticked above.

Patient signature | **Patient printed name** | **Date**

× | | |

Parent / guardian / legal representative signature (if applicable) | **Printed name** | **Relationship to patient** | **Date**

× | | | |

Clinician statement. I have explained the nature, purpose, benefits, risks and alternatives of the procedure, answered the patient's questions, and believe the patient understands and consents.

Surgeon / clinician signature | **Name & credentials** | **Date**

× | | |

Interpreter name & ID (if used) | **Language** | **Witness signature** | **Witness name**

| | × | |