

Experiment 3: kB for Gain 150

Madeleine Allen Edward Piper

2/6/2019

Recording the resistor values measured during lab.

Experiment 3: Johnson Noise - Boltzmann Constant

```
experiment3data1<-read.csv("/Users/mallen/Documents/128AL/JohnsonNoise128AL/experiment3data1.csv")
```

Calculate Vmeas, V, and Vsystem

```
Vsys<- experiment3data1[1,7] #first row 7th column  
VsysError <- experiment3data1[1,9]
```

```
Vmeask1<- (experiment3data1[2,7])  
Vmeask10<-experiment3data1[3,7]  
Vmeask20 <-experiment3data1[4,7]  
Vmeask32<-experiment3data1[5,7]  
Vmeask48<-experiment3data1[6,7]  
Vmeask100<-experiment3data1[7,7]
```

```
Vmeas<-c(Vmeask1, Vmeask10, Vmeask20, Vmeask32, Vmeask48, Vmeask100)
```

```
VmeasError<-sqrt((sum(experiment3data1[2:7,9])^2))
```

```
V<- sqrt(-Vsys^2+Vmeas^2)  
Verror<- sqrt(VmeasError^2+ VsysError^2)
```

Calculating G

```
gain <- data.frame(Frequency = vout1$x, Gain = (m_vout[2]/m_in))
```

```
capacitance <-87.875*(10^-12)  
capacitanceError <- .594*(10^-12)  
#df is just the x component  
#also changed here  
riemanSum <- function(f){  
  area<-(125/2)*(f[1]+2*sum(f[2:398])+f[399])  
  return(area)  
}
```

```
C = capacitance  
integrand <- data.frame(  
  gain[2]^2/(1+(2*pi*C*vin1$x*short)^2),  
  gain[2]^2/(1+(2*pi*C*vin1$x*k1)^2),  
  gain[2]^2/(1+(2*pi*C*vin1$x*k10)^2),  
  gain[2]^2/(1+(2*pi*C*vin1$x*k20)^2),
```

```

    gain[2]^2/(1+(2*pi*C*vin1$x*k35)^2),
    gain[2]^2/(1+(2*pi*C*vin1$x*k48)^2),
    gain[2]^2/(1+(2*pi*C*vin1$x*k100)^2)
  )
area <- data.frame(
  G1 =0,
  G2 =0,
  G3 =0,
  G4 =0,
  G5 =0,
  G6 =0,
  G7 =0

)
for(i in 1:length(integrand))
{
  area[i] <- riemanSum(unlist(integrand[i]))
}

```

So this returns a gain value G for each resistor (called “area”)

```

#prepare the data for graphing
kb<- 1.38064852 *10^-23 #m2 kg s-2 K-1

area2<-area[2:7]/100 #take away the short's data
resistors2 <-resistors[1:6]
y_value2<- (V^2)/(4*resistors2*area2)
y2<- unlist(y_value2, use.names=FALSE)

#calculate the temperatures
Temperature<- ((V^2)/(4*kb*area2*resistors2))
Temperature2<-unlist(Temperature, use.names = FALSE)
fit <- lm(y2~Temperature2)

Temperature2<-Temperature2/100
Temperature2[2]<- Temperature2[2]/10
print(Temperature2)

```

```
## [1] 70.75617 66.41194 66.79345 66.13097 68.47735 64.37755
```

```
Temperature2Error<-Temperature2*sqrt((V^2/Verror^2)+(resistors2/resistorserror)^2)
```

Calculating Errors

```

tempAvg<-mean(Temperature2)
t1 = (((tempAvg-Temperature2[1])^2))/sqrt(6)
t2 = (((tempAvg-Temperature2[2])^2))/sqrt(6)
t3 = (((tempAvg-Temperature2[3])^2))/sqrt(6)
t4 = (((tempAvg-Temperature2[4])^2))/sqrt(6)
t5 = (((tempAvg-Temperature2[5])^2))/sqrt(6)
t6 = (((tempAvg-Temperature2[6])^2))/sqrt(6)

Temperature2Error2<- sqrt(c(t1, t2, t3, t4, t5, t6))


```

```

library(ggplot2)
qplot((Temperature2), (y2))+geom_errorbar(aes(x=(Temperature2), ymin=(y2-Temperature2Error2), ymax=(y2+T

```

```
geom_smooth(method="lm", se=FALSE, fullrange=TRUE, level=0.95)+labs(title = "Temperature vs Voltage/G
```



```
summary(fit)
```

```
## Warning in summary.lm(fit): essentially perfect fit: summary may be
## unreliable
```

```
##
```

```
## Call:
```

```
## lm(formula = y2 ~ Temperature2)
```

```
##
```

```
## Residuals:
```

```
##      1      2      3      4      5      6
```

```
## 8.087e-35 -6.240e-37 -2.308e-35 -2.306e-35 -1.108e-35 -2.303e-35
```

```
##
```

```
## Coefficients:
```

```
##              Estimate Std. Error  t value Pr(>|t|)
```

```
## (Intercept)  7.863e-35  2.319e-35  3.391e+00  0.0275 *
```

```
## Temperature2 1.381e-23  8.340e-40  1.655e+16  <2e-16 ***
```

```
## ---
```

```
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
##
```

```
## Residual standard error: 4.544e-35 on 4 degrees of freedom
```

```
## Multiple R-squared:  1, Adjusted R-squared:  1
```

```
## F-statistic: 2.741e+32 on 1 and 4 DF, p-value: < 2.2e-16
```

```
plot(fit$residuals, main = "Residuals of the fit line", ylab= "Residuals")
```

Residuals of the fit line

