

Experiment 2: Temperatures for Gain 150

Madeleine Allen, Edward Piper

2/7/2019

Experiment 2: Johnson Noise 1

Recording the resistor values measured during lab.

```
#everything will be in ohms
short<- .03
shortError<-.001

k20<-20090
k20error<-1

k35 <- 35230 #secretly 35.2 but that would be an ugly variable name
k35error<-1

k100 <- 100700
k100error <- 1

k10 <- 999.05
k10error<- .01

k1 <- 998.17
k1error <- .01

k48 <- 48650 #secretly 48.7k but again that would be an ugly variable name
k48error<- 1

resistors<-c(k1,k10,k20,k35, k48,k100)
resistorerror<-c(k1error, k10error, k20error, k35error, k48error, k100error)
```

Import Band Voltage measurements from experiment 2

```
experiment2data<-read.csv("/Users/mallen/Documents/128AL/JohnsonNoise128AL/experiment2data1.csv")
```

Calculate Vmeas, V, and Vsystem

```
Vsys<- experiment2data[1,7] #first row 7th column
VsysError <- experiment2data[1,9]

Vmeask1<- (experiment2data[2,7])
Vmeask10<-experiment2data[3,7]
Vmeask20 <-experiment2data[4,7]
Vmeask32<-experiment2data[5,7]
Vmeask48<-experiment2data[6,7]
Vmeask100<-experiment2data[7,7]

Vmeas<-c(Vmeask1, Vmeask10, Vmeask20, Vmeask32, Vmeask48, Vmeask100)

#need to redo the error later (2/5)
```

```
VmeasError<-sqrt((sum(experiment2data[2:7,9])^2))

V<- sqrt(-Vsys^2+Vmeas^2)
Verror<- sqrt(VmeasError^2+ VsysError^2)
```

Calculating G

```
capacitance <-87.875*(10^-12)
capacitanceError <- .594*(10^-12)
#df is just the x component

riemanSum <- function(fa,fb){
  area <-0.5*(125)*(fb-fa)+fa*125
  return(area)
}

#resistors<-read.csv("experiment2data1.csv")

C = capacitance
integrand <- data.frame(
  gain[2]^2/(1+(2*pi*C*vin1$x*short)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k1)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k10)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k20)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k35)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k48)^2),
  gain[2]^2/(1+(2*pi*C*vin1$x*k100)^2)
)
area <- data.frame(
  G1 =0,
  G2 =0,
  G3 =0,
  G4 =0,
  G5 =0,
  G6 =0,
  G7 =0

)
for(i in 1:length(integrand))
{
  for(l in 1:398)
  {
    if(is.na(integrand[l+1,i]))
    {
      break
    }
    else
    {
      area[i] <- area[i]+ riemanSum(integrand[l,i],integrand[l+1,i])
    }
  }
}
}
```

```
area2error=sqrt((capacitance/capacitanceError)^2+(resistors/resistorerror)^2)
```

So this returns a gain value G for each resistor (called "area")

```
#resistors<-c(0,1000,10000,20000,32500, 48700,100000)
#plot(resistors,area, main= "Resistor value vs G", ylab = "Values of G", xlab = "Resistor Values (Ohm)".
```

Plotting R as a function of V^2 , kB, and G

```
kb<- 1.38064852 *10^-23 #m2 kg s-2 K-1
```

```
area2<-area[2:7] #take away the short's data
y_value<- (V^2)/(4*kb*area2) #area is the vector that contains all G's
```

```
#prepare data for graphing
resistors2 <-resistors[1:7]
y<- unlist(y_value, use.names=FALSE)
```

```
#I'll try finding temperatures
```

```
Temperature<- ((V^2)/(4*kb*area2*resistors2))
Temperature2<-unlist(Temperature, use.names = FALSE)
Temperature2[2]<-Temperature2[2]/10
print(Temperature2) #these are the correct values
```

```
## [1] 252.4061 251.8954 255.7764 252.9490 252.5065 243.3532
```

Create a fit line

```
#
#resistor as the x axis and the other term as the y axis.
#
resistors3<-resistors2[1:6]
resistors3error <- resistors3*sqrt((V^2/Verror^2)+((area2/area2error)^2))
fit <- lm(y~0+resistors3)
```

```
rm(gain)
```

```
rm(gainerror)
```


```
gain=data.frame()
```

```
gainerror=data.frame()
```

```
library(ggplot2)
```

```
qplot(unlist(resistors3),unlist(y))+geom_errorbar(aes(x=unlist(resistors3), ymin=unlist(y-resistorerror),
  geom_smooth(method="lm", se=FALSE, fullrange=TRUE, level=0.95))+labs(title = "Resistance as a function
```

Resistance as a function of Gain and Voltage with a 1D Fit




```
summary(fit)
```

```
##
## Call:
## lm(formula = y ~ 0 + resistors3)
##
## Residuals:
##      1      2      3      4      5      6
##  6154 2270554 191574 236337 304835 -290761
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## resistors3    246.241      8.756   28.12 1.06e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1042000 on 5 degrees of freedom
## Multiple R-squared:  0.9937, Adjusted R-squared:  0.9925
## F-statistic: 790.9 on 1 and 5 DF, p-value: 1.064e-06
```

```
plot(fit$residuals, main = "Residuals of the fit line", ylab= "Residuals")
```

Residuals of the fit line



```
#lines(resistors3, predict(fit, data.frame(resistors3)), col="red")  
#legend("topleft", legend=c("Fit Line"), col=c("red"), lty=1:2, cex=0.8)
```

Then, to find the value of absolute 0 in Celsius, we use the recorded room temperature (20.5 C) and the average Temperature (277.75K). Solving for absolute 0, you get -257.3. We expected to find -273.15.