# Thursday 26 March 2026

Published 25 Mar 2026, 17:00:00
Valid from 25 Mar 2026, 17:00:00 until 26 Mar 2026, 17:00:00
Written by Avalanche Service Bavaria
**translated with DeepL**

## The avalanche prone locations are more frequent in high and congested areas.




| 1 | 2 | 3 | 4 | 5 |
|---|---|---|---|---|
| low | moderate | considerable | high | very high |

# Thursday 26 March 2026

Published 25 Mar 2026, 17:00:00
Valid from 25 Mar 2026, 17:00:00 until 26 Mar 2026, 17:00:00
Written by Avalanche Service Bavaria
**translated with DeepL**

## Danger Level 2 - Moderate




## Weak layers in new and wind slab snow

The avalanche risk is moderate. New fallen snow is the main problem. The avalanche prone locations are in steep terrain adjacent to ridgelines in all aspects, in gullies and bowls and behind slope discontinuities. Slab avalanches can be triggered with little additional load. They usually remain small, but can also reach medium size in heavily snow-covered areas.

Small gliding avalanches can release on their own on previously apery, very steep meadow slopes.

## Snowpack

Around 30 to 40 cm of new fallen snow with strong, sometimes stormy winds from a north-westerly direction. Where there is still snow, the new and wind-slab snow is lying on a slightly damp surface of old snow. On very steep north-facing slopes at higher altitudes, it is lying on soft snow. The snowdrift accumulations can become large in places and be covered by new fallen snow. Weak layers prone to triggering are embedded within the wind slab. The old snowpack is generally well stabilised and slightly moist. In many places, the new fallen snow covers the previously apery ground and starts gliding snow.

## Tendency

The avalanche danger can increase further with wind.