Friday 27 February 2026

Published 26 Feb 2026, 17:00:00
Valid from 26 Feb 2026, 17:00:00 until 27 Feb 2026, 17:00:00
Written by Avalanche Service Bavaria
translated with DeepL

Spring conditions with persistent weak layer problem and risk of cornice breakage

earlier



later



1	2	3	4	5
low	moderate	considerable	high	very high

Friday 27 February 2026

Published 26 Feb 2026, 17:00:00
Valid from 26 Feb 2026, 17:00:00 until 27 Feb 2026, 17:00:00
Written by Avalanche Service Bavaria
translated with DeepL

Danger Level 3 - Considerable

earlier



later



Danger of falling on a hard frozen snow surface!

The avalanche danger above 1500 m increases from moderate to considerable during the daytime changes. Below that, it is low. Until the morning, gliding snow is the main problem at medium altitudes. Small and medium-sized gliding avalanches will spontaneously release themselves on a few steep slopes with a smooth surface. Staying under glide cracks should be avoided.

When the sun shines, the activity of sliding snow increases on sunny slopes and small and medium-sized, wet loose snow avalanches can be expected from extremely steep terrain that has not yet been discharged.

In addition, weak persistent layers above 1500 metres are problematic. With large additional loads, for example due to cornice breakage, large slab avalanches can also be triggered in isolated cases in the west,

north and east aspects. Take particular care in transition areas from a little to a lot of snow.

Snowpack

On the surface, the snowpack freezes overnight in many places and is capable of bearing loads; in the sun, it softens again during the day and soaking progresses. Up to high altitudes, the snowpack is slightly moist, in medium altitudes it is sometimes completely soaked to the ground. In shady slopes at higher altitudes, faceted crystals can be found deeper in the snowpack.

Tendency

Slow decline in the wet snow problem - the persistent weak layer problem remains.