Tuesday 3 March 2026

Published 2 Mar 2026, 17:00:00
Valid from 2 Mar 2026, 17:00:00 until 3 Mar 2026, 17:00:00
Written by Avalanche Service Bavaria
translated with DeepL

Avoid large additional loads on the snowpack.



1	2	3	4	5
low	moderate	considerable	high	very high

Tuesday 3 March 2026

Published 2 Mar 2026, 17:00:00
Valid from 2 Mar 2026, 17:00:00 until 3 Mar 2026, 17:00:00
Written by Avalanche Service Bavaria
translated with DeepL

Danger Level 2 - Moderate





There is a risk of falling on hard snow surfaces.

The avalanche risk is moderate above 1700 metres and low below that. Persistent weak layer is a persistent weak layer. With large additional loads, medium slab avalanches can be triggered on shady slopes. Take particular care at high altitudes in transition areas from little to much snow, for example when entering gullies and bowls.

At medium altitudes, gliding avalanches are possible on slopes with a smooth surface. Gliding avalanches can also occur on shady slopes and at any time of day or night. They can be small to medium in size.

Snowpack

In shady slopes at higher altitudes, weak layers consisting of faceted crystals can be found deep in the snowpack. These are usually covered by compact or crusted can form layers of snow. Early in the morning, the snowpack at medium altitudes is frozen hard and firns in the sun during the day. Towards the ground, the snowpack is partly wet and can glide snow. In the foothills, the ground is increasingly pitting on the south side.

Tendency

Slow decline in the wet snow problem - the persistent weak layer problem remains.