

# Spheroid, basement membrane, matrix:

the three-entity model and the arrows between them

the container note · prototype/ecm · log/okuda\_ECM · companions: note\_junction, note\_fibre, note\_surface,  
note\_sheet · 10 August 2026

## 1. Mechanism

An epithelial spheroid growing inside an extracellular matrix is *three* entities and not two, and the middle one is not a passive spacer. The literature fixes its proportions, and they are not the convenient ones.

**The basement membrane is thin, stiff and squat-anchored.** It is  $\sim 100$  nm thick; its adhesions are clusters, not molecules: an integrin is a heterodimer, integrins assemble into nanoclusters of 20–50 at  $\sim 35$ –100 nm [14], and those cluster again into adhesions with a measured  $\sim 555$  nm centre-to-centre spacing [15]. The **plaque** set is the MIDDLE level. Vertically the cluster sits  $\sim 40$  nm from the actin it pulls on [16], so an adhesion is several times *wider* than the sheet is thick and a fraction of its own spacing across. And the sheet is 10 to 100 times stiffer than the stroma outside it, not equal to it.

**It is not a permanent solid.** Collagen IV in the fly embryo has a half-life of 7–10 hours [13] and in mouse hair follicle about 3, and slowing that turnover disrupts morphogenesis. A sheet enclosing a tissue that triples its radius does not stretch to 2.4; it is *replaced* at the new size while it is being stretched.

**And it is the interface.** While the sheet is intact the epithelium does not touch the stroma: load goes cell  $\rightarrow$  plaque  $\rightarrow$  sheet  $\rightarrow$  anchoring fibril  $\rightarrow$  stroma [17]. An epithelium that *also* pushes the stroma directly is counting the same push twice. That single observation is what removes **cell\_to\_ecm** from the model and puts two edge sets in its place.

## 2. Equations and symbols

Each equation below belongs to one arrow of Fig. 1b. The companion notes carry the mechanics of each entity; this note carries what passes *between* them.

**Mass, not just mechanics.** The sheet carries an areal density  $\rho$  of collagen IV produced by the epithelium and removed by proteolysis, on a surface whose area is growing:

$$\frac{D\rho}{Dt} = \underbrace{s(\mathbf{x}, t)}_{\text{secretion}} - \underbrace{\frac{\rho}{\tau_{\text{bm}}}}_{\text{turnover}} - \underbrace{\rho \frac{\dot{A}}{A}}_{\text{dilution}}, \quad s^* = \rho \left( \frac{1}{\tau_{\text{bm}}} + \frac{\dot{A}}{A} \right), \quad \tau_{\text{bm}} = \frac{t_{1/2}}{\ln 2} \approx 4\text{--}14 \text{ h}. \quad (1)$$

The two removal terms are the same size here — a spheroid tripling its radius over a day has  $\dot{A}/A \sim 10^{-1} \text{ h}^{-1}$  against  $1/\tau_{\text{bm}} \sim 10^{-1} \text{ h}^{-1}$ . Turnover is not a correction to growth; it is the same size as growth, which is why the sheet can accommodate it at all.

**A nonlinear sheet.** A single Young’s modulus is wrong by construction: the same membrane reads  $\sim 0.4$  MPa at low strain and  $\sim 3$  MPa at high [12], and a growing spheroid takes it across that range:

$$E(\lambda) = E_0 [1 + \beta(\lambda - 1)], \quad \lambda = \sigma_{\text{max}}(\mathbf{F}), \quad \frac{E_{\text{bm}}}{E_{\text{ecm}}} \in [10, 100]. \quad (2)$$

**MPa is not a quantity this box has** — it is dimensionless and the stroma’s modulus is the number 15 — so the falsifiable forms are the two ratios and never the pascals.

**Adhesion, with its reaction and with slip.** A plaque is two-sided: whatever force it applies to the sheet it applies equally and oppositely to the cell, which a replayed epithelium discards. And the sheet is not pinned — it *slides* — so the tangential law is friction against relative velocity and not a spring to a frozen direction:

$$\mathbf{f}_p^{\text{adh}} = \kappa_n (\ell_p - \ell_0) \hat{\mathbf{n}}_p - \xi (\mathbf{v}_{\text{bm}} - \mathbf{v}_{\text{epi}})_{\parallel}, \quad \mathbf{f}_p^{\text{epi}} = -\mathbf{f}_p^{\text{adh}}, \quad (3)$$

$$n_p = \Sigma^{-2} \text{ plaques per unit area, } \Sigma \approx 7T, \quad \ell_0 \approx 0.3T. \quad (4)$$

$\ell_0$  is a material property and not a tuning constant: it is what sets the standoff.

**Sheet to stroma.** Anchoring fibrils [17] load the stroma *through* the sheet, which is also what lets a stiff sheet spread a local epithelial push over a soft matrix:

$$\mathbf{t} = k_a (\mathbf{x}_{\text{ecm}} - \mathbf{x}_{\text{bm}}) \text{ on anchored points, } \mathbf{f}^{\text{bm}} = - \sum_{\text{anchors}} \mathbf{t}. \quad (5)$$




Figure 1: (a) The three entities in section, at the proportions the literature measures rather than the ones that are convenient: the adhesion is a squat plaque, not a thread, and the sheet is thin against its own spacing. (b) Every operator, in the direction it acts. Filled bullets carry a force and must therefore have a reaction; open bullets carry a signal or a material flux. There is no arrow between the epithelium and the matrix.

The barrier is geometric rather than mechanical: pores of 10–130 nm pass solutes, stop cells [11], and are what a proteolytic operator has to open before anything crosses.

**The matrix pushes back on the cell cycle.** The force the stroma returns, collected by direction and divided by the area each direction covers, is a pressure; a cell facing a stressed matrix grows more slowly, which is what a spheroid grown under a measured external stress does [9, 10]:

$$g = f + \frac{1 - f}{1 + (P/p_{1/2})^n}, \quad s_{\text{now}} \leftarrow s_{\text{prev}} (1 + g (f_{\text{grow}} - 1)). \quad (6)$$

Because it is the growth *rate* that is slowed and not the size directly, a small difference compounds over hundreds of frames — which is how a modest difference in stress between one direction and another becomes a difference in shape.

| symbol                      | is                                                 | of what                                                                                        |
|-----------------------------|----------------------------------------------------|------------------------------------------------------------------------------------------------|
| $\rho, \tau_{\text{bm}}, s$ | areal density, turnover time, secretion            | of collagen IV on the sheet, Eq. (1)                                                           |
| $A, \dot{A}/A$              | sheet area, its relative growth rate               | the dilution term; the SAME size as turnover                                                   |
| $E_0, \beta, \lambda$       | low-strain modulus, stiffening, principal stretch  | of the sheet, Eq. (2); report RATIOS, never MPa                                                |
| $T, \Sigma, \ell_0$         | thickness, plaque spacing, linkage rest length     | $\Sigma \approx 7T, \ell_0 \approx 0.3T$ ; $T$ never appears as geometry                       |
| $\kappa_n, \xi$             | plaque normal stiffness, tangential friction       | against RELATIVE velocity — $\xi$ is what lets the sheet stay attached while the tissue slides |
| $k_a$                       | fibril stiffness                                   | sheet to stroma, Eq. (5)                                                                       |
| $P, p_{1/2}, n, f$          | matrix pressure, gate half-point, sharpness, floor | Eq. (6); $f \leq g \leq 1$ multiplies the growth RATE                                          |

### 3. The Plexus2 container

**Two structural claims, and both differ from what the archived line built.** An adhesion is not a body and a fibril is not a particle: each is a *relation between two entities*, which in Plexus2 is an edge set carrying `.pre` and `.post` index buffers rather than a set of nodes with positions. Writing the hemidesmosome as an edge set is what makes its reaction automatic — one operator returns a delta to both endpoints, so the force the sheet feels and the force the cell feels cannot get out of step. The archived line made the adhesion first a field on the membrane and then a column of MPM particles, and both lost the reaction.

| set          | kind                                                    | one element is                | state it carries                                     |
|--------------|---------------------------------------------------------|-------------------------------|------------------------------------------------------|
| spheroid     | node, depth 0                                           | the organ                     | centroid, cell count                                 |
| cell         | node, depth 1                                           | one epithelial cell           | $A, P, V$ and targets, age, $\rho_f$                 |
| vertex       | node, depth 1                                           | a corner of the apical mesh   | <b>pos</b>                                           |
| junction     | <b>edge</b> $\subseteq$ <b>vertex</b> <sup>2</sup>      | a cell-cell contact           | $n_e, \ell_e$                                        |
| bm           | node on a surface mesh                                  | a patch of membrane           | <b>pos</b> , <b>F</b> , $\rho$ , age, <b>occ</b>     |
| bm_face      | <b>hyperedge</b> $\subseteq$ <b>bm</b> <sup>3</sup>     | a triangle of membrane        | $\mathbf{D}_m^{-1}$ , $\mathbf{C}_0$ , $A^0$ , $Y_2$ |
| plaque       | <b>edge</b> <b>cell</b> $\rightarrow$ <b>bm</b>         | one hemidesmosome             | $\ell_0$ , load, bound                               |
| fibril       | <b>edge</b> <b>bm</b> $\rightarrow$ <b>mpm_particle</b> | one anchoring fibril          | rest length, load                                    |
| mpm_particle | node, depth 1                                           | a patch of <i>stroma only</i> | <b>pos</b> , <b>F</b> , fibre direction              |
| mpm_grid     | <b>field</b>                                            | the stroma's momentum         | mass, momentum                                       |
| protease     | <b>field</b>                                            | MMP concentration             | scalar, diffusing                                    |

| operator                                                     | kind             | acts on         | mechanism                                       | where its gates are |
|--------------------------------------------------------------|------------------|-----------------|-------------------------------------------------|---------------------|
| <i>epithelium</i>                                            |                  |                 |                                                 |                     |
| grow_3d, divide_3d                                           | Divide           | cell            | volume growth and cytokinesis [6, 7]            | 00; note_junction   |
| shape_energy_3d                                              | Lateral          | junction        | vertex-model force balance [8]                  | 00                  |
| medioapical_myosin,                                          | Lateral,         | cell,           | two myosin pools and the ring                   | note_junction J1–   |
| junction_myosin,                                             | Structural       | junction        |                                                 | J11                 |
| cytokinetic_ring                                             |                  |                 |                                                 |                     |
| <i>epithelium</i> $\rightarrow$ <i>basement membrane</i>     |                  |                 |                                                 |                     |
| bm_secrete,                                                  | Divide, Die      | bm_face         | Eq. (1) [13, 18]                                | note_sheet G24–     |
| bm_degrade                                                   |                  |                 |                                                 | G26, G19–G23        |
| bm_stiffness,                                                | Lateral          | bm_face         | Eq. (2)                                         | note_sheet G1–G8    |
| bm_elastic                                                   |                  |                 |                                                 |                     |
| <i>basement membrane</i> $\leftrightarrow$ <i>epithelium</i> |                  |                 |                                                 |                     |
| plaque_seed,                                                 | Rewire,          | plaque          | Eqs. (3)–(4), <i>with the reaction</i> [16]     | note_sheet G9–      |
| plaque_pull,                                                 | Lateral          |                 |                                                 | G13                 |
| plaque_rupture                                               |                  |                 |                                                 |                     |
| <i>basement membrane</i> $\leftrightarrow$ <i>matrix</i>     |                  |                 |                                                 |                     |
| fibril_seed,                                                 | Rewire,          | fibril          | Eq. (5) [17]                                    | open                |
| fibril_pull                                                  | Lateral          |                 |                                                 |                     |
| mesh_contact                                                 | Lateral          | contact         | particle-to-surface, both endpoints in one call | note_surface S1–    |
|                                                              |                  |                 |                                                 | S19                 |
| bm_barrier                                                   | Exchange         | bm_face, fields | pores pass solutes, stop cells [11]             | open                |
| <i>matrix</i>                                                |                  |                 |                                                 |                     |
| seed_ecm,                                                    | mpm_*, Seed, Ex- | mpm_particle    | MLS-MPM [4, 5]                                  | note_fibre F1–F18   |
| ecm_stress                                                   | change, Lateral  |                 |                                                 |                     |
| ecm_growth_gate_3d                                           | Lateral          | cell            | Eq. (6) [9, 10]                                 | 58; open            |

**One schedule, split at exactly one place.** Anything carrying a force runs inside the substep block, at the stroma's rate; anything changing topology or mass runs once per frame, because a set whose membership changed mid-substep would be integrated against two different index sets in one step:

frame ( $\Delta t = 1$ ): cell\_geometry\_3d  $\rightarrow$  grow\_3d  $\rightarrow$  bm\_growth\_gate  $\rightarrow$  medioapical\_myosin  $\rightarrow$  junction\_myosin  
 $\rightarrow$  shape\_energy\_3d  $\rightarrow$  reconnect\_t1\_3d  $\rightarrow$  divide\_3d  $\rightarrow$  cytokinetic\_ring  $\rightarrow$  junction\_myosin  
 $\rightarrow$  bm\_refine  $\rightarrow$  bm\_secrete  $\rightarrow$  bm\_degrade  $\rightarrow$  plaque\_seed  $\rightarrow$  plaque\_rupture

substep ( $\Delta t_{\text{sub}} = \Delta t/8$ ): bm\_stiffness  $\rightarrow$  bm\_elastic  $\rightarrow$  plaque\_pull  $\rightarrow$  fibril\_pull  
 $\rightarrow$  mpm\_strain  $\rightarrow$  mpm\_scatter  
 $\rightarrow$  mpm\_grid\_update  $\rightarrow$  mpm\_gather

**Four rules, each buying something specific.** *The membrane never enters the grid* — it is  $\sim 100$  nm on a  $\sim 200 \mu\text{m}$  spheroid, so resolving it as MPM material needs a  $3500^3$  lattice; as a surface mesh it costs what the epithelium costs and  $\Delta x$  is set by the stroma alone. *The epithelium stays out of the substep block* — it is overdamped and massless, so it does not need the matrix's CFL step, and putting its relaxation inside would multiply its cost by eight and change nothing. *The sheet is massless too*, so its stiffness costs drag rather than substeps: the bound is  $\Delta t(zk_b + \kappa)/\gamma < 2$ , a rate and not a wave speed, and the 10–100 stiffness ratio is paid for with  $\gamma$ . *Particles go on the shell that carries stress, not on the box* — the far field is quiescent, and filling the box at a usable density costs a million particles.

**And one thing this deletes.** cell\_to\_ecm and cell\_exclude\_3d go, because while the sheet is intact the basement membrane *is* the interface (§1). MPM fibres also stop being asked to act as ropes, which note\_fibre measured they cannot.




Figure 2: **(a)** Every set and field, and which kind each is. A node set carries positions; an edge or hyperedge set is a *relation* and carries state belonging to neither endpoint — a junction to neither cell, a plaque to neither the cell nor the sheet, a triangle to none of its three nodes. **(b)** Every operator, on the set it acts on, coloured by kind. Anything carrying a *force* sits on a set whose elements have positions; anything carrying *mass* or *number* sits on the set that owns the quantity. **bm\_refine** is the only operator here that is numerical rather than biological.

### 3.1 The two diffusible species, and the one that is not

Proteolysis introduces the first *fields* in this model, and deciding which of the three species deserves to be one is not a modelling preference — it is arithmetic.

| species        | what it is                                       | where it lives                 | why                                                                |
|----------------|--------------------------------------------------|--------------------------------|--------------------------------------------------------------------|
| <b>MMP</b>     | soluble MMP-2/9, $\sim 72$ kDa, cuts collagen IV | <b>field</b> on <b>bm_face</b> | it diffuses freely, $D \sim 10\text{--}100 \mu\text{m}^2/\text{s}$ |
| <b>TIMP</b>    | soluble inhibitor, binds MMP 1:1                 | <b>field</b> on <b>bm_face</b> | likewise — and it is the <i>sink</i>                               |
| <b>MT1-MMP</b> | membrane-type MMP14, transmembrane               | a per-cell <i>state</i>        | it never enters the extracellular space                            |

**A soluble protease cannot open a hole, and the arithmetic says so before any run.** In one 600 s frame a soluble MMP travels  $\sqrt{4Dt} = 155\text{--}490 \mu\text{m}$ , against a spheroid  $318 \mu\text{m}$  across and a cell of  $10 \mu\text{m}$ . It is therefore *uniform over the entire sheet within a single frame*, and a uniform protease dissolves rather than perforates — the same failure uniform starvation produced, for the same reason. Localising to one cell would need  $D < 0.042 \mu\text{m}^2/\text{s}$ ,  $240\text{--}2400\times$  slower than any protein of that size. That is not a parameter range, it is a different molecule, and the different molecule is MT1-MMP: *tethered*, acting where its cell put it. This is why focal basement-membrane breach is done by membrane-type MMPs and invadopodia rather than by soluble enzyme.

**What makes the soluble field worth solving at all is the sink.** With  $Dc/Dt = D\nabla_s^2 c + s - kc$  the source decays over  $\sqrt{D/k}$ , so the field acquires a *length* instead of filling the sphere — and  $k$  here is not a fitted decay but the reaction with TIMP. The breach diameter becomes a *prediction* of the MMP/TIMP balance: a  $10 \mu\text{m}$  breach needs an effective  $k$  of  $0.1\text{--}0.5 \text{s}^{-1}$  (half-life  $1.4\text{--}6.9$  s), a  $50 \mu\text{m}$  breach  $0.004\text{--}0.02 \text{s}^{-1}$  ( $35\text{--}170$  s). A number to check, not to choose.

**Three new operators, and one costs what the others do not.** **protease\_secrete** (Lateral, on **cell**) is the source; **protease\_inhibit** (Exchange) is the 1:1 reaction; **protease\_diffuse** (Exchange) is the only operator in this model carrying a *spatial* coupling, and so the only one whose stability limit involves the mesh. Explicit diffusion needs  $\Delta t D/h^2 < \frac{1}{2}$  — at these  $D$ ,  $1,300\text{--}13,400$  substeps per frame, *quadrupling at every 1 $\rightarrow$ 4 refinement*, unlike the sheet's own bound which is a rate and mesh-independent. It is therefore solved semi-implicitly,  $(\mathbf{I} + \Delta t D\mathbf{L})c = c_{\text{old}}$  by conjugate gradient on a finite-volume face Laplacian: unconditionally stable, one solve per frame, indifferent to  $h$ .

**The same argument applied to the integrin gives the opposite answer for the opposite reason.** A receptor is well mixed because it is *confined* to one cell's basal face ( $\sqrt{4Dt} = 15.5 \mu\text{m}$  over a  $10 \mu\text{m}$  cell) and needs

no field; a soluble protease would be well mixed because it *travels far*. **Confinement, not diffusivity, decides whether a species is worth resolving as a field.**

## Proteolysis, and the mechanism the literature forced

05g\_degrade put the three species on the entities the confinement argument assigns them and measured which one can make a hole. The answer was unambiguous and one-sided:

| arm                 | faces dead  | under MT1 | elsewhere   | sheet breached |
|---------------------|-------------|-----------|-------------|----------------|
| both arms           | 317         | 317       | 0           | 6.2%           |
| soluble only        | <b>0</b>    | 0         | 0           | 0%             |
| tethered only       | 317         | 317       | 0           | 6.2%           |
| no protease         | 0           | 0         | 0           | 0%             |
| <b>no inhibitor</b> | <b>5120</b> | 317       | <b>4803</b> | <b>100%</b>    |

The last row is the argument of §3.1 as an experiment: **remove TIMP and the sheet dissolves entirely**, with 4803 of the deaths nowhere near a cell that expresses MT1-MMP. With the inhibitor present the diffusible arm kills nothing at all and the tethered arm kills exactly its own footprint. A soluble protease has no length scale unless something quenches it.

**Two defects that the runs found rather than the design.** First, neither field had a *clearance* term, so TIMP accumulated without bound (0.0318 at frame 20, 0.0438 at frame 25) — its only sink was a 1:1 reaction with an MMP twenty times scarcer, so there was no steady state for the pair to reach. A first-order clearance is not a refinement: the  $\sqrt{D/k}$  that justifies solving the field *is* that  $k$ . With it, both fields plateau and drift by  $< 10^{-4}$  over twenty frames. Second, a five-point sweep of the MMP:TIMP source ratio over a hundredfold range returned *identically* 317 dead at every point, because the direct tethered cut had been set  $350\times$  larger than anything the soluble arm could contribute — the sweep axis had no effect on the answer, and the holes were the source stencil in every case.

## TIMP-2 is the adaptor as well as the inhibitor, and that predicts a bell

The pure-inhibitor law of 05g is half of the mechanism. In the canonical model [1] TIMP-2 is the *bridging adaptor* that makes activation possible: its C-terminus tethers proMMP-2 and its N-terminus binds MT1-MMP, and the prodomain is then cleaved by a *second*, *TIMP-2-free* MT1-MMP [2]. TIMP-2 therefore both presents the zymogen and poisons the enzyme that must cut it, and the dose response is biphasic. The dissection is clean: an artificial receptor that tethers proMMP-2 *without* inhibiting MT1-MMP has no descending limb [3], so the fall-off is inhibition and not saturation.

With  $T$  the tethered enzyme and  $K$  the MT1–TIMP dissociation constant, occupancy is Langmuir and the rate needs one occupied site and one free one:

$$[\text{MT1}\cdot\text{TIMP}] = T \frac{c_T/K}{1 + c_T/K}, \quad [\text{MT1}_{\text{free}}] = \frac{T}{1 + c_T/K}, \quad \text{act} \propto T^2 c_{\text{pro}} \frac{c_T/K}{(1 + c_T/K)^2}, \quad (7)$$

which is bell-shaped and **peaks at**  $c_T = K$ . Both the shape and the location are therefore predictions, and 05g's pure-inhibitor law cannot produce a peak at all — it can only decay — so one sweep distinguishes them and 05g is the control.

**Measured** (05h\_ternary, eight TIMP levels, four species): the activation rate rises and falls,  $4.5 \times 10^{-2} \rightarrow 6.2 \times 10^{-2} \rightarrow 2.4 \times 10^{-2} \rightarrow 7.5 \times 10^{-3} \rightarrow 2.5 \times 10^{-3}$ , and the pure-inhibitor control over the same sweep is monotone decreasing, as Eq. (7) requires of it. Two arrangements had to be corrected before the curve could be read at all, and both are worth recording because they are measurement errors rather than modelling ones: the activation had to be made *rate-limiting* (at a large  $k_{\text{act}}$  the whole zymogen pool converts every frame whatever TIMP does, and saturation erases the dose response); and the rate has to be read *while the substrate still exists*, since at the TIMP levels where the mechanism works best it destroys every MT1-bearing face and a late-window average then reports exactly zero at the peak.

## 4. Gates

A gate is a number with a threshold decided *before* the run and a stated consequence if it fails. The per-entity gates live in the companion notes and are indexed in §3; this table is the *system* level — what has to be true of the assembled model, and which arrows are not yet built.

| gate                                                                                  | thresholdmeasured                                                                                                                                 |
|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>the epithelium is unchanged by everything attached to it</i> (00 is the reference) |                                                                                                                                                   |
| B1 cells, radius, aspect, T1 rate of the bare tissue                                  | the num-<br>bers every<br>later run<br>must re-<br>produce bit-<br>identical 200 → 6076, × <b>3.56</b> , <b>1.021</b> , <b>0.0089</b> /cell/frame |
| B2 adding per-junction myosin at $m_e \equiv 1$ changes nothing                       | <b>bit-identical</b>                                                                                                                              |
| <i>each arrow carries its reaction</i>                                                |                                                                                                                                                   |
| B3 epithelium → matrix, momentum residual                                             | $< 10^{-6}$ <b>1.5–1.8e-7</b> across every sweep run ( <b>note_surface</b> S5, S12)                                                               |
| B4 cell ↔ sheet, as bookkeeping and as motion                                         | $< 10\epsilon_{64}$ ; <b>1.4e-16</b> , <b>4.4e-15</b> ( <b>note_sheet</b> G9, G10)                                                                |
|                                                                                       | drift $< 10^{-12}$                                                                                                                                |
| B5 sheet → stroma: the stroma is loaded THROUGH the sheet                             | displacement <b>bt built</b> — <b>fibril_pull</b> is the missing arrow with the sheet present vs absent                                           |
| <i>the middle entity does what a membrane does</i>                                    |                                                                                                                                                   |
| B6 the sheet reports its own deformation                                              | $> 0.95$ of the geomet-<br>ric stretch within 10% <b>0.9935</b> as a surface; <b>0.13</b> as MPM material (run 130)                               |
| B7 $\rho$ held flat while $A$ triples                                                 | <b>1/11.1</b> with no secretion ( <b>note_sheet</b> G24)                                                                                          |
| B8 the standoff is the linkage's rest length                                          | $\ell/\ell_0 \in [0.9, 1.1]$ <b>3.43</b> ( <b>note_sheet</b> G12)                                                                                 |
| <i>the coupling is mutual</i>                                                         |                                                                                                                                                   |
| B9 the tissue feels the matrix: shape differs from the free control                   | any separation oblateness <b>one-way</b> — the reaction is computed, recorded and dropped against the uncon-<br>fined run                         |
| B10 the growth gate changes the shape it is given                                     | <b>1.43</b> grown vs <b>1.41</b> pressed (58, 48)                                                                                                 |

**One convention makes these mean what they say. A reaction is measured on its own force pair**, inside the substep and before any drive is added; with the drive in the sum it stops being a statement about the reaction. That is why B3 and B4 are separate rows with different thresholds: B3 is float32 through a grid, B4 is float64 on an explicit edge.

**What the gates license.** The epithelium is reproducible on its own and nothing attached to it has moved it (B1, B2); every arrow that has been built returns its reaction to machine precision (B3, B4); the sheet, as a surface, reports the deformation it is given (B6); and the growth gate can produce a shape that pressing produces too, which is the one place the matrix has been shown to change the tissue (B10). **What they do not license** is the assembled model: B5 is not built, B7 and B8 fail on the sheet's own testbench, and B9 is the architecture — the tissue is a replay, so it drives and is never driven. Until B9 passes, the stroma's modulus is very nearly a free parameter: the tissue never feels the matrix, and the growth gate normalises the pressure map to its own late-run median, so a softer stroma changes the substep it needs and not the shape that comes out. That is the clearest statement of what one-way coupling costs, and **mesh\_contact** returning **vertex\_force** to a live epithelium is the run that ends it.

**What sixty-six archived runs settled, and it is four things.** Runs 91–164 asked where the sheet sits and what holds it there, and they answer that question and no other. The standoff is a *tracking lag*,  $\delta - \dot{R}\Delta t\gamma/\kappa$ , and not a force balance, which two fibre lengths differing fourfold confirmed to  $3 \times 10^{-5}$  box units. A force routed through the grid leaves the sheet carrying 100% of its true stretch while the same force integrated by the engine leaves it carrying 13% (B6). An MPM fibre cannot act as a rope, because  $\mathbf{F}$  is advected by the grid's velocity gradient and never measures the distance between two particles. And every MPM body in the prototype silently carried the *parent* set's modulus, so a membrane declaring  $E = 400$  ran at the stroma's 15 until each body was given its own parent — which is why run 146 came back bit-identical to run 144 at ten times the stiffness. None of the four is a model derived from what a basement membrane *is*; that is what **note\_sheet** builds.

## 5. References

### References

- [1] Karagiannis, E.D., Popel, A.S. (2004) A theoretical model of type I collagen proteolysis by matrix metalloproteinase (MMP) 2 and membrane type 1 MMP in the presence of tissue inhibitor of metalloproteinase 2. *J. Biol. Chem.*

- 279**(37):39105–39114. — the canonical ODE network for MT1-MMP / TIMP-2 / proMMP-2 / MMP-2, and the source of the reaction topology used here.
- [2] Sato, H., Takino, T. (2010) Coordinate action of membrane-type matrix metalloproteinase-1 (MT1-MMP) and MMP-2 enhances pericellular proteolysis and invasion. *Cancer Sci.* **101**(4):843–847.
  - [3] Activation of MMP-2 by MT1-MMP through an artificial receptor for proMMP-2 generates active MMP-2. *Cancer Res.* (2008) **68**(21):9096. — the dissection showing the descending limb is MT1-MMP inhibition and not receptor saturation.
  - [4] Sulsky, D., Chen, Z., Schreyer, H.L. (1994) A particle method for history-dependent materials. *Comput. Methods Appl. Mech. Eng.* **118**:179–196.
  - [5] Hu, Y. *et al.* (2018) A moving least squares material point method with displacement discontinuity and two-way rigid body coupling. *ACM Trans. Graph.* **37**(4):150.
  - [6] Honda, H., Tanemura, M., Nagai, T. (2004) A three-dimensional vertex dynamics cell model of space-filling polyhedra. *J. Theor. Biol.* **226**(4):439–453.
  - [7] Okuda, S., Inoue, Y., Eiraku, M., Sasai, Y., Adachi, T. (2013) Reversible network reconnection model for simulating large deformation in dynamic tissue morphogenesis. *Biomech. Model. Mechanobiol.* **12**(4):627–644.
  - [8] Okuda, S., Miura, T., Inoue, Y., Adachi, T., Eiraku, M. (2018) Combining Turing and 3D vertex models reproduces autonomous multicellular morphogenesis. *Sci. Rep.* **8**:2386.
  - [9] Helmlinger, G., Netti, P.A., Lichtenbeld, H.C., Melder, R.J., Jain, R.K. (1997) Solid stress inhibits the growth of multicellular tumor spheroids. *Nat. Biotechnol.* **15**(8):778–783.
  - [10] Montel, F. *et al.* (2011) Stress clamp experiments on multicellular tumor spheroids. *Phys. Rev. Lett.* **107**:188102.
  - [11] Yurchenco, P.D. (2011) Basement membranes: cell scaffoldings and signaling platforms. *Cold Spring Harb. Perspect. Biol.* **3**(2):a004911.
  - [12] Candiello, J. *et al.* (2007) Biomechanical properties of native basement membranes. *FEBS J.* **274**(11):2897–2908.
  - [13] Matsubayashi, Y. *et al.* (2020) Rapid homeostatic turnover of embryonic ECM during tissue morphogenesis. *Dev. Cell* **54**(1):33–42.
  - [14] Changede, R., Sheetz, M. (2017) Integrin and cadherin clusters: a robust way to organize adhesions for cell mechanics. *BioEssays* **39**(1):1600123. — nascent adhesions are  $\sim 100$  nm and contain  $\sim 50$  integrins; large adhesions are loose aggregates of such clusters.
  - [15] *Differential spatial regulation and activation of integrin nanoclusters inside focal adhesions.* eLife reviewed preprint 105270. — STORM: nanocluster median size 35–40 nm, centre-to-centre  $\sim 555$  nm, spacing set cell-intrinsically. Cited as a preprint.
  - [16] Kanchanawong, P. *et al.* (2010) Nanoscale architecture of integrin-based cell adhesions. *Nature* **468**:580–584.
  - [17] Keene, D.R., Sakai, L.Y., Lunstrum, G.P., Morris, N.P., Burgeson, R.E. (1987) Type VII collagen forms an extended network of anchoring fibrils. *J. Cell Biol.* **104**(3):611–621.
  - [18] Kelley, L.C. *et al.* (2019) Adaptive F-actin polymerization and localized ATP production drive basement membrane invasion in the absence of MMPs. *Dev. Cell* **48**(3):313–328.