

June 14, 2021

### Abstract

We calculate quantum corrections to the mass of the vortex in  $N = 2$  supersymmetric abelian Higgs model in  $2+1$  dimensions. We put the system in a box and apply the zeta function regularization. The boundary conditions inevitably violate a part of the supersymmetries. Remaining supersymmetry is however enough to ensure isospectrality of relevant operators in bosonic and fermionic sectors. A non-zero correction to the mass of the vortex comes from finite renormalization of couplings.

PACS: 11.27.+d; 12.60.Jv

$$\rho_B(\omega) - \rho_F(\omega) \propto \delta(\omega), \quad (1)$$

$$\Delta E \propto \int d\omega \omega (\rho_B(\omega) - \rho_F(\omega)). \quad (2)$$

$$\Delta E = \Delta E_B - \Delta E_F, \quad \Delta E_{B,F} = \frac{1}{2} \sum_{\omega_{B,F}} \omega_{B,F}, \quad (3)$$

$$\Delta E_{B,F}^{\text{reg}} = \frac{1}{2} \sum_{\omega_{B,F} \neq 0} \omega_{B,F}^{1-2s}, \quad (4)$$

$$\mathcal{L} = \mathcal{L}_B + \mathcal{L}_F, \quad (5)$$

$$\mathcal{L}_B = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - |D_\mu \phi|^2 - \frac{1}{2} (\partial_\mu w)^2 - \frac{e^2}{2} (|\phi|^2 - v^2)^2 - e^2 w^2 |\phi|^2, \quad (6)$$

$$\mathcal{L}_F = i\bar{\psi}\gamma^\mu D_\mu \psi + i\bar{\chi}\gamma^\mu \partial_\mu \chi - i\sqrt{2}e(\bar{\psi}\chi\phi - \bar{\chi}\psi\phi^*) + ew\bar{\psi}\psi, \quad (7)$$

$$\begin{aligned} \delta A_\mu &= i(\bar{\eta}\gamma_\mu \chi - \bar{\chi}\gamma_\mu \eta), \\ \delta \phi &= \sqrt{2}\bar{\eta}\psi, \quad \delta w = i(\bar{\chi}\eta - \bar{\eta}\chi), \\ \delta \chi &= \gamma^\mu \eta \left( \partial_\mu w + \frac{i}{2} \epsilon_{\mu\nu\lambda} F^{\nu\lambda} \right) + i\eta(e|\phi|^2 - ev^2), \\ \delta \psi &= -\sqrt{2}(i\gamma^\mu \eta D_\mu \phi - ew\phi) \end{aligned} \quad (8)$$

$$\gamma^0 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \gamma^2 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \quad (9)$$

$$\gamma^\mu \gamma^\nu = -g^{\mu\nu} - i\epsilon^{\mu\nu\rho} \gamma_\rho. \quad (10)$$

$$\eta = \begin{pmatrix} \eta_+ \\ \eta_- \end{pmatrix}, \quad (11)$$

$$(D_1 + iD_2)\phi = 0, \quad (12)$$

$$F_{12} + e(|\phi|^2 - v^2) = 0. \quad (13)$$

$$\phi = f(r)e^{in\theta}, \quad eA_j = \epsilon_{jk} \frac{x^k}{r^2} (a(r) - n) \quad (14)$$

$$\begin{aligned} \frac{1}{r} \frac{d}{dr} a(r) &= e^2 (f^2(r) - v^2), \\ r \frac{d}{dr} \ln f(r) &= a(r). \end{aligned} \quad (15)$$

$$f(0) = 0, \quad f(\infty) = v, \quad (16)$$

$$a(0) = n, \quad a(\infty) = 0. \quad (17)$$

$$E^{\text{cl}} = 2\pi n v^2. \quad (18)$$

$$\mathcal{L}_{\text{gf}} = -\frac{1}{2} [\partial_\mu \alpha^\mu - ie(\varphi^* \phi - \varphi \phi^*)]^2, \quad (19)$$

$$\mathcal{L}_{\text{ghost}} = \sigma^* (\partial_\mu \partial^\mu - 2e^2 \phi \phi^*) \sigma. \quad (20)$$

### Abstract

We discuss a new class of brane models (extending both  $p$ -brane and  $Dp$ -brane cases) where the brane tension appears as an *additional dynamical degree of freedom* instead of being put in by hand as an *ad hoc* dimensionfull scale. Consistency of dynamics naturally involves the appearance of additional higher-rank antisymmetric tensor gauge fields on the world-volume which can couple to charged lower-dimensional branes living on the original  $Dp$ -brane world-volume. The dynamical tension has the physical meaning of electric-type field strength of the additional higher-rank world-volume gauge fields. It obeys Maxwell (or Yang-Mills) equations of motion (in the string case  $p = 1$ ) or their higher-rank gauge theory analogues (in the  $Dp$ -brane case). This in particular triggers a simple classical mechanism of (“color”) charge confinement.

$$\omega_F^2 \begin{pmatrix} U \\ V \end{pmatrix} = \begin{pmatrix} D_F D_F^\dagger, & 0 \\ 0, & D_F^\dagger D_F \end{pmatrix} \begin{pmatrix} U \\ V \end{pmatrix}, \quad (21)$$

$$U = \begin{pmatrix} \psi_+ \\ \chi_- \end{pmatrix}, \quad V = \begin{pmatrix} \psi_- \\ \chi_+ \end{pmatrix} \quad (22)$$

$$D_F = \begin{pmatrix} D_+, & -\sqrt{2}e\phi \\ -\sqrt{2}e\phi^*, & \partial_- \end{pmatrix}, \quad -D_F^\dagger = \begin{pmatrix} D_-, & \sqrt{2}e\phi \\ \sqrt{2}e\phi^*, & \partial_+ \end{pmatrix}. \quad (23)$$

$$D_\pm := D_1 \pm iD_2, \quad \partial_\pm := \partial_1 \pm i\partial_2. \quad (24)$$

$$\Delta_w = -\partial_j \partial_j + 2e^2 |\phi|^2. \quad (25)$$

$$\omega_B^2 \begin{pmatrix} \varphi \\ i\alpha_+ \sqrt{2} \end{pmatrix} := -\partial_0^2 \begin{pmatrix} \varphi \\ i\alpha_+ \sqrt{2} \end{pmatrix} = D_F^\dagger D \begin{pmatrix} \varphi \\ i\alpha_+ \sqrt{2} \end{pmatrix}, \quad (26)$$

$$D_F^\dagger D_F = - \begin{pmatrix} D_j^2 - e^2(3|\phi|^2 - v^2), & -\sqrt{2}e(D_- \phi) \\ -\sqrt{2}e(D_+ \phi^*), & \partial_j^2 - 2e^2|\phi|^2 \end{pmatrix}, \quad (27)$$

$$D_F D_F^\dagger = - \begin{pmatrix} D_j^2 - e^2(|\phi|^2 + v^2), & 0 \\ 0, & \partial_j^2 - 2e^2|\phi|^2 \end{pmatrix}. \quad (28)$$

$$u_\pm = e^{\pm i\theta} \left( u_r \pm \frac{i}{r} u_\theta \right), \quad (29)$$

$$\mathcal{B}^{[\alpha]} \alpha_\mu|_{\partial M} = 0, \quad \mathcal{B}^{[\sigma]} \sigma|_{\partial M} = 0, \quad (30)$$

$$\mathcal{B}^{[\alpha]} \partial_\mu \sigma|_{\partial M} = 0. \quad (31)$$

$$\alpha_0|_{\partial M} = 0, \quad \alpha_\theta|_{\partial M} = 0, \quad \left( \partial_r + \frac{1}{r} \right) \alpha_r|_{\partial M} = 0, \quad \sigma|_{\partial M} = 0. \quad (32)$$

$$\left( \partial_r + \frac{1}{r} \right) \partial_r \sigma = [-\Delta_w \sigma] + \left[ -\frac{1}{r^2} \partial_\theta^2 + 2\phi\phi^* \right] \sigma, \quad (33)$$

$$[-\Delta_w \sigma]|_{\partial M} = -\sum_k \omega_k^2 \sigma_k|_{\partial M} = 0. \quad (34)$$

$$w|_{\partial M} = 0. \quad (35)$$

$$\alpha_r = \Re(e^{-i\theta} \alpha_+) , \quad \alpha_\theta = r \Im(e^{-i\theta} \alpha_+) . \quad (36)$$

$$\Re(e^{-i\theta}\chi_+)|_{\partial M} = 0, \quad \left(\partial_r + \frac{1}{r}\right)\Im(e^{-i\theta}\chi_+)|_{\partial M} = 0. \quad (37)$$

$$V(\omega) = \omega^{-2}D_F^\dagger U(\omega), \quad (38)$$

$$U(\omega) = \omega^{-2}D_F V(\omega) \quad (39)$$

$$U_1(\omega) = \omega^{-2}\left(D_+ V_1(\omega) - \sqrt{2}e\phi V_2(\omega)\right). \quad (40)$$

$$\Re(e^{-i\theta}\phi^* U_1)|_{\partial M} = \Re(e^{-i\theta}\phi^* \psi_+)|_{\partial M} = 0. \quad (41)$$

$$U_2(\omega) = \omega^2\left(\partial_- V_2 - \sqrt{2}e\phi^* V_1\right). \quad (42)$$

$$\Im(U_2)|_{\partial M} = 0, \quad \Im(\phi^* V_1)|_{\partial M} = 0. \quad (43)$$

$$\Im(\phi^* \varphi)|_{\partial M} = 0, \quad (\partial_r - 2(\partial_r \ln \phi^*))\Re(\phi^* \varphi)|_{\partial M} = 0. \quad (44)$$

$$\partial_r \Re(\chi_-)|_{\partial M} = 0, \quad \left(\partial_r + \frac{1}{r}\right)\Im(e^{-i\theta}\phi^* \psi_+)|_{\partial M} = 0. \quad (45)$$

$$\delta\alpha_+ = 2i\eta_-^* \chi_+. \quad (46)$$

$$\Re(\eta_-) = 0 \quad (47)$$

$$\Delta E^{\text{ren}} = \Delta E(V+B)^{\text{ren}} - \Delta E(B)^{\text{ren}} + \Delta E^{\text{f.r.}}, \quad (48)$$

$$\Delta E_B^{\text{reg}} = \frac{1}{2} \sum_{\omega_B \neq 0} \omega_B^{1-2s} = \frac{1}{2} \sum_{\omega_F \neq 0} \omega_F^{1-2s} = \Delta E_F^{\text{reg}}. \quad (49)$$

$$\Delta E(V+B)^{\text{ren}} = 0. \quad (50)$$

$$\tilde{\Delta} = -\partial_j^2 + 2e^2 v^2. \quad (51)$$

$$\partial_r \phi = -\frac{i}{r} D_\theta \phi. \quad (52)$$

$$(D_- \phi) = e^{-i\theta} 2\partial_r \phi \rightarrow 0 \quad (53)$$

$$\tilde{\varphi} = e^{i\beta(r)\theta} \varphi, \quad \tilde{\psi} = e^{i\beta(r)\theta} \psi, \quad (54)$$

$$\begin{aligned} \left( \partial_r + \frac{1}{r} \right) \Im \left( e^{-i\theta} (i\alpha_+, \chi_+, \tilde{\psi}_+) \right) |_{\partial M} &= 0, \\ \Re \left( e^{-i\theta} (i\alpha_+, \chi_+, \tilde{\psi}_+) \right) |_{\partial M} &= 0. \end{aligned} \quad (55)$$

$$\begin{aligned} \partial_r \Re \left( (\tilde{\varphi}, \tilde{\psi}_-, \chi_-) \right) |_{\partial M} &= 0, \\ \Im \left( (\tilde{\varphi}, \tilde{\psi}_-, \chi_-) \right) |_{\partial M} &= 0. \end{aligned} \quad (56)$$

$$\Delta E(B)^{\text{ren}} = 0. \quad (57)$$

$$W_m(s) = \frac{1}{2} \sum \omega(m)^{1-2s} = \frac{1}{2} \zeta_m \left( s - \frac{1}{2} \right), \quad (58)$$

$$\zeta_m \left( s - \frac{1}{2} \right) = \Gamma \left( s - \frac{1}{2} \right)^{-1} \int d^2x \int_0^\infty dt t^{s-\frac{1}{2}-1} K(t, x). \quad (59)$$

$$K(t, x) = \langle x | e^{-t\Delta_m} | x \rangle = (4\pi t)^{-1} e^{-m^2 t}. \quad (60)$$

$$\mathcal{W}_m = -\frac{m^3}{12\pi}. \quad (61)$$

$$\mathcal{W}^{1-\text{loop}} = -\frac{e^3}{6\pi} \left[ (3|\phi|^2 - v^2)^{3/2} - (2|\phi|^2)^{3/2} \right]. \quad (62)$$

$$\mathcal{W}^{\text{tot}} = \mathcal{W}^{\text{cl}}(e + \hbar \delta e, v + \hbar \delta v) + \hbar \mathcal{W}^{1-\text{loop}}, \quad (63)$$

$$\mathcal{W}^{\text{cl}}(e, v) = \frac{e^2}{2} (|\phi|^2 - v^2) \quad (64)$$

$$\delta v = -\frac{e}{4\sqrt{2}\pi}. \quad (65)$$

$$\Delta E^{\text{f.r.}} = \hbar(\delta v) \frac{dE^{\text{cl}}}{dv} = -\frac{evn\hbar}{\sqrt{2}}. \quad (66)$$

$$\Delta E^{\text{ren}} = -\frac{evn\hbar}{\sqrt{2}}. \quad (67)$$

$$\delta \chi_- = -\eta_- \left( \partial_0 w + i\epsilon_{ojk} \partial^j \alpha^k - 2ie\Re(\phi^* \varphi) \right). \quad (68)$$

$$\Im(\delta \chi_-)|_{\partial M} \sim -\partial_0 w|_{\partial M} = 0, \quad (69)$$

$$0 = \partial_r \Re(\delta\chi_-)|_{\partial M} \sim \partial_r \left( -\epsilon_{ojk} \partial^j \alpha^k + 2e\Re(\phi^* \varphi) \right) |_{\partial M}. \quad (70)$$

$$X_i = \frac{x_i}{1-w} \quad \text{and} \quad \tilde{X}_i = \frac{x_i}{1+w}. \quad (71)$$

$$g = \text{diag} \left( 1, \frac{-4}{1+R^2}, \frac{-4}{1+R^2}, \frac{-4}{1+R^2} \right), \quad \text{where } R^2 = X_1^2 + X_2^2 + X_3^2. \quad (72)$$

$$\tilde{X}_i = \frac{1}{R^2} X_i. \quad (73)$$

$$\partial_{\tilde{X}_i} = (R^2 \delta_{ij} - 2X_i X_j) \partial_{X_j}. \quad (74)$$

$$\begin{aligned} e_0^0 &= 1, & \tilde{e}_0^0 &= 1, \\ e_i^i &= -\frac{1+R^2}{2}, & \tilde{e}_i^i &= \frac{1+\tilde{R}^2}{2}, \end{aligned} \quad \text{and} \quad (75)$$

$$e_\alpha^\mu e_\beta^\nu g_{\mu\nu} = \eta_{\alpha\beta}, \quad (76)$$

$$\hat{\tilde{e}}_i = \frac{1}{R^2} (2X_i X_j - \delta_{ij}) \hat{e}_j. \quad (77)$$

$$d\hat{\theta}^\alpha + \omega^\alpha{}_\beta \wedge \hat{\theta}^\beta = 0, \quad (78)$$

$$\omega^{\alpha\beta} = \frac{2}{1+R^2} (X^\alpha dX^\beta - X^\beta dX^\alpha), \quad (79)$$

$$\mathcal{L}_{\text{fermion}} = \bar{\psi} (i\gamma^\alpha e_\alpha{}^\kappa (\partial_\kappa + \Omega_\kappa)) \psi. \quad (80)$$

$$\mathcal{L}_{\text{fermion}} = \bar{\psi}(X_i, t) \left( i\gamma^0 \partial_t - i\gamma^i \left( \frac{1+R^2}{2} \partial_{X_i} - X_i \right) \right) \psi(X_i, t), \quad (81)$$

$$\mathcal{L}_{\text{fermion}} = \bar{\tilde{\psi}}(\tilde{X}_i, t) \left( i\gamma^0 \partial_t + i\gamma^i \left( \frac{1+\tilde{R}^2}{2} \partial_{\tilde{X}_i} - \tilde{X}_i \right) \right) \tilde{\psi}(\tilde{X}_i, t), \quad (82)$$

$$\tilde{\psi}(\tilde{X}_i) = \rho \psi(X_i). \quad (83)$$

$$T(\alpha)_{ij} = \cos \alpha \, \delta_{ij} + \sin \alpha \, \epsilon_{ijk} \hat{X}_k + \hat{X}_i \hat{X}_j (1 - \cos \alpha) \quad (84)$$

$$\rho = i\gamma_0\gamma_5\gamma_j\hat{X}_j. \quad (85)$$

$$\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}. \quad (86)$$

$$\tau_k\psi_i = \psi_i\sigma_k^T, \quad (87)$$

$$= -\psi_i\sigma_2\sigma_k\sigma_2, \quad (88)$$

$$f(\mu) = 2 \arctan \left( k \tan \frac{\mu}{2} \right). \quad (89)$$

$$\left( i\gamma^0\partial_t - i\gamma^i \left( \frac{1+R^2}{2}\partial_{X_i} - X_i \right) - gU^{\gamma_5} \right) \psi(X_i, t) = 0, \quad (90)$$

$$\left( i\gamma^0\partial_t + i\gamma^i \left( \frac{1+\tilde{R}^2}{2}\partial_{\tilde{X}_i} - \tilde{X}_i \right) - gU^{\gamma_5} \right) \tilde{\psi}(\tilde{X}_i, t) = 0, \quad (91)$$

$$U^{\gamma_5} = \exp \left( if(\mu(R))\gamma_5\tau_i \frac{X_i}{R} \right), \quad (92)$$

$$x = (\sin \mu \sin \theta \cos \phi, \sin \mu \sin \theta \sin \phi, \sin \mu \cos \theta, \cos \mu). \quad (93)$$

$$\begin{aligned} \mu &= \begin{cases} \arctan \frac{2R}{R^2-1} + \pi & \text{for } R^2 < 1, \\ \frac{\pi}{2} & \text{for } R^2 = 1, \\ \arctan \frac{2R}{R^2-1} & \text{for } R^2 > 1, \end{cases} \\ \theta &= \arctan \frac{\sqrt{X_1^2 + X_2^2}}{X_3}, \\ \phi &= \arctan \frac{X_2}{X_1}. \end{aligned} \quad (94)$$

$$\mu = \begin{cases} \arctan \frac{2\tilde{R}}{1-\tilde{R}^2} & \text{for } \tilde{R}^2 < 1, \\ \frac{\pi}{2} & \text{for } \tilde{R}^2 = 1, \\ \arctan \frac{2\tilde{R}}{1-\tilde{R}^2} + \pi & \text{for } \tilde{R}^2 > 1. \end{cases} \quad (95)$$

$$e_\mu = \begin{pmatrix} \sin \theta \cos \phi \\ \sin \theta \sin \phi \\ \cos \theta \end{pmatrix}, \quad e_\theta = \begin{pmatrix} \cos \theta \cos \phi \\ \cos \theta \sin \phi \\ -\sin \theta \end{pmatrix} \quad \text{and} \quad e_\phi = \begin{pmatrix} -\sin \phi \\ \cos \phi \\ 0 \end{pmatrix}. \quad (96)$$

$$\gamma^0 = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma_i \\ -\sigma_i & 0 \end{pmatrix}, \quad \text{and } \gamma_5 = \begin{pmatrix} 0 & I_2 \\ I_2 & 0 \end{pmatrix}. \quad (97)$$

$$\tilde{\psi}(\tilde{X}_i) = \begin{pmatrix} -i\sigma_j \hat{X}_j & 0 \\ 0 & -i\sigma_j \hat{X}_j \end{pmatrix} \psi(X_i). \quad (98)$$

$$\begin{aligned} L_x &= i \left( \sin \phi \frac{\partial}{\partial \theta} + \cot \theta \cos \phi \frac{\partial}{\partial \phi} \right), \\ L_y &= i \left( -\cos \phi \frac{\partial}{\partial \theta} + \cot \theta \sin \phi \frac{\partial}{\partial \phi} \right), \\ L_z &= -i \frac{\partial}{\partial \phi}. \end{aligned} \quad (99)$$

$$\hat{P} \psi(X_i) = \gamma_0 \psi(-X_i) \quad \text{and} \quad \hat{P} X_i \hat{P}^{-1} = -X_i. \quad (100)$$

$$\begin{aligned} \hat{P} \gamma_0 \hat{P}^{-1} &= \gamma_0 \quad \text{and} \quad \hat{P} \gamma_i \hat{P}^{-1} = -\gamma_i, \\ \hat{P} \partial_0 \hat{P}^{-1} &= \partial_0 \quad \text{and} \quad \hat{P} \partial_i \hat{P}^{-1} = -\partial_i. \end{aligned} \quad (101)$$

$$\hat{P} \tilde{\psi}(\tilde{X}_i) = -\gamma_0 \tilde{\psi}(\tilde{X}_i). \quad (102)$$

$$\frac{d}{d\mu} \begin{pmatrix} \tilde{F} \\ \tilde{G} \end{pmatrix} = \begin{pmatrix} -\frac{1+3\cos\mu}{2\sin\mu} + g \sin f & E + g \cos f \\ -(E - g \cos f) & \frac{3(1-\cos\mu)}{2\sin\mu} - g \sin f \end{pmatrix} \begin{pmatrix} \tilde{F} \\ \tilde{G} \end{pmatrix}. \quad (103)$$

$$+ \frac{3gf' \sin f}{2(E + g \cos f)} \frac{(1+u)}{\sqrt{1-u^2}} - \frac{g^2 f' \sin^2 f}{E + g \cos f} - gf' \cos f \Big) G(u) = 0, \quad (104)$$

$$+ \frac{3gf' \sin f}{2(E - g \cos f)} \frac{(1-u)}{\sqrt{1-u^2}} - \frac{g^2 f' \sin^2 f}{E - g \cos f} + gf' \cos f \Big) \tilde{G}(u) = 0. \quad (105)$$

$$F(u) = \sqrt{\frac{1+u}{1-u}} \tilde{G}(u). \quad (106)$$

$$F(u) = \frac{1}{E + g \cos f} \left( \left( -\frac{3(1+u)}{2\sqrt{1-u^2}} + g \sin f \right) G(u) - \sqrt{1-u^2} \frac{d}{du} G(u) \right), \quad (107)$$

$$\tilde{F}(u) = \frac{1}{E - g \cos f} \left( \left( \frac{3(1-u)}{2\sqrt{1-u^2}} - g \sin f \right) \tilde{G}(u) - \sqrt{1-u^2} \frac{d}{du} \tilde{G}(u) \right). \quad (108)$$



$$y''(u) + P(u)y'(u) + Q(u)y(u) = 0. \quad (109)$$

$$y(u) = \sum_{n=0}^{\infty} a_n (u - u_0)^n. \quad (110)$$

$$P(u) = \frac{p(u)}{u - u_0}, \quad (111)$$

$$Q(u) = \frac{q(u)}{(u - u_0)^2}, \quad (112)$$

$$p(u) = p_0 + p_1(u - u_0) + \dots \quad \text{and} \quad q(u) = q_0 + q_1(u - u_0) + \dots, \quad (113)$$

$$y(u) = (u - u_0)^\rho \sum_{n=0}^{\infty} a_n (u - u_0)^n. \quad (114)$$

$$\rho^2 + (p_0 - 1)\rho + q_0 = 0, \quad (115)$$

$$(u - u_0)^2 z''(u) + \left( 2(u - u_0)^2 \frac{y_1'(u)}{y_1(u)} + (u - u_0)p(u) \right) z'(u) = 0. \quad (116)$$

$$E(A, \Phi) = SW(A, \Phi) - 8\pi^2 Q, \quad (117)$$

$$F_A^+ = \sigma^+(\Phi \otimes \Phi^\dagger)_0, \quad (118)$$

$$\mathcal{D}_A \Phi = 0, \quad (119)$$

$$F_A^+ + \chi^+ = \sigma^+(\Phi \otimes \Phi^\dagger)_0, \quad (120)$$

$$\mathcal{D}_A \Phi = 0, \quad (121)$$

$$SW_\chi(A, \Phi) = E_\chi(A, \Phi) + 16\pi^2 K_\chi + 8\pi^2 Q. \quad (122)$$

$$[x^\alpha, x^\beta]_\star := x^\alpha \star x^\beta - x^\beta \star x^\alpha = i\theta^{\alpha\beta}. \quad (123)$$

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha + [A_\alpha, A_\beta]_\star \quad (124)$$

$$D_{z^a} \phi := \partial_{z^a} \phi + A_{+z^a} \star \phi - \phi \star A_{-z^a}. \quad (125)$$

$$-p_0 - 2\rho_1 = -s - 1. \quad (126)$$

$$g(u) = 1 + \sum_{n=1}^{\infty} g_n (u - u_0)^n. \quad (127)$$

$$y(u) = Ay_1(u) + B (g_s y_1(u) \log(u - u_0) + \tilde{y}_2(u)), \quad (128)$$

$$\tilde{y}_2(u) = (u - u_0)^{\rho_2} \left( -\frac{1}{s} + \sum_{n=1}^{\infty} h_n (u - u_0)^n \right), \quad (129)$$

$$y(u) = Ay_1(u) + B \left( y_1(u) \log(u - u_0) + (u - u_0)^{\rho_2} \sum_{n=1}^{\infty} h_n (u - u_0)^n \right). \quad (130)$$

$$\rho_1^{(\infty)} + \rho_2^{(\infty)} + \sum_{i=1}^n \left( \rho_1^{(i)} + \rho_2^{(i)} \right) = n - 1. \quad (131)$$

$$\begin{aligned} G(u) &= \left( \frac{1+u}{1-u} \right)^{\frac{1}{2}} \left( -i\sigma \cdot e_\mu \ i\sigma \cdot e_\mu \tilde{F}(u) \right), \\ &= \left( \frac{1+u}{1-u} \right)^{\frac{1}{2}} \tilde{F}(u), \end{aligned} \quad (132)$$

$$\tilde{F}(u) = \frac{1}{E-g} \left( -\frac{3(1-u)}{2\sqrt{1-u^2}} \tilde{G}(u) + \sqrt{1-u^2} \frac{d}{du} \tilde{G}(u) \right). \quad (133)$$

$$(1-u^2) (E+gu) \frac{d^2 G}{du^2} - ((4u+1) (E+gu) + g(1-u^2)) \frac{dG}{du} + \quad (134)$$

$$D_{z^a} \phi = \partial_{z^a} \phi + \mathcal{A}_{z^a} \star \phi. \quad (135)$$

$$D_{z^a} \phi = \partial_{z^a} \phi - \phi \star \mathcal{B}_{z^a} \quad (136)$$

$$G(u) = \sum_{n=0}^{\infty} a_n (u+1)^n. \quad (137)$$

$$a_n = \prod_{i=0}^{n-1} b_i. \quad (138)$$

$$\begin{aligned}
R_{\text{conv.}} &= \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|, \\
&= \lim_{n \rightarrow \infty} \frac{1}{|b_n|}, \\
&= 2.
\end{aligned} \tag{139}$$

$$G(1) = \sum_{n=0}^{\infty} a_n 2^n. \tag{140}$$

$$\begin{aligned}
y_1'(u) &= y_2(u), \\
y_2'(u) &= h(y_1(u), y_2(u), E).
\end{aligned} \tag{141}$$

$$\begin{aligned}
y_1'(u) &= y_2(u), \\
y_2'(u) &= h(y_1(u), y_2(u), y_3(u)), \\
y_3'(u) &= 0.
\end{aligned} \tag{142}$$

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_C \frac{f(z)dz}{(z-u)^{n+1}}, \tag{143}$$

$$G(u) = \sum_{k=0}^{\infty} a_k u^k, \tag{144}$$

$$G(u) = \frac{1}{\pi} \Re \left( \int_0^\pi \frac{G(e^{i\phi}) e^{i\phi} d\phi}{e^{i\phi} - u} \right), \tag{145}$$

$$Q = \int_{S^3} \bar{\psi} \psi. \tag{146}$$

$$Q = \int_0^\pi \tilde{Q}(\mu) \, \mathrm{d}\mu, \tag{147}$$

$$= 4\pi \int_0^\pi \left( (1 - \cos \mu) G^2 + (1 + \cos \mu) \tilde{G}^2 \right) \sin^2 \mu \, \mathrm{d}\mu. \tag{148}$$

$$\tilde{G}(1) = -\frac{3}{2(E+g)}. \tag{149}$$

$$\tilde{B}(\mu) = \frac{2}{\pi} f'(\mu) \sin^2 f(\mu). \quad (150)$$

$$\tilde{Q}(\mu) = 8\pi a_0^2 \sin^2 \mu. \quad (151)$$

$$\hat{D}_{z^a} \phi = \hat{\partial}_{z^a} \hat{\phi} + \hat{A}_{+z^a} \hat{\phi} - \hat{\phi} \hat{A}_{-z^a}. \quad (152)$$

$$\mathcal{T} = \sum_{i=\pm} (F_{i\mu\nu} * F_{i\mu\nu} + \nabla_{i\mu} \mathcal{J}_{i\mu}), \quad (153)$$

$$\nabla_{\pm\mu} \cdot := \partial_\mu \cdot + [A_{\pm\mu}, \cdot], \quad (154)$$

$$\begin{aligned} -(\Delta\alpha)_r &= \left( \partial_r^2 + \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_\theta^2 - \frac{1}{r^2} \right) \alpha_r - \frac{2}{r^3} \partial_\theta \alpha_\theta, \\ -(\Delta\alpha)_\theta &= \left( \partial_r^2 - \frac{1}{r} \partial_r + \frac{1}{r^2} \partial_\theta^2 \right) \alpha_\theta + \frac{2}{r} \partial_\theta \alpha_r. \end{aligned} \quad (155)$$

$$-\frac{1}{r}(\Delta\alpha)_\theta = -\frac{1}{r}\omega^2\alpha_\theta - 2e\left(\varphi^*\partial_r\phi + \varphi\partial_r\phi^*\right). \quad (156)$$

$$\partial_r \Re(\delta\chi_-)|_{\partial M} \sim \left[ -\frac{1}{r}\omega^2\alpha_\theta + e\Re(\phi^*\partial_r\varphi - \varphi\partial_r\phi^*) \right] |_{\partial M} = 0 \quad (157)$$

$$\begin{aligned} G_0(u) &= a_0, \\ G_n(u) &= \sum_{k=0}^n a_k (u+1)^k, \end{aligned}$$

$$\Psi = \Psi_0 + \Phi, \quad (158)$$

$$Q(f) = \oint \frac{dw}{2\pi i} f(w) J_{\mathrm{B}}(w), \quad C(f) = \oint \frac{dw}{2\pi i} f(w) c(w). \quad (159)$$

$$h_a(w) = \log \left( 1 + \frac{a}{2} \left( w + \frac{1}{w} \right)^2 \right), \quad (160)$$

$$\Phi' = e^{K(h_a)} \Phi, \quad (161)$$

$$K(f) = \oint \frac{dw}{2\pi i} f(w) \left( J_{\text{gh}}(w) - \frac{3}{2} w^{-1} \right). \quad (162)$$

$$K(h_a) = -\tilde{q}_0 \log(1 - Z(a))^2 - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} (q_{2n} + q_{-2n}) Z(a)^n, \quad (163)$$

$$b_0 \Phi = 0. \quad (164)$$

$$L(a) = (1+a)L_0 + \frac{a}{2}(L_2 + L_{-2}) + a(q_2 - q_{-2}) + 4aZ(a), \quad (165)$$

$$L(a) = (1+a)L'_0 + \frac{a}{2}(L'_2 + L'_{-2}) + 4aZ(a) - 1 - a. \quad (166)$$

$$L(a) = 2(1+a)l_0 + a(l_2 + l_{-2}) + 4aZ(a) - 4(1+a), \quad (167)$$

$$l_0 = \frac{1}{2}(L'_0 + 3), \quad l_{\pm 2} = \frac{1}{2}L'_{\pm 2}, \quad (168)$$

$$[l_0, l_{\pm 2}] = \mp l_{\pm 2}, \quad [l_2, l_{-2}] = 2l_0. \quad (169)$$

$$U(s, t, u) = \exp(s l_{-2}) \exp(t l_0) \exp(u l_2), \quad (170)$$

$$au^2 + 2(1+a)u + a = 0, \quad (171)$$

$$a e^t - 2asu - 2(1+a)s = 0. \quad (172)$$

$$a e^t \mp 2\sqrt{1+2a} s = 0. \quad (173)$$

$$U'(a)L(a)U'(a)^{-1} = \sqrt{1+2a} L_0, \quad (174)$$

$$f_a(w) = \left( \frac{w^2 + Z(a)}{Z(a)w^2 + 1} \right)^{\frac{1}{2}}. \quad (175)$$

$$U'(a)\phi(w)U'(a)^{-1} = \left( \frac{df_a(w)}{dw} \right)^h \phi(f_a(w)). \quad (176)$$

$$L(a) = \oint \frac{dw}{2\pi i} w e^{h_a(w)} T'(w) + 4Z(a) - 1 - a, \quad (177)$$