

Acoustic Biodiversity Mapping

A global location-level biodiversity metric from Xeno-canto recordings

Generated 2026-07-14

OBJECTIVE

Derive a single metric that classifies a geographic location as biodiversity 'good' or 'bad', using a global archive of wildlife sound recordings and their metadata.

DATA

- 759,767 recordings across 5 continents (Xeno-canto), each with 42 acoustic indices computed by a SLURM batch pipeline (compute_indice.py over Butterworth-filtered audio).
- Metadata flattened from 766,747 API records (species, coordinates, quality, device).
- Acoustic scores joined to metadata on recording id (>99.99% match).


METHOD

1. Aggregate all recordings into 0.1-degree (~11 km) grid cells -> 40,914 cells.
2. Per cell, compute species richness from recorded species (genus+species and the 'also' co-occurring-species field), corrected for sampling effort by Hurlbert rarefaction to 10 recordings (S_{rare10}).
3. Test whether the acoustic indices predict richness (validation against ground truth).
4. Label each well-sampled cell good / moderate / bad relative to its 10-degree latitude band (region-relative tertiles).



HEADLINE RESULT

- 11,088 cells had enough recordings (≥ 10) to score confidently.
- The acoustic indices did NOT predict species richness (best |Spearman rho| ~ 0.13). The defensible biodiversity metric is therefore effort-controlled species richness, not any acoustic index -- the indices are unreliable on targeted (non-soundscape) recordings that dominate the archive.
- Final labels: 3692 good, 3695 moderate, 3701 bad.

Region-relative biodiversity label per 0.1-degree cell (11,088 well-sampled cells)



Validation: does the audio carry the biodiversity signal?



The Metric & Results

METRIC DEFINITION

location = 0.1 deg x 0.1 deg grid cell (~11 km)
biodiversity value = S_{rare10} , the expected number of distinct recorded species in 10 recordings (Hurlbert sample-based rarefaction; controls for recording effort)
label = tertile of S_{rare10} WITHIN the cell's 10-deg latitude band: good = top third, bad = bottom third
scope = cells with ≥ 10 recordings (11,088 of 40,914 cells)

GLOBAL VALUE RANGE (S_{rare10})

min 1.0 p25 8.2 median 10.1 p75 13.3 p90 17.2 max 46

Median richness by continent (well-sampled cells)

Continent	Median richness	Scored cells
america	10.9	3415
europa	10.2	4643
australia	9.9	404
africa	9.7	1003
asia	9.3	1623



KEY LIMITATIONS (must accompany any use of this metric)

1. Acoustic indices are non-predictive here. ACI/ADI/NDISI etc. were built for passive soundscape monitoring; 69% of archive clips are single-target recordings (median 24 s), so per-clip indices do not reflect site diversity.
2. 'Richness' = RECORDED species, not true species. It reflects recordist effort and interest. Rarefaction controls sample SIZE but not observer bias.
3. Weak latitude gradient. Effort-controlled richness is nearly flat across latitude (thresholds ~10-13 everywhere) -- evidence the signal is effort-driven, not the true tropical biodiversity peak. Region-relative labelling is used so scores mean 'rich for its region'.
4. Coverage. Only 27% of cells are confidently scored; the rest are 'insufficient_data' (< 10 recordings).

OUTPUT FILES

grid_cells.csv	- per-cell features + richness (all 40,914 cells)
grid_cells_labeled_regional.csv	- region-relative good/moderate/bad labels
biodiversity_map.csv	- lat, lon, label, code -> ready for GIS/folium

Year-by-year (2015-2025): temporal slices of the metric



COVERAGE: 14,487 scored cell-years; 2697 cells scored in ≥ 2 years, 10 in all 11.

CAUTION: even an 11-year window cannot show real biodiversity change. These movements reflect WHICH cells were recorded and by WHOM each year (effort + observer turnover), not ecological gain or loss. The global change (9.7 $\rightarrow$ 9.3) tracks recording effort, not nature. Use year slices to study sampling coverage over time -- not as a biodiversity time series.

OUTPUTS: grid_cells_yearly.csv (per cell-year), cell_change.csv (per tracked cell).