

# Contents

<b>1 Definition</b>	<b>2</b>	<b>4 Tables</b>	<b>5</b>
1.1 Extensions to Angles $> 90^\circ$ . . . . .	2	4.1 Exact Values of Trigonometric Functions . . .	5
1.2 Degrees and Radians . . . . .	2	4.2 Relations Between Trig Functions . . . . .	6
1.3 Signs and Variations . . . . .	2	<b>5 Graphs</b>	<b>6</b>
<b>2 Properties</b>	<b>3</b>	5.1 $y = \sin x$ . . . . .	6
2.1 $\sin x$ . . . . .	3	5.2 $y = \cos x$ . . . . .	6
2.2 $\cos x$ . . . . .	3	5.3 $y = \tan x$ . . . . .	7
2.3 $\tan x$ . . . . .	3	<b>6 Inverse Trigonometric Functions</b>	<b>8</b>
2.4 $\cot x$ . . . . .	3	6.1 Principal Values . . . . .	8
<b>3 Identities</b>	<b>3</b>	6.2 Identities . . . . .	8
3.1 Basic Identities . . . . .	3	6.2.1 Reciprocal Identities . . . . .	8
3.1.1 Reciprocal Identities . . . . .	3	6.2.2 Parity Identities . . . . .	8
3.1.2 Ratio Identities . . . . .	3	6.2.3 Pythagorean Identities (sort of) . . . .	8
3.1.3 Pythagorean Identities . . . . .	3	6.2.4 Sum and Difference . . . . .	8
3.1.4 Co-function Identities . . . . .	3	<b>7 Relationships Between Sides and Angles</b>	<b>8</b>
3.1.5 Parity Identities . . . . .	3	7.1 Law of Sines . . . . .	8
3.2 Sum and Difference . . . . .	4	7.2 Law of Cosines . . . . .	8
3.3 Double Angle . . . . .	4	7.3 Law of Tangents . . . . .	8
3.4 Half Angle . . . . .	4	7.4 Law of Cotangents . . . . .	9
3.5 Multiple Angle . . . . .	4	7.5 Mollweide's Formula . . . . .	9
3.6 Power Reduction . . . . .	4	7.6 Stewart's Theorem . . . . .	9
3.7 Product to Sum . . . . .	4	<b>8 Solving Triangles</b>	<b>9</b>
3.8 Sum to Product . . . . .	4	8.1 AAA Triangle . . . . .	9
3.9 Other Related Identities . . . . .	4	8.2 AAS/ASA Triangle . . . . .	9
3.10 Identities without Variables . . . . .	4	8.3 SAS Triangle . . . . .	9
		8.4 SSS Triangle . . . . .	9
		8.5 SSA Triangle . . . . .	9

# 1 Definition

Triangle  $ABC$  has a right angle at  $C$  and sides of length  $a$ ,  $b$ ,  $c$ . The trigonometric functions of angle  $A$  are defined as follows:

1.  $\sin A = \frac{a}{c} = \frac{\text{opposite}}{\text{hypotenuse}}$
2.  $\cos A = \frac{b}{c} = \frac{\text{adjacent}}{\text{hypotenuse}}$
3.  $\tan A = \frac{a}{b} = \frac{\text{opposite}}{\text{adjacent}}$
4.  $\csc A = \frac{b}{a} = \frac{\text{hypotenuse}}{\text{opposite}}$
5.  $\sec A = \frac{c}{a} = \frac{\text{hypotenuse}}{\text{adjacent}}$
6.  $\cot A = \frac{b}{a} = \frac{\text{adjacent}}{\text{opposite}}$

## 1.3 Signs and Variations

Quadrant	$\sin A$	$\cos A$	$\tan A$
I	$\begin{matrix} + \\ (0, 1) \end{matrix}$	$\begin{matrix} + \\ (1, 0) \end{matrix}$	$\begin{matrix} + \\ (0, \infty) \end{matrix}$
II	$\begin{matrix} + \\ (1, 0) \end{matrix}$	$\begin{matrix} - \\ (0, -1) \end{matrix}$	$\begin{matrix} - \\ (-\infty, 0) \end{matrix}$
III	$\begin{matrix} - \\ (0, -1) \end{matrix}$	$\begin{matrix} - \\ (-1, 0) \end{matrix}$	$\begin{matrix} + \\ (0, \infty) \end{matrix}$
IV	$\begin{matrix} - \\ (-1, 0) \end{matrix}$	$\begin{matrix} + \\ (0, 1) \end{matrix}$	$\begin{matrix} - \\ (-\infty, 0) \end{matrix}$

Quadrant	$\cot A$	$\sec A$	$\csc A$
I	$\begin{matrix} + \\ (\infty, 0) \end{matrix}$	$\begin{matrix} + \\ (1, \infty) \end{matrix}$	$\begin{matrix} + \\ (\infty, 1) \end{matrix}$
II	$\begin{matrix} - \\ (0, -\infty) \end{matrix}$	$\begin{matrix} - \\ (\infty, -1) \end{matrix}$	$\begin{matrix} + \\ (1, \infty) \end{matrix}$
III	$\begin{matrix} + \\ (\infty, 0) \end{matrix}$	$\begin{matrix} - \\ (-1, \infty) \end{matrix}$	$\begin{matrix} - \\ (\infty, -1) \end{matrix}$
IV	$\begin{matrix} - \\ (0, -\infty) \end{matrix}$	$\begin{matrix} + \\ (\infty, 1) \end{matrix}$	$\begin{matrix} - \\ (-1, \infty) \end{matrix}$

### 1.1 Extensions to Angles $> 90^\circ$

A point  $P$  in the Cartesian plane has coordinates  $(x, y)$ , where  $x$  is considered as positive along  $OX$  and negative along  $OX'$ , while  $y$  is considered as positive along  $OY'$  and negative along  $OY$ . The distance from origin  $O$  to point  $P$  is positive and denoted by  $r = \sqrt{x^2 + y^2}$ . The angle  $A$  described *counterclockwise* from  $OX$  is considered *positive*. If it is described *clockwise* from  $OX$  it is considered *negative*.

For an angle  $A$  in any quadrant, the trigonometric functions of  $A$  are defined as follows:

1.  $\sin A = \frac{y}{r}$
2.  $\cos A = \frac{x}{r}$
3.  $\tan A = \frac{y}{x}$
4.  $\csc A = \frac{r}{y}$
5.  $\sec A = \frac{r}{x}$
6.  $\cot A = \frac{x}{y}$

### 1.2 Degrees and Radians

A *radian* is that angle  $\theta$  subtended at center  $O$  of a circle by an arc  $MN$  equal to the radius  $r$ . Since  $2\pi$  radians  $= 360^\circ$  we have:

$$1 \text{ radian} = 180^\circ / \pi = 57.29577951308232 \dots^\circ$$

$$1^\circ = \pi / 180 \text{ radians} = 0.017453292519943 \dots \text{ radians}$$

## 2 Properties

### 2.1 $\sin x$

**Domain:**  $\{x|x \in \mathbb{R}\}$

**Range:**  $\{y|-1 \leq y \leq 1\}$

**Period:**  $2\pi$

**VA:** none

**$x$ -intercepts:**  $k\pi$  where  $k \in \mathbb{Z}$

**Parity:** odd

### 2.2 $\cos x$

**Domain:**  $\{x|x \in \mathbb{R}\}$

**Range:**  $\{y|-1 \leq y \leq 1\}$

**Period:**  $2\pi$

**VA:** none

**$x$ -intercepts:**  $\frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**Parity:** even

### 2.3 $\tan x$

**Domain:**  $\{x|x \neq \frac{\pi}{2} + k\pi, k \in \mathbb{Z}\}$

**Range:**  $\{y|y \in \mathbb{R}\}$

**Period:**  $\pi$

**VA:**  $x = \frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:** midway between asymptotes

**Parity:** odd

### 2.4 $\cot x$

**Domain:**  $\{x|x \neq k\pi, k \in \mathbb{Z}\}$

**Range:**  $\{y|y \in \mathbb{R}\}$

**Period:**  $\pi$

**VA:**  $x = k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:** midway between asymptotes

**Parity:** odd

## 3 Identities

### 3.1 Basic Identities

#### 3.1.1 Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta}; \sin \theta = \frac{1}{\csc \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}; \cos \theta = \frac{1}{\sec \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}; \tan \theta = \frac{1}{\cot \theta}$$

$$\sin \theta \csc \theta = \cos \theta \sec \theta = \tan \theta \cot \theta = 1$$

#### 3.1.2 Ratio Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}; \cos \theta = \frac{\sin \theta}{\tan \theta}; \sin \theta = \cos \theta \tan \theta$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}; \sin \theta = \frac{\cos \theta}{\cot \theta}; \cos \theta = \sin \theta \cot \theta$$

#### 3.1.3 Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1; \sin^2 \theta = 1 - \cos^2 \theta; \cos^2 \theta = 1 - \sin^2 \theta$$

$$\tan^2 \theta + 1 = \sec^2 \theta; \tan^2 \theta = \sec^2 \theta - 1; \sec^2 \theta - \tan^2 \theta = 1$$

$$\cot^2 \theta + 1 = \csc^2 \theta; \cot^2 \theta = \csc^2 \theta - 1; \csc^2 \theta - \cot^2 \theta = 1$$

#### 3.1.4 Co-function Identities

$$\sin \left( \frac{\pi}{2} - \theta \right) = \cos \theta$$

$$\cos \left( \frac{\pi}{2} - \theta \right) = \sin \theta$$

$$\tan \left( \frac{\pi}{2} - \theta \right) = \cot \theta$$

$$\csc \left( \frac{\pi}{2} - \theta \right) = \sec \theta$$

$$\sec \left( \frac{\pi}{2} - \theta \right) = \csc \theta$$

$$\cot \left( \frac{\pi}{2} - \theta \right) = \tan \theta$$

#### 3.1.5 Parity Identities

$$\sin(-A) = -\sin A$$

$$\cos(-A) = \cos A$$

$$\tan(-A) = -\tan A$$

$$\csc(-A) = -\csc A$$

$$\sec(-A) = \sec A$$

$$\cot(-A) = -\cot A$$

### 3.2 Sum and Difference

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

$$\cot(\alpha \pm \beta) = \frac{\cot \alpha \cot \beta \mp 1}{\cot \beta \pm \cot \alpha}$$

### 3.3 Double Angle

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 1 - 2 \sin^2 \alpha = 2 \cos^2 \alpha - 1$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$$

### 3.4 Half Angle

Let  $\mathcal{Q}_n$ , where  $n \in \{1, 2, 3, 4\}$ , denote the set of all points within the  $n$ th quadrant of the Cartesian plane.

$$\sin \frac{\alpha}{2} = \begin{cases} \sqrt{\frac{1 - \cos \alpha}{2}} & \frac{\alpha}{2} \in (\mathcal{Q}_1 \cup \mathcal{Q}_2) \\ -\sqrt{\frac{1 - \cos \alpha}{2}} & \frac{\alpha}{2} \in (\mathcal{Q}_3 \cup \mathcal{Q}_4) \end{cases}$$

$$\cos \frac{\alpha}{2} = \begin{cases} \sqrt{\frac{1 + \cos \alpha}{2}} & \frac{\alpha}{2} \in (\mathcal{Q}_1 \cup \mathcal{Q}_4) \\ -\sqrt{\frac{1 + \cos \alpha}{2}} & \frac{\alpha}{2} \in (\mathcal{Q}_2 \cup \mathcal{Q}_3) \end{cases}$$

$$\tan \frac{\alpha}{2} = \frac{\sin \alpha}{1 + \cos \alpha} = \frac{1 - \cos \alpha}{\sin \alpha} = \csc \alpha - \cot \alpha$$

### 3.5 Multiple Angle

$$\sin 3\alpha = 3 \sin \alpha - 4 \sin^3 \alpha$$

$$\cos 3\alpha = 4 \cos^3 \alpha - 3 \cos \alpha$$

$$\tan 3\alpha = \frac{3 \tan \alpha - \tan^3 \alpha}{1 - 3 \tan^2 \alpha}$$

$$\sin 4\alpha = 4 \sin \alpha \cos \alpha - 8 \sin^3 \alpha \cos \alpha$$

$$\cos 4\alpha = 8 \cos^4 \alpha - 8 \cos^2 \alpha + 1$$

$$\tan 4\alpha = \frac{4 \tan \alpha - 4 \tan^3 \alpha}{1 - 6 \tan^2 \alpha + \tan^4 \alpha}$$

$$\sin(n\alpha) = \sum_{i=0}^n \binom{n}{i} \cos^i \alpha \sin^{n-i} \alpha \sin\left(\frac{(n-i)\pi}{2}\right)$$

$$\cos(n\alpha) = \sum_{i=0}^n \binom{n}{i} \cos^i \alpha \sin^{n-i} \alpha \cos\left(\frac{(n-i)\pi}{2}\right)$$

### 3.6 Power Reduction

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

$$\tan^2 \theta = \frac{1 - \cos 2\theta}{1 + \cos 2\theta}$$

$$\sin^3 \theta = \frac{3 \sin \theta - \sin 3\theta}{4}$$

$$\cos^3 \theta = \frac{3 \cos \theta + \cos 3\theta}{4}$$

$$\sin^4 \theta = \frac{3 - 4 \cos 2\theta + \cos 4\theta}{8}$$

$$\cos^4 \theta = \frac{3 + 4 \cos 2\theta + \cos 4\theta}{8}$$

$$\sin^5 \theta = \frac{10 \sin \theta - 5 \sin 3\theta + \sin 5\theta}{16}$$

$$\cos^5 \theta = \frac{10 \cos \theta + 5 \cos 3\theta + \cos 5\theta}{16}$$

### 3.7 Product to Sum

$$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\cos \alpha \sin \beta = \frac{1}{2} [\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2} [\cos(\alpha + \beta) + \cos(\alpha - \beta)]$$

$$\sin \alpha \sin \beta = \frac{1}{2} [\cos(\alpha + \beta) - \cos(\alpha - \beta)]$$

### 3.8 Sum to Product

$$\sin \theta \pm \sin \varphi = 2 \sin \frac{\theta \pm \varphi}{2} \cos \frac{\theta \mp \varphi}{2}$$

$$\cos \theta \pm \cos \varphi = \pm 2 \sin \frac{\theta + \varphi}{2} \cos \frac{\theta - \varphi}{2}$$

### 3.9 Other Related Identities

- If  $x + y + z = \pi$ , then  $\sin 2x + \sin 2y + \sin 2z = 4 \sin x \sin y \sin z$ .
- *Triple Tangent Identity.* If  $x + y + z = \pi$ , then  $\tan x + \tan y + \tan z = \tan x \tan y \tan z$ .
- *Triple Cotangent Identity.* If  $x + y + z = \frac{\pi}{2}$ , then  $\cot x + \cot y + \cot z = \cot x \cot y \cot z$ .

### 3.10 Identities without Variables

$$\cos 20^\circ \cdot \cos 40^\circ \cdot \cos 80^\circ = \frac{1}{8}$$

$$\sin 20^\circ \cdot \sin 40^\circ \cdot \sin 80^\circ = \frac{\sqrt{3}}{8}$$

$$\cos 24^\circ + \cos 48^\circ + \cos 96^\circ + \cos 168^\circ = \frac{1}{2}$$

## 4 Tables

### 4.1 Exact Values of Trigonometric Functions


$A^\circ$	$A$ rad	$\sin A$	$\cos A$	$\tan A$	$\cot A$	$\sec A$	$\csc A$
$0^\circ$	0	0	1	0	$\infty$	1	$\infty$
$15^\circ$	$\pi/12$	$\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$2 - \sqrt{3}$	$2 + \sqrt{3}$	$\sqrt{6} - \sqrt{2}$	$\sqrt{6} + \sqrt{2}$
$30^\circ$	$\pi/6$	$\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	2
$45^\circ$	$\pi/4$	$\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
$60^\circ$	$\pi/3$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	2	$\frac{2}{3}\sqrt{3}$
$75^\circ$	$5\pi/12$	$\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$2 + \sqrt{3}$	$2 - \sqrt{3}$	$\sqrt{6} + \sqrt{2}$	$\sqrt{6} - \sqrt{2}$
$90^\circ$	$\pi/2$	1	0	$\pm\infty$	0	$\pm\infty$	1
$105^\circ$	$7\pi/12$	$\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$-\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$-(2 + \sqrt{3})$	$-(2 - \sqrt{3})$	$-(\sqrt{6} + \sqrt{2})$	$\sqrt{6} - \sqrt{2}$
$120^\circ$	$2\pi/3$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	-2	$\frac{2}{3}\sqrt{3}$
$135^\circ$	$3\pi/4$	$\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	-1	-1	$-\sqrt{2}$	$\sqrt{2}$
$150^\circ$	$5\pi/6$	$\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	2
$165^\circ$	$11\pi/12$	$\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$-\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$-(2 - \sqrt{3})$	$-(2 + \sqrt{3})$	$-(\sqrt{6} - \sqrt{2})$	$\sqrt{6} + \sqrt{2}$
$180^\circ$	$\pi$	0	-1	0	$\mp\infty$	-1	$\pm\infty$
$195^\circ$	$13\pi/12$	$-\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$-\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$2 - \sqrt{3}$	$2 + \sqrt{3}$	$-(\sqrt{6} - \sqrt{2})$	$-(\sqrt{6} + \sqrt{2})$
$210^\circ$	$7\pi/6$	$\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	-2
$225^\circ$	$5\pi/4$	$-\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	1	1	$-\sqrt{2}$	$-\sqrt{2}$
$240^\circ$	$4\pi/3$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	-2	$-\frac{2}{3}\sqrt{3}$
$255^\circ$	$17\pi/12$	$-\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$-\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$2 + \sqrt{3}$	$2 - \sqrt{3}$	$-(\sqrt{6} + \sqrt{2})$	$-(\sqrt{6} - \sqrt{2})$
$270^\circ$	$3\pi/2$	-1	0	$\pm\infty$	0	$\mp\infty$	-1
$285^\circ$	$19\pi/12$	$-\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$-(2 + \sqrt{3})$	$-(2 - \sqrt{3})$	$\sqrt{6} + \sqrt{2}$	$-(\sqrt{6} - \sqrt{2})$
$300^\circ$	$5\pi/3$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	2	$-\frac{2}{3}\sqrt{3}$
$315^\circ$	$7\pi/4$	$-\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	-1	-1	$\sqrt{2}$	$-\sqrt{2}$
$330^\circ$	$11\pi/6$	$-\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	-2
$345^\circ$	$23\pi/12$	$-\frac{1}{4}(\sqrt{6} - \sqrt{2})$	$\frac{1}{4}(\sqrt{6} + \sqrt{2})$	$-(2 + \sqrt{3})$	$-(2 - \sqrt{3})$	$\sqrt{6} - \sqrt{2}$	$-(\sqrt{6} + \sqrt{2})$
$360^\circ$	$2\pi$	0	1	0	$\mp\infty$	1	$\mp\infty$

## 4.2 Relations Between Trig Functions


	$\sin \theta = u$	$\cos \theta = u$	$\tan \theta = u$	$\csc \theta = u$	$\sec \theta = u$	$\cot \theta = u$
$\sin \theta$	$u$	$\sqrt{1-u^2}$	$\frac{u}{\sqrt{1+u^2}}$	$\frac{1}{u}$	$\frac{\sqrt{u^2-1}}{u}$	$\frac{1}{\sqrt{1+u^2}}$
$\cos \theta$	$\sqrt{1-u^2}$	$u$	$\frac{1}{\sqrt{1+u^2}}$	$\frac{\sqrt{u^2-1}}{u}$	$\frac{1}{u}$	$\frac{u}{\sqrt{1+u^2}}$
$\tan \theta$	$\frac{u}{\sqrt{1-u^2}}$	$\frac{\sqrt{1-u^2}}{u}$	$u$	$\frac{1}{\sqrt{u^2-1}}$	$\sqrt{u^2-1}$	$\frac{1}{u}$
$\csc \theta$	$\frac{1}{u}$	$\frac{1}{\sqrt{1-u^2}}$	$\frac{\sqrt{1+u^2}}{u}$	$u$	$\frac{u}{\sqrt{u^2-1}}$	$\sqrt{1+u^2}$
$\sec \theta$	$\frac{1}{\sqrt{1-u^2}}$	$\frac{1}{u}$	$\sqrt{1+u^2}$	$\frac{u}{\sqrt{u^2-1}}$	$u$	$\frac{\sqrt{1+u^2}}{u}$
$\cot \theta$	$\frac{\sqrt{1-u^2}}{u}$	$\frac{u}{\sqrt{1-u^2}}$	$\frac{1}{u}$	$\sqrt{u^2-1}$	$\frac{1}{\sqrt{u^2-1}}$	$u$


## 5 Graphs

### 5.1 $y = \sin x$



### 5.2 $y = \cos x$



**5.3**  $y = \tan x$ 

## 6 Inverse Trigonometric Functions

If  $x = \sin y$ , then  $y = \sin^{-1} x$ , i.e. the angle whose sine is  $x$  or inverse sine of  $x$ , is a multiple-valued function of  $x$  which is a collection of single-valued functions called *branches*. Similarly, the other inverse trigonometric functions are multiple-valued.

For many purposes, a particular branch is required. This is called the *principal branch* and the values for this branch are called *principal values*.

### 6.1 Principal Values

Principal values for $x \geq 0$	Principal values for $x < 0$
$0 \leq \sin^{-1} x \leq \frac{\pi}{2}$	$-\frac{\pi}{2} \leq \sin^{-1} x < 0$
$0 \leq \cos^{-1} x \leq \frac{\pi}{2}$	$\frac{\pi}{2} < \cos^{-1} x \leq \pi$
$0 \leq \tan^{-1} x < \frac{\pi}{2}$	$-\frac{\pi}{2} < \tan^{-1} x < 0$
$0 < \cot^{-1} x \leq \frac{\pi}{2}$	$\frac{\pi}{2} < \cot^{-1} x < \pi$
$0 \leq \sec^{-1} x < \frac{\pi}{2}$	$\frac{\pi}{2} < \sec^{-1} x \leq \pi$
$0 < \csc^{-1} x \leq \frac{\pi}{2}$	$-\frac{\pi}{2} \leq \csc^{-1} x < 0$

### 6.2 Identities

In all cases it is assumed that principal values are used.

#### 6.2.1 Reciprocal Identities

$$\csc^{-1} x = \sin^{-1} \left( \frac{1}{x} \right)$$

$$\sec^{-1} x = \cos^{-1} \left( \frac{1}{x} \right)$$

$$\cot^{-1} x = \tan^{-1} \left( \frac{1}{x} \right)$$

#### 6.2.2 Parity Identities

$$\sin^{-1}(-x) = -\sin^{-1} x$$

$$\cos^{-1}(-x) = \pi - \cos^{-1} x$$

$$\tan^{-1}(-x) = -\tan^{-1} x$$

$$\csc^{-1}(-x) = -\csc^{-1} x$$

$$\sec^{-1}(-x) = \pi - \sec^{-1} x$$

$$\cot^{-1}(-x) = \pi - \cot^{-1} x$$

#### 6.2.3 Pythagorean Identities (sort of)

$$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$$

$$\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$$

$$\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}$$

#### 6.2.4 Sum and Difference

$$\sin^{-1} \alpha \pm \sin^{-1} \beta = \sin^{-1} \left( \alpha \sqrt{1 - \beta^2} \pm \beta \sqrt{1 - \alpha^2} \right)$$

$$\cos^{-1} \alpha \pm \cos^{-1} \beta = \cos^{-1} \left( \alpha \beta \mp \sqrt{(1 - \alpha^2)(1 - \beta^2)} \right)$$


$$\tan^{-1} \alpha \pm \tan^{-1} \beta = \tan^{-1} \left( \frac{\alpha \pm \beta}{1 \mp \alpha \beta} \right)$$

## 7 Relationships Between Sides and Angles

### 7.1 Law of Sines

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$$

#### Extended Law of Sines



$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma} = 2R,$$

where  $R$  is the circumradius of the triangle.

### 7.2 Law of Cosines

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}; \quad a = \sqrt{b^2 + c^2 - 2bc \cos \alpha}$$

$$\cos \beta = \frac{a^2 + c^2 - b^2}{2ac}; \quad b = \sqrt{a^2 + c^2 - 2ac \cos \beta}$$

$$\cos \gamma = \frac{a^2 + b^2 - c^2}{2ab}; \quad c = \sqrt{a^2 + b^2 - 2ab \cos \gamma}$$

### 7.3 Law of Tangents

$$\frac{a - b}{a + b} = \frac{\tan \frac{1}{2}(\alpha - \beta)}{\tan \frac{1}{2}(\alpha + \beta)}$$

$$\frac{b - c}{b + c} = \frac{\tan \frac{1}{2}(\beta - \gamma)}{\tan \frac{1}{2}(\beta + \gamma)}$$

$$\frac{c - a}{c + a} = \frac{\tan \frac{1}{2}(\gamma - \alpha)}{\tan \frac{1}{2}(\gamma + \alpha)}$$



## 7.4 Law of Cotangents

Let  $s$  be the semi-perimeter, that is,  $s = \frac{(a+b+c)}{2}$ , and  $r$  be the radius of the inscribed circle, then

$$\frac{\cot(\frac{\alpha}{2})}{s-a} = \frac{\cot(\frac{\beta}{2})}{s-b} = \frac{\cot(\frac{\gamma}{2})}{s-c} = \frac{1}{r}$$

and furthermore that the inradius is given by  $r = \sqrt{\frac{(s-a)(s-b)(s-c)}{s}}$ .


## 7.5 Mollweide's Formula

Each of these identities uses all six parts of the triangle—the three angles and the lengths of the three sides.

$$\frac{a+b}{c} = \frac{\cos\left(\frac{\alpha-\beta}{2}\right)}{\sin\left(\frac{\gamma}{2}\right)}$$

$$\frac{a-b}{c} = \frac{\sin\left(\frac{\alpha-\beta}{2}\right)}{\cos\left(\frac{\gamma}{2}\right)}$$

## 7.6 Stewart's Theorem



Let  $D$  be a point in  $\overline{BC}$  of  $\triangle ABC$ . If  $|BD| = m$ ,  $|CD| = n$ , and  $|AD| = d$ , then  $b^2m + c^2n = a(d^2 + mn)$ .

# 8 Solving Triangles

## 8.1 AAA Triangle

1. Write “no solution” as your answer.

## 8.2 AAS/ASA Triangle

1. Solve for the missing angle.
2. Use the Law of Sines twice for the other two sides.

## 8.3 SAS Triangle

1. Use the Law of Cosines for the other non-included angle.
2. Use the Law of Sines for the missing side.
3. Solve for the missing angle.

## 8.4 SSS Triangle

1. Use the Law of Cosines twice for the two angles.
2. Solve for the missing angle.

## 8.5 SSA Triangle

When  $\alpha$  is acute

$0 < a < h$	$a = h$	$a > b$	$h < a < b$
0	1	1	2

When  $\alpha$  is obtuse

$0 < a \leq b$	$a > b$
0	1