


1 Definition

Triangle  $ABC$  has a right angle at  $C$  and sides of length  $a$ ,  $b$ ,  $c$ . The trigonometric functions of angle  $A$  are defined as follows:

- 1.  $\sin A = \frac{a}{c} = \frac{\text{opposite}}{\text{hypotenuse}}$
- 2.  $\cos A = \frac{b}{c} = \frac{\text{adjacent}}{\text{hypotenuse}}$
- 3.  $\tan A = \frac{a}{b} = \frac{\text{opposite}}{\text{adjacent}}$
- 4.  $\csc A = \frac{c}{a} = \frac{\text{hypotenuse}}{\text{opposite}}$
- 5.  $\sec A = \frac{c}{b} = \frac{\text{hypotenuse}}{\text{adjacent}}$
- 6.  $\cot A = \frac{b}{a} = \frac{\text{adjacent}}{\text{opposite}}$

1.2 The Unit Circle



1.3 Degrees and Radians

A *radian* is that angle  $\theta$  subtended at center  $O$  of a circle by an arc  $MN$  equal to the radius  $r$ . Since  $2\pi$  radians  $= 360^\circ$  we have:

1 radian  $= 180^\circ / \pi = 57.29577951308232 \dots^\circ$   
1 $^\circ = \pi / 180$  radians  $= 0.017453292519943 \dots$  radians

1.4 Signs and Variations

Quadrant	$\sin A$	$\cos A$	$\tan A$
I	$\frac{+}{(0,1)}$	$\frac{+}{(1,0)}$	$\frac{+}{(0,\infty)}$
II	$\frac{+}{(1,0)}$	$\frac{-}{(0,-1)}$	$\frac{-}{(-\infty,0)}$
III	$\frac{-}{(0,-1)}$	$\frac{-}{(-1,0)}$	$\frac{+}{(0,\infty)}$
IV	$\frac{-}{(-1,0)}$	$\frac{+}{(0,1)}$	$\frac{-}{(-\infty,0)}$

Quadrant	$\cot A$	$\sec A$	$\csc A$
I	$\frac{+}{(\infty,0)}$	$\frac{+}{(1,\infty)}$	$\frac{+}{(\infty,1)}$
II	$\frac{-}{(0,-\infty)}$	$\frac{-}{(\infty,-1)}$	$\frac{+}{(1,\infty)}$
III	$\frac{+}{(\infty,0)}$	$\frac{-}{(-1,\infty)}$	$\frac{-}{(\infty,-1)}$
IV	$\frac{-}{(0,-\infty)}$	$\frac{+}{(\infty,1)}$	$\frac{-}{(-1,-\infty)}$

Trigonometry Cram Sheet

November 23, 2015

Contents

1	Definition	2
1.1	Extensions to Angles $> 90^\circ$	2
1.2	The Unit Circle	2
1.3	Degrees and Radians	2
1.4	Signs and Variations	2
2	Properties and General Forms	3
2.1	Properties	3
2.1.1	$\sin x$	3
2.1.2	$\cos x$	3
2.1.3	$\tan x$	3
2.1.4	$\csc x$	3
2.1.5	$\sec x$	3
2.1.6	$\cot x$	3
2.2	General Forms of Trigonometric Functions	3
3	Identities	4
3.1	Basic Identities	4
3.2	Sum and Difference	4
3.3	Double Angle	4
3.4	Half Angle	4
3.5	Multiple Angle	4
3.6	Power Reduction	5
3.7	Product to Sum	5
3.8	Sum to Product	5
3.9	Linear Combinations	5
3.10	Other Related Identities	5
3.11	Identities without Variables	6
4	Graphs	6
4.1	$y = \sin x$	6
4.2	$y = \cos x$	6
4.3	$y = \tan x$	6
4.4	$y = \csc x$	6
4.5	$y = \sec x$	6
4.6	$y = \cot x$	6
5	Tables	7
5.1	Exact Values of Trigonometric Functions	7
5.2	Relations Between Trig Functions	8
6	Inverse Trigonometric Functions	8
6.1	Principal Values	8
6.2	Identities	8
7	Relationships Between Sides and Angles	9
7.1	Law of Sines	9
7.2	Law of Cosines	9
7.3	Law of Tangents	9
7.4	Law of Cotangents	9
7.5	Mollweide's Formula	9
7.6	Stewart's Theorem	9
7.7	Angles in Terms of Sides	9
8	Solving Triangles	10
8.1	AAS/ASA Triangle	10
8.2	SAS Triangle	10
8.3	SSS Triangle	10
8.4	SSA Triangle	11
8.5	Right Triangle	11
9	Polar Coordinates	11
9.1	Properties	11
9.2	Coordinate Transformation	11
10	Special Polar Graphs	11
10.1	Limaçon of Pascal	12
10.2	Rose	13
10.3	Spiral of Archimedes	13
10.4	Lemniscate of Bernoulli	13
10.5	Folium of Descartes	14
10.6	Spiral of Fermat	14
10.7	Cissoid of Diocles	14
10.8	Epispiral	14
10.9	Lituus	15
10.10	Eight Curve	15
10.11	Butterfly Curve	15
10.12	Strophoid	15
10.13	Cochleoid	16
10.14	Cycloid of Ceva	16
10.15	Freeth's Nephroid	16
11	Miscellaneous Stuff	17
11.1	Pythagorean Triples	17
11.2	Triangle Centers	17
11.3	Area of the Triangle	18

## 2 Properties and General Forms

### 2.1 Properties

#### 2.1.1 $\sin x$

**Domain:**  $\{x|x \in \mathbb{R}\}$  or  $(-\infty, +\infty)$

**Range:**  $\{y| -1 \leq y \leq 1\}$  or  $[-1, 1]$

**Period:**  $2\pi$

**VA:** none

**$x$ -intercepts:**  $k\pi$  where  $k \in \mathbb{Z}$

**Parity:** odd

#### 2.1.2 $\cos x$

**Domain:**  $\{x|x \in \mathbb{R}\}$  or  $(-\infty, +\infty)$

**Range:**  $\{y| -1 \leq y \leq 1\}$  or  $[-1, 1]$

**Period:**  $2\pi$

**VA:** none

**$x$ -intercepts:**  $\frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**Parity:** even

#### 2.1.3 $\tan x$

**Domain:**  $\{x|x \neq \frac{\pi}{2} + k\pi, k \in \mathbb{Z}\}$  or  $\bigcup_{k \in \mathbb{Z}} \left( \frac{(k-1)\pi}{2}, \frac{(k+1)\pi}{2} \right)$

**Range:**  $\{y|y \in \mathbb{R}\}$  or  $(-\infty, +\infty)$

**Period:**  $\pi$

**VA:**  $x = \frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:**  $k\pi$  where  $k \in \mathbb{Z}$

**Parity:** odd

#### 2.1.4 $\csc x$

**Domain:**  $\{x|x \neq k\pi, k \in \mathbb{Z}\}$  or  $\bigcup_{k \in \mathbb{Z}} (k\pi, (k+1)\pi)$

**Range:**  $\{y|y \leq -1 \cup y \geq 1\}$  or  $(-\infty, -1] \cup [1, +\infty)$

**Period:**  $2\pi$

**VA:**  $x = k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:** none

**Parity:** odd

#### 2.1.5 $\sec x$

**Domain:**  $\{x|x \neq \frac{\pi}{2} + k\pi, k \in \mathbb{Z}\}$  or  $\bigcup_{k \in \mathbb{Z}} \left( \frac{(k-1)\pi}{2}, \frac{(k+1)\pi}{2} \right)$

**Range:**  $\{y|y \leq -1 \cup y \geq 1\}$  or  $(-\infty, -1] \cup [1, +\infty)$

**Period:**  $2\pi$

**VA:**  $x = \frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:** none

**Parity:** even

#### 2.1.6 $\cot x$

**Domain:**  $\{x|x \neq k\pi, k \in \mathbb{Z}\}$  or  $\bigcup_{k \in \mathbb{Z}} (k\pi, (k+1)\pi)$

**Range:**  $\{y|y \in \mathbb{R}\}$  or  $(-\infty, +\infty)$

**Period:**  $\pi$


**VA:**  $x = k\pi$  where  $k \in \mathbb{Z}$

**$x$ -intercepts:**  $\frac{\pi}{2} + k\pi$  where  $k \in \mathbb{Z}$

**Parity:** odd

## 2.2 General Forms of Trigonometric Functions

Given some trigonometric function  $f(x)$ , its general form is represented as  $y = Af(B(x - C)) + D$ , where its amplitude is  $|A|$ , its period is  $\frac{2\pi}{|B|}$  or  $\frac{\pi}{|B|}$  (for tangent and cotangent), its phase shift is  $C$ , and its vertical translation is  $D$  units upward (if  $D > 0$ ) or  $D$  units downward (if  $D < 0$ ). The maximum and minimum value for  $\sin x$  and  $\cos x$  is  $A + D$  and  $-A + D$  respectively.



## 3 Identities

### 3.1 Basic Identities

#### Reciprocal Identities

$$\csc \theta = \frac{1}{\sin \theta}; \quad \sin \theta = \frac{1}{\csc \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}; \quad \cos \theta = \frac{1}{\sec \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}; \quad \tan \theta = \frac{1}{\cot \theta}$$

$$\sin \theta \csc \theta = \cos \theta \sec \theta = \tan \theta \cot \theta = 1$$

#### Ratio Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}; \quad \cos \theta = \frac{\sin \theta}{\tan \theta}; \quad \sin \theta = \cos \theta \tan \theta$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}; \quad \sin \theta = \frac{\cos \theta}{\cot \theta}; \quad \cos \theta = \sin \theta \cot \theta$$

#### Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1; \quad \sin^2 \theta = 1 - \cos^2 \theta; \quad \cos^2 \theta = 1 - \sin^2 \theta$$

$$\tan^2 \theta + 1 = \sec^2 \theta; \quad \tan^2 \theta = \sec^2 \theta - 1; \quad \sec^2 \theta - \tan^2 \theta = 1$$

$$\cot^2 \theta + 1 = \csc^2 \theta; \quad \cot^2 \theta = \csc^2 \theta - 1; \quad \csc^2 \theta - \cot^2 \theta = 1$$

#### Co-function Identities

$$\sin\left(\frac{\pi}{2} - \theta\right) = \cos \theta$$

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin \theta$$

$$\tan\left(\frac{\pi}{2} - \theta\right) = \cot \theta$$

$$\csc\left(\frac{\pi}{2} - \theta\right) = \sec \theta$$

$$\sec\left(\frac{\pi}{2} - \theta\right) = \csc \theta$$

$$\cot\left(\frac{\pi}{2} - \theta\right) = \tan \theta$$

#### Parity Identities

$$\sin(-A) = -\sin A$$

$$\cos(-A) = \cos A$$

$$\tan(-A) = -\tan A$$

$$\csc(-A) = -\csc A$$

$$\sec(-A) = \sec A$$

$$\cot(-A) = -\cot A$$

### 3.2 Sum and Difference

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

$$\cot(\alpha \pm \beta) = \frac{\cot \alpha \cot \beta \mp 1}{\cot \beta \pm \cot \alpha}$$

### 3.3 Double Angle

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha$$

$$\cos 2\alpha = \cos^2 \alpha - \sin^2 \alpha = 1 - 2 \sin^2 \alpha = 2 \cos^2 \alpha - 1$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha}$$

### 3.4 Half Angle

Let  $Q_n$ , where  $n \in \{1, 2, 3, 4\}$ , denote the set of all angles within the  $n^{\text{th}}$  quadrant of the Cartesian plane.

$$\sin \frac{\alpha}{2} = \begin{cases} \sqrt{\frac{1 - \cos \alpha}{2}} & \text{if } \frac{\alpha}{2} \in (Q_1 \cup Q_2) \\ -\sqrt{\frac{1 - \cos \alpha}{2}} & \text{if } \frac{\alpha}{2} \in (Q_3 \cup Q_4) \end{cases}$$

$$\cos \frac{\alpha}{2} = \begin{cases} \sqrt{\frac{1 + \cos \alpha}{2}} & \text{if } \frac{\alpha}{2} \in (Q_1 \cup Q_4) \\ -\sqrt{\frac{1 + \cos \alpha}{2}} & \text{if } \frac{\alpha}{2} \in (Q_2 \cup Q_3) \end{cases}$$

$$\tan \frac{\alpha}{2} = \frac{\sin \alpha}{1 + \cos \alpha} = \frac{1 - \cos \alpha}{\sin \alpha} = \csc \alpha - \cot \alpha$$

### 3.5 Multiple Angle

$$\sin 3\alpha = 3 \sin \alpha - 4 \sin^3 \alpha$$

$$\cos 3\alpha = 4 \cos^3 \alpha - 3 \cos \alpha$$

$$\tan 3\alpha = \frac{3 \tan \alpha - \tan^3 \alpha}{1 - 3 \tan^2 \alpha}$$

$$\sin 4\alpha = 4 \sin \alpha \cos \alpha - 8 \sin^3 \alpha \cos \alpha$$

$$\cos 4\alpha = 8 \cos^4 \alpha - 8 \cos^2 \alpha + 1$$

$$\tan 4\alpha = \frac{4 \tan \alpha - 4 \tan^3 \alpha}{1 - 6 \tan^2 \alpha + \tan^4 \alpha}$$

$$\sin(n\alpha) = \sum_{i=0}^n \binom{n}{i} \cos^i \alpha \sin^{n-i} \alpha \sin\left(\frac{(n-i)\pi}{2}\right)$$

$$\cos(n\alpha) = \sum_{i=0}^n \binom{n}{i} \cos^i \alpha \sin^{n-i} \alpha \cos\left(\frac{(n-i)\pi}{2}\right)$$

### 3.6 Power Reduction

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2}$$

$$\tan^2 \theta = \frac{1 - \cos 2\theta}{1 + \cos 2\theta}$$

$$\sin^3 \theta = \frac{3 \sin \theta - \sin 3\theta}{4}$$

$$\cos^3 \theta = \frac{3 \cos \theta + \cos 3\theta}{4}$$

$$\sin^4 \theta = \frac{3 - 4 \cos 2\theta + \cos 4\theta}{8}$$

$$\cos^4 \theta = \frac{3 + 4 \cos 2\theta + \cos 4\theta}{8}$$

$$\sin^5 \theta = \frac{10 \sin \theta - 5 \sin 3\theta + \sin 5\theta}{16}$$

$$\cos^5 \theta = \frac{10 \cos \theta + 5 \cos 3\theta + \cos 5\theta}{16}$$

### 3.7 Product to Sum

$$\sin \alpha \cos \beta = \frac{1}{2} [\sin (\alpha + \beta) + \sin (\alpha - \beta)]$$

$$\cos \alpha \sin \beta = \frac{1}{2} [\sin (\alpha + \beta) - \sin (\alpha - \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2} [\cos (\alpha + \beta) + \cos (\alpha - \beta)]$$

$$\sin \alpha \sin \beta = \frac{1}{2} [\cos (\alpha + \beta) - \cos (\alpha - \beta)]$$

### 3.8 Sum to Product

$$\sin \theta \pm \sin \varphi = 2 \sin \frac{\theta \pm \varphi}{2} \cos \frac{\theta \mp \varphi}{2}$$

$$\cos \theta + \cos \varphi = 2 \cos \frac{\theta + \varphi}{2} \cos \frac{\theta - \varphi}{2}$$

$$\cos \theta - \cos \varphi = -2 \sin \frac{\theta + \varphi}{2} \sin \frac{\theta - \varphi}{2}$$

### 3.9 Linear Combinations

For some purposes it is important to know that any linear combination of sine waves of the same period or frequency but different phase shifts is also a sine wave with the same period or frequency, but a different phase shift.

### Definition

The two-argument form of the arctangent function, denoted by  $\tan^{-1}(y, x)$  gathers information on the signs of the inputs in order to return the appropriate quadrant of the computed angle. Thus, it is defined as:

$$\tan^{-1}(y, x) = \begin{cases} \tan^{-1}\left(\frac{y}{x}\right) & \text{if } x > 0, \\ \tan^{-1}\left(\frac{y}{x}\right) + \pi & \text{if } x < 0 \text{ and } y \geq 0, \\ \tan^{-1}\left(\frac{y}{x}\right) - \pi & \text{if } x < 0 \text{ and } y < 0, \\ +\frac{\pi}{2} & \text{if } x = 0 \text{ and } y > 0, \\ -\frac{\pi}{2} & \text{if } x = 0 \text{ and } y < 0, \\ \text{undefined} & \text{if } x = 0 \text{ and } y = 0. \end{cases}$$

### Sine and Cosine

In the case of a non-zero linear combination of a sine and cosine wave (which is just a sine wave with a phase shift of  $\frac{\pi}{2}$ ), we have:

$$a \sin x + b \cos x = c \sin(x + \theta)$$

where  $c = \pm \sqrt{a^2 + b^2}$  and  $\theta$  satisfies the equations  $c \cos \theta = a$  and  $c \sin \theta = b$ , or  $\theta = \tan^{-1}(b, a)$ .

### Arbitrary Phase Shift

More generally, for an arbitrary phase shift, we have:

$$a \sin x + b \sin(x + \theta) = c \sin(x + \varphi)$$

where  $c = \pm \sqrt{a^2 + b^2 + 2ab \cos \theta}$ , and  $\varphi$  satisfies the equations  $c \cos \varphi = a + b \cos \theta$  and  $c \sin \varphi = b \sin \theta$  or  $\varphi = \tan^{-1}(b \sin \theta, a + b \cos \theta)$ .

### 3.10 Other Related Identities

• If  $x + y + z = \pi$ , then  $\sin 2x + \sin 2y + \sin 2z = 4 \sin x \sin y \sin z$ .

• *Triple Tangent Identity.* If  $x + y + z = \pi$ , then  $\tan x + \tan y + \tan z = \tan x \tan y \tan z$ .

• *Triple Cotangent Identity.* If  $x + y + z = \frac{\pi}{2}$ , then  $\cot x + \cot y + \cot z = \cot x \cot y \cot z$ .

• *Ptolemy's Theorem.* If  $w + x + y + z = \pi$ , then  $\sin(w + x) \sin(x + y) = \sin w \sin y + \sin x \sin z$ .

•  $\cot x \cot y + \cot y \cot z + \cot z \cot x = 1$

•  $a \cos x + b \sin x = \sqrt{a^2 + b^2} \cos(x - \tan^{-1}(b, a))$

• *Tangent of an Average.*  $\tan\left(\frac{\alpha + \beta}{2}\right) = \frac{\sin \alpha + \sin \beta}{\cos \alpha + \cos \beta} = \frac{\cos \alpha - \cos \beta}{\sin \alpha - \sin \beta}$


•  $\tan x + \sec x = \tan\left(\frac{x}{2} + \frac{\pi}{4}\right)$

•  $\sum_{i=0}^n \sin(\varphi + i\alpha) = \frac{\sin \frac{(n+1)\alpha}{2} \sin\left(\varphi + \frac{n\alpha}{2}\right)}{\sin \frac{\alpha}{2}}$


- $\sum_{i=0}^n \cos(\varphi + i\alpha) = \frac{\sin \frac{(n+1)\alpha}{2} \cos\left(\varphi + \frac{n\alpha}{2}\right)}{\sin \frac{\alpha}{2}}$
  - $\sum_{n=1}^{\infty} \prod_{m=1}^n \cos \frac{m\pi}{2n+1} = 1$
- ### 3.11 Identities without Variables
- *Morrie's Law.*  $\cos 20^\circ \cdot \cos 40^\circ \cdot \cos 80^\circ = \frac{1}{8}$
  - $\sin 20^\circ \cdot \sin 40^\circ \cdot \sin 80^\circ = \frac{\sqrt{3}}{8}$
  - $\cos 24^\circ + \cos 48^\circ + \cos 96^\circ + \cos 168^\circ = \frac{1}{2}$
  - $\cos \frac{2\pi}{31} + \cos \left(2 \cdot \frac{2\pi}{31}\right) + \cos \left(4 \cdot \frac{2\pi}{31}\right) + \cos \left(5 \cdot \frac{2\pi}{31}\right) + \cos \left(8 \cdot \frac{2\pi}{31}\right) + \cos \left(10 \cdot \frac{2\pi}{31}\right) = \frac{1}{2}$
  - $\cos \frac{\pi}{5} = \cos 36^\circ = \frac{1}{4}(\sqrt{5} + 1) = \frac{1}{2}\varphi$
  - $\sin \frac{\pi}{10} = \sin 18^\circ = \frac{1}{4}(\sqrt{5} - 1) = \frac{1}{2}\varphi^{-1}$
  - $\sin^2 18^\circ + \sin^2 30^\circ = \sin^2 36^\circ$

## 4 Graphs


### 4.1 $y = \sin x$




### 4.4 $y = \csc x$




### 4.2 $y = \cos x$




### 4.5 $y = \sec x$



### 4.3 $y = \tan x$



### 4.6 $y = \cot x$



5 Tables

5.1 Exact Values of Trigonometric Functions

$A^\circ$	$A$ rad	$\sin A$	$\cos A$	$\tan A$	$\cot A$	$\sec A$	$\csc A$
$0^\circ$	0	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	1	0	$\infty$	1	$\infty$
$15^\circ$	$\pi/12$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$2-\sqrt{3}$	$2+\sqrt{3}$	$\sqrt{6}-\sqrt{2}$	$\sqrt{6}+\sqrt{2}$
$30^\circ$	$\pi/6$	$\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	2
$45^\circ$	$\pi/4$	$\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
$60^\circ$	$\pi/3$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	2	$\frac{2}{3}\sqrt{3}$
$75^\circ$	$5\pi/12$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$2+\sqrt{3}$	$2-\sqrt{3}$	$\sqrt{6}+\sqrt{2}$	$\sqrt{6}-\sqrt{2}$
$90^\circ$	$\pi/2$	1	0	$\pm\infty$	0	$\pm\infty$	1
$105^\circ$	$7\pi/12$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-(2+\sqrt{3})$	$-(2-\sqrt{3})$	$-(\sqrt{6}+\sqrt{2})$	$\sqrt{6}-\sqrt{2}$
$120^\circ$	$2\pi/3$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	-2	$\frac{2}{3}\sqrt{3}$
$135^\circ$	$3\pi/4$	$\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	-1	-1	$-\sqrt{2}$	$\sqrt{2}$
$150^\circ$	$5\pi/6$	$\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	2
$165^\circ$	$11\pi/12$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-(2-\sqrt{3})$	$-(2+\sqrt{3})$	$-(\sqrt{6}-\sqrt{2})$	$\sqrt{6}+\sqrt{2}$
$180^\circ$	$\pi$	0	-1	0	$\mp\infty$	-1	$\pm\infty$
$195^\circ$	$13\pi/12$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$2-\sqrt{3}$	$2+\sqrt{3}$	$-(\sqrt{6}-\sqrt{2})$	$-(\sqrt{6}+\sqrt{2})$
$210^\circ$	$7\pi/6$	$\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	-2
$225^\circ$	$5\pi/4$	$-\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	1	1	$-\sqrt{2}$	$-\sqrt{2}$
$240^\circ$	$4\pi/3$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	-2	$-\frac{2}{3}\sqrt{3}$
$255^\circ$	$17\pi/12$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$2+\sqrt{3}$	$2-\sqrt{3}$	$-(\sqrt{6}+\sqrt{2})$	$-(\sqrt{6}-\sqrt{2})$
$270^\circ$	$3\pi/2$	-1	0	$\pm\infty$	0	$\mp\infty$	-1
$285^\circ$	$19\pi/12$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-(2+\sqrt{3})$	$-(2-\sqrt{3})$	$\sqrt{6}+\sqrt{2}$	$-(\sqrt{6}-\sqrt{2})$
$300^\circ$	$5\pi/3$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	2	$-\frac{2}{3}\sqrt{3}$
$315^\circ$	$7\pi/4$	$-\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	-1	-1	$\sqrt{2}$	$-\sqrt{2}$
$330^\circ$	$11\pi/6$	$-\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	-2
$345^\circ$	$23\pi/12$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-(2+\sqrt{3})$	$-(2+\sqrt{3})$	$\sqrt{6}-\sqrt{2}$	$-(\sqrt{6}+\sqrt{2})$
$360^\circ$	$2\pi$	0	1	0	$\mp\infty$	1	$\mp\infty$

5.2 Relations Between Trig Functions

	$\sin \theta = u$	$\cos \theta = u$	$\tan \theta = u$	$\csc \theta = u$	$\sec \theta = u$	$\cot \theta = u$
$\sin \theta$	$u$	$\sqrt{1-u^2}$	$\frac{u}{\sqrt{1+u^2}}$	$\frac{1}{u}$	$\frac{\sqrt{u^2-1}}{u}$	$\frac{1}{\sqrt{1+u^2}}$
$\cos \theta$	$\sqrt{1-u^2}$	$u$	$\frac{1}{\sqrt{1+u^2}}$	$\frac{1}{\sqrt{u^2-1}}$	$\frac{1}{u}$	$\frac{u}{\sqrt{1+u^2}}$
$\tan \theta$	$\frac{u}{\sqrt{1-u^2}}$	$\frac{\sqrt{1-u^2}}{u}$	$u$	$\frac{1}{\sqrt{u^2-1}}$	$\frac{1}{\sqrt{u^2-1}}$	$\frac{1}{u}$
$\csc \theta$	$\frac{1}{u}$	$\frac{1}{\sqrt{1-u^2}}$	$\frac{\sqrt{1+u^2}}{u}$	$u$	$\frac{u}{\sqrt{u^2-1}}$	$\frac{\sqrt{1+u^2}}{u}$
$\sec \theta$	$\frac{1}{\sqrt{1-u^2}}$	$\frac{1}{u}$	$\frac{\sqrt{1+u^2}}{u}$	$\frac{u}{\sqrt{u^2-1}}$	$u$	$\frac{\sqrt{1+u^2}}{u}$
$\cot \theta$	$\frac{\sqrt{1-u^2}}{u}$	$\frac{u}{\sqrt{1-u^2}}$	$\frac{1}{u}$	$\frac{1}{\sqrt{u^2-1}}$	$\frac{1}{\sqrt{u^2-1}}$	$u$

6 Inverse Trigonometric Functions

Reciprocal Identities

If  $x = \sin y$ , then  $y = \sin^{-1} x$ , i.e. the angle whose sine is  $x$  or arcsine of  $x$ , is a multiple-valued function of  $x$  which is a collection of single-valued functions called *branches*. Similarly, the other inverse trigonometric functions are multiple valued.

For many purposes, a particular branch is required. This is called the *principal branch* and the values for this branch are called *principal values*.

6.1 Principal Values

Since none of the six trigonometric functions are one-to-one, they are restricted in order to have inverse functions. Therefore the ranges of the inverse functions are proper subsets of the domains of the original functions.

Negative Identities

$$\begin{aligned} \sin^{-1}(-x) &= -\sin^{-1} x \\ \cos^{-1}(-x) &= \pi - \cos^{-1} x \\ \tan^{-1}(-x) &= -\tan^{-1} x \\ \csc^{-1}(-x) &= -\csc^{-1} x \\ \sec^{-1}(-x) &= \pi - \sec^{-1} x \\ \cot^{-1}(-x) &= \pi - \cot^{-1} x \end{aligned}$$

Complementary Identities

$$\begin{aligned} \sin^{-1} x + \cos^{-1} x &= \frac{\pi}{2} \\ \tan^{-1} x + \cot^{-1} x &= \frac{\pi}{2} \end{aligned}$$

6.2 Identities

In all cases it is assumed that principal values are used.

$\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}$

Sum and Difference Identities


$$\begin{aligned}\sin^{-1} \alpha \pm \sin^{-1} \beta &= \sin^{-1} \left( \alpha \sqrt{1-\beta^2} \pm \beta \sqrt{1-\alpha^2} \right) \\ \cos^{-1} \alpha \pm \cos^{-1} \beta &= \cos^{-1} \left( \alpha \beta \pm \sqrt{(1-\alpha^2)(1-\beta^2)} \right) \\ \tan^{-1} \alpha \pm \tan^{-1} \beta &= \tan^{-1} \left( \frac{\alpha \pm \beta}{1 \mp \alpha \beta} \right)\end{aligned}$$

7 Relationships Between Sides and Angles

7.1 Law of Sines

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$$

Extended Law of Sines



$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma} = 2R,$$

where  $R$  is the circumradius of the triangle.

7.2 Law of Cosines

$$\begin{aligned}\cos \alpha &= \frac{b^2 + c^2 - a^2}{2bc}; & a &= \sqrt{b^2 + c^2 - 2bc \cos \alpha} \\ \cos \beta &= \frac{a^2 + c^2 - b^2}{2ac}; & b &= \sqrt{a^2 + c^2 - 2ac \cos \beta} \\ \cos \gamma &= \frac{a^2 + b^2 - c^2}{2ab}; & c &= \sqrt{a^2 + b^2 - 2ab \cos \gamma}\end{aligned}$$

7.3 Law of Tangents

$$\begin{aligned}\frac{a-b}{a+b} &= \frac{\tan \frac{1}{2}(\alpha-\beta)}{\tan \frac{1}{2}(\alpha+\beta)} \\ \frac{b-c}{b+c} &= \frac{\tan \frac{1}{2}(\beta-\gamma)}{\tan \frac{1}{2}(\beta+\gamma)} \\ \frac{c-a}{c+a} &= \frac{\tan \frac{1}{2}(\gamma-\alpha)}{\tan \frac{1}{2}(\gamma+\alpha)}\end{aligned}$$

7.4 Law of Cotangents

Let  $s$  be the semiperimeter, that is,  $s = \frac{a+b+c}{2}$ , and  $r$  be the radius of the inscribed circle, then:

$$\frac{\cot \frac{\alpha}{2}}{s-a} = \frac{\cot \frac{\beta}{2}}{s-b} = \frac{\cot \frac{\gamma}{2}}{s-c} = \frac{1}{r}$$

and furthermore that the inradius is given by:


$$r = \sqrt{\frac{(s-a)(s-b)(s-c)}{s}}$$

7.5 Mollweide's Formula

Each of these identities uses all six parts of the triangle—the three angles and the lengths of the three sides.

$$\frac{a+b}{c} = \frac{\cos \frac{\alpha-\beta}{2}}{\sin \frac{\gamma}{2}}; \quad \frac{a-b}{c} = \frac{\sin \frac{\alpha-\beta}{2}}{\cos \frac{\gamma}{2}}$$

7.6 Stewart's Theorem



Let  $D$  be a point in  $\overline{BC}$  of  $\triangle ABC$ . If  $|BD| = m$ ,  $|CD| = n$ , and  $|AD| = d$ , then  $b^2m + c^2n = a(d^2 + mn)$ .

7.7 Angles in Terms of Sides

Let  $s = \frac{a+b+c}{2}$  be the semiperimeter of the triangle, then:

$$\begin{aligned}\alpha &= \sin^{-1} \left( \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)} \right) \\ \beta &= \sin^{-1} \left( \frac{2}{ac} \sqrt{s(s-a)(s-b)(s-c)} \right) \\ \gamma &= \sin^{-1} \left( \frac{2}{ab} \sqrt{s(s-a)(s-b)(s-c)} \right)\end{aligned}$$

8 Solving Triangles

A general form triangle has six main characteristics: three linear (side lengths  $a, b, c$ ) and three angular ( $\alpha, \beta, \gamma$ ). The classical plane trigonometry problem is to specify three of the six characteristics and determine the other three. A triangle can be uniquely determined in this sense when given any of the following:

- Three sides (SSS)
- Two sides and the included angle (SAS)
- Two sides and an angle not included between them (SSA), if the side length adjacent to the angle is shorter than the other side length.
- A side and the two angles adjacent to it (ASA)
- A side, the angle opposite to it and an angle adjacent to it (AAS).


For all cases in the plane, at least one of the side lengths must be specified. If only the angles are given, the side lengths cannot be determined, because any similar triangle is a solution.

Notes

- To find an unknown angle, the law of cosines is safer than the law of sines. The reason is that the value of sine for the angle does not uniquely determine this angle. For example, if  $\sin \beta = 0.5$ , the angle  $\beta$  can be equal either  $30^\circ$  or  $150^\circ$ . Using the law of cosines avoids this problem: within the interval from  $0^\circ$  to  $180^\circ$  the cosine value unambiguously determines its angle. On the other hand, if the angle is small (or close to  $180^\circ$ ), then it is more robust numerically to determine it from its sine than its cosine because the arc-cosine function has a divergent derivative at 1 (or  $-1$ ).

- We assume that the relative position of specified characteristics is known. If not, the mirror reflection of the triangle will also be a solution. For example, three side lengths uniquely define either a triangle or its reflection.

8.1 AAS/ASA Triangle



Let three side lengths  $a, b, c$  be specified. To find the angles  $\alpha, \beta$ , the law of cosines can be used:

$$\alpha = \cos^{-1} \frac{b^2 + c^2 - a^2}{2bc}; \quad \beta = \cos^{-1} \frac{a^2 + c^2 - b^2}{2ac}.$$

Then angle  $\gamma = 180^\circ - \alpha - \beta$ .


The known characteristics are the side  $c$  and the angles  $\alpha, \beta$ . The third angle  $\gamma = 180^\circ - \alpha - \beta$ .

Two unknown side can be calculated from the law of sines:

$$a = \frac{c \sin \alpha}{\sin \gamma}; \quad b = \frac{c \sin \beta}{\sin \gamma}.$$

The procedure for solving an AAS triangle is same as that for an ASA triangle: First, find the third angle by using the angle sum property of a triangle, then find the other two sides using the law of sines.

8.2 SAS Triangle



Here the lengths of sides  $a, b$  and the angle  $\gamma$  between these sides are known. The third side can be determined from the law of cosines:


$$c = \sqrt{a^2 + b^2 - 2ab \cos \gamma}$$

Now we use law of cosines to find the second angle:

$$\alpha = \cos^{-1} \frac{b^2 + c^2 - a^2}{2bc}$$

Finally,  $\beta = 180^\circ - \alpha - \gamma$ .

8.3 SSS Triangle




Let three side lengths  $a, b, c$  be specified. To find the angles  $\alpha, \beta$ , the law of cosines can be used:

$$\alpha = \cos^{-1} \frac{b^2 + c^2 - a^2}{2bc}; \quad \beta = \cos^{-1} \frac{a^2 + c^2 - b^2}{2ac}.$$

Then angle  $\gamma = 180^\circ - \alpha - \beta$ .

### 8.4 SSA Triangle



This case is not solvable in all cases; a solution is guaranteed to be unique only if the side length adjacent to the angle is shorter than the other side length. Assume that two sides  $b$ ,  $c$  and the angle  $\beta$  are known. The equation for the angle  $\gamma$  can be implied from the law of sines:

$$\sin \gamma = \frac{c \sin \beta}{b} \quad (\text{equation's right side}).$$

We denote further  $D = \frac{c \sin \beta}{b}$  (equation's right side). There are four possible cases:

1. If  $D > 1$ , no such triangle exists because the side  $b$  does not reach line  $BC$ . For the same reason a solution does not exist if the angle  $\beta \geq 90^\circ$  and  $b \leq c$ .
2. If  $D = 1$ , a unique solution exists:  $\gamma = 90^\circ$ , i.e., the triangle is right-angled.
3. If  $D < 1$ , two alternatives are possible.



- (a) If  $b < c$ , the angle  $\gamma$  may be acute:  $\gamma = \sin^{-1} D$  or obtuse:  $\gamma' = 180^\circ - \gamma$ . The picture above shows the point  $C$ , the side  $b$  and the angle  $\gamma$  as the first solution, and the point  $C'$ , side  $b'$  and the angle  $\gamma'$  as the second solution.
- (b) If  $b \geq c$  then  $\beta \geq \gamma$  (the larger side corresponds to a larger angle). Since no triangle can have two obtuse angles,  $\gamma$  is acute angle and the solution  $\gamma = \sin^{-1} D$  is unique.

Once  $\gamma$  is obtained, the third angle  $\alpha = 180^\circ - \beta - \gamma$ . The third side can then be found from the law of sines:

$$a = \frac{b \sin \alpha}{\sin \beta}$$

### 8.5 Right Triangle

Solving right triangles is simply using the definitions of the trigonometric functions and the Pythagorean theorem to determine the other parts. The right angle  $\gamma = 90^\circ$  is always assumed to be given.

## 9 Polar Coordinates

A point  $P$  can be located by rectangular coordinates  $(x, y)$  or polar coordinates  $(r, \theta)$ .

The angle  $\theta$  is a *directed angle*, that is, it is positive if it is measured counterclockwise from the initial side to the terminal side, and negative if it is measured clockwise.

The value  $r$  is a *directed distance*, it is positive if the point  $P$  lies on the terminal side of  $\theta$  and negative if  $P$  is on the extension of the terminal side.

### 9.1 Properties

- Every ordered pair of polar coordinates  $(r, \theta)$  locates a unique point in the plane.
- However, a point  $P$  on the plane may be specified by an infinite number of ordered pairs  $(r, \theta)$ .
- The pole  $O$  may be specified by the ordered pair  $(0, \theta)$  where  $\theta \in \mathbb{R}$ .
- Let  $P(r, \theta)$  be a point in the polar plane. Then  $(r, \theta + 2k\pi)$  are also coordinates of the point  $P$  for any  $k \in \mathbb{Z}$ .
- It can also be shown that  $((-1)^n r, \theta + n\pi)$  are also coordinates of  $P$ , where  $n \in \mathbb{Z}$ .

## 9.2 Coordinate Transformation

### Polar to Rectangular

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$$

### Rectangular to Polar

$$\begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1} \left( \frac{y}{x} \right) \approx \tan^{-1} \left( \frac{y}{x} \right) \end{cases}$$

where  $\tan^{-1} (y, x)$  is the two-argument form of the arctangent function (see section 3.9).

## 10 Special Polar Graphs

### Theorem

A polar graph is:

1. **symmetric with respect to the polar axis** if an equivalent equation is obtained when  $(r, \theta)$  is replaced by either  $(r, -\theta)$  or  $(-r, \pi - \theta)$ .
2. **symmetric with respect to the  $\frac{\pi}{2}$ -axis** if an equivalent equation is obtained when  $(r, \theta)$  is replaced by either  $(r, \pi - \theta)$  or  $(-r, -\theta)$ .

- 3. **symmetric with respect to the pole** if an equivalent equation is obtained when  $(r, \theta)$  is replaced by either  $(-r, \theta)$  or  $(r, \pi + \theta)$ .


### 10.1 Limaçon of Pascal

A polar equation of the form  $r = a + b \cos \theta$  or  $r = a + b \sin \theta$  has a polar graph which is called a *limaçon*.

Let  $OQ$  be a line joining origin  $O$  to any point in  $Q$  on a circle of diameter  $b$  passing through  $O$ . Then the curve is the locus of all points in  $P$  such that  $|PQ| = a$

### Types of Limaçons

- Looped limaçon if  $0 < \left| \frac{a}{b} \right| < 1$
- Dimpled limaçon if  $1 < \left| \frac{a}{b} \right| < 2$
- Convex limaçon if  $\left| \frac{a}{b} \right| \geq 2$



### Symmetry and Direction


For  $a > 0$  and  $b > 0$ :

	symmetry	points to the
$r = a + b \cos \theta$	polar axis	right
$r = a - b \cos \theta$	polar axis	left
$r = a + b \sin \theta$	$\frac{\pi}{2}$ -axis	top
$r = a - b \sin \theta$	$\frac{\pi}{2}$ -axis	bottom




### 10.2 Rose

A rose with  $n$  leaves has a polar equation  $r = a \cos(n\theta)$  or  $r = a \sin(n\theta)$  where  $a$  is a constant and  $n$  is an odd integer.



For an even integer  $n$ , the polar graph of an equation  $r = a \cos(n\theta)$  or  $r = a \sin(n\theta)$  is a rose with  $2n$  leaves.




#### Properties

- The length of one leaf in the polar graph of a rose is  $|a|$ .
- If  $n$  is odd, then the graph of the polar equation  $r = a \cos(n\theta)$  is symmetric with respect to the polar axis.
- If  $n$  is odd, then the graph of the polar equation  $r = a \sin(n\theta)$  is symmetric with respect to the  $\frac{\pi}{2}$ -axis.

- A rose with an even number of leaves is symmetric with respect to the polar axis, the  $\frac{\pi}{2}$ -axis, and the pole.


### 10.3 Spiral of Archimedes

The polar graph of a polar equation  $r = a\theta$  where  $\theta > 0$  and  $a \in \mathbb{R}$  is called a *spiral*.




### 10.4 Lemniscate of Bernoulli

A polar equation  $r^2 = a \cos 2\theta$  or  $r^2 = a \sin 2\theta$  has a polar graph that is called a *lemniscate*.




### 10.5 Folium of Descartes

A *folium* is a plane curve proposed by Descartes to challenge Fermat's extremum-finding techniques. It has a polar equation  $r = \frac{3a \sec \theta \tan \theta}{1 + \tan^3 \theta}$ .




### 10.8 Epispiral

The *epispiral* is a plane curve with a polar equation  $r = a \sec(n\theta)$ . Then there are  $n$  sections if  $n$  is odd (in blue), or  $2n$  sections if  $n$  is even (in red). A slightly more symmetric version considers instead  $r = a |\sec(n\theta)|$ .



### 10.6 Spiral of Fermat

The *Fermat's spiral*, also known as the parabolic spiral, has a polar equation  $r^2 = a^2 \theta$ . The resulting spiral is symmetric with respect to the origin.




### 10.7 Cissoid of Diocles

Given an origin  $O$  and a point  $P$  on the curve, let  $S$  be the point where the extension of the line  $OP$  intersects the line  $x = 2a$  and  $R$  be the intersection of the circle of radius  $a$


### 10.9 Lituus

The *lituus* is the locus of the point  $P$  moving such that the area of a circular sector remains constant. It means a "crook," in the sense of a bishop's crozier. It has a polar equation  $r^2\theta = a^2$ .




### 10.11 Butterfly Curve

The *butterfly curve* is a transcendental plane curve discovered by Temple H. Fay. The curve is given by the polar equation  $r = e^{\sin \theta} - 2 \cos 4\theta + \sin^5 \left( \frac{1}{24} (2\theta - \pi) \right)$ .




### 10.10 Eight Curve

The eight curve, also known as the *lemniscate of Gerono*, is given by the polar equation  $r^2 = a^2 \sec^4 \theta \cos(2\theta)$ . It has vertical tangents  $(\pm a, 0)$  and horizontal tangents at  $(\pm \frac{\sqrt{2}}{2}a, \pm \frac{1}{2}a)$ .




### 10.12 Strophoid

Let  $C$  be a curve, let  $O$  be a fixed point (the pole), and let  $O'$  be a second fixed point. Let  $P$  and  $P'$  be points on a line through  $O$  meeting  $C$  at  $Q$  such that  $P'Q = QP = QO'$ . The locus of  $P$  and  $P'$  is called the *strophoid* of  $C$  with respect to the pole  $O$  and fixed point  $O'$ . Its polar equation is  $r = \frac{b \sin(a - 2\theta)}{\sin(a - \theta)}$ .




### 10.13 Cochleoid

A *cochleoid* is a snail-shaped curve similar to a strophoid which can be represented by the polar equation  $r = \frac{a \sin \theta}{\theta}$ .




### 10.15 Freeth's Nephroid

It is a strophoid of a circle with the pole  $O$  at the center of the circle and the fixed point  $P$  on the circumference of the circle. It has a polar equation  $r = a(1 + 2 \sin \frac{\theta}{2})$ .



### 10.14 Cycloid of Ceva

The *cycloid of Ceva* is a polar curve that can be used for angle trisection. It has the polar equation  $r = 1 + 2 \cos 2\theta$ .





## 11 Miscellaneous Stuff

### 11.1 Pythagorean Triples

A Pythagorean triple  $(a, b, c)$  is a triple of positive integers such that  $a^2 + b^2 = c^2$ . It is primitive if the greatest common divisor of  $a$ ,  $b$ , and  $c$  is 1. For any primitive Pythagorean triple, either  $m$  or  $n$  is even, but not both (i.e.  $m \not\equiv n \pmod{2}$ ). When that is done, then every primitive Pythagorean triple  $(a, b, c)$  is of the form  $(a, b, c) = (m^2 - n^2, 2mn, m^2 + n^2)$  where  $m$  and  $n$  are relatively prime (i.e.  $\gcd(m, n) = 1$ ) and  $1 \leq n < m$ .

$m$	$n$	$a$	$b$	$c$	$m$	$n$	$a$	$b$	$c$
2	1	3	4	5	11	8	57	176	185
3	2	5	12	13	11	10	21	220	221
4	1	15	8	17	12	1	143	24	145
4	3	7	24	25	12	3	135	72	153
5	2	21	20	29	12	5	119	120	169
5	4	9	40	41	12	7	95	168	193
6	1	35	12	37	12	9	63	216	225
6	3	27	36	45	12	11	23	264	265
6	5	11	60	61	13	2	165	52	173
7	2	45	28	53	13	4	153	104	185
7	4	33	56	65	13	6	133	156	205
7	6	13	84	85	13	8	105	208	233
8	1	63	16	65	13	10	69	260	269
8	3	55	48	73	13	12	25	312	313
8	5	39	80	89	14	1	195	28	197
8	7	15	112	113	14	3	187	84	205
9	2	77	36	85	14	5	171	140	221
9	4	65	72	97	14	7	147	196	245
9	6	45	108	117	14	9	115	252	277
9	8	17	144	145	14	11	75	308	317
10	1	99	20	101	14	13	27	364	365
10	3	91	60	109	15	2	221	60	229
10	5	75	100	125	15	4	209	120	241
10	7	51	140	149	15	6	189	180	261
10	9	19	180	181	15	8	161	240	289
11	2	117	44	125	15	10	125	300	325
11	4	105	88	137	15	12	81	360	369
11	6	85	132	157	15	14	29	420	421

### 11.2 Triangle Centers

#### Barycenter/Centroid

The geometric centroid (center of mass) of the polygon vertices of a triangle is the point  $G$  which is also the intersection of the triangle's three triangle medians. The point is therefore sometimes called the median point. The centroid is always in the interior of the triangle.

For a triangle with Cartesian vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$ ,  $(x_3, y_3)$ , the Cartesian coordinates of the centroid are given by:

$$G(x_0, y_0) = \left( \frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right)$$

#### Incenter

The incenter  $I$  is the center of the incircle for a triangle. The corresponding radius of the incircle is known as the inradius.

The incenter can be constructed as the intersection of angle bisectors. It is also the interior point for which distances to the sides of the triangle are equal.

For a triangle with Cartesian vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$ ,  $(x_3, y_3)$ , the Cartesian coordinates of the incenter are given by:

$$I(x_0, y_0) = \left( \frac{ax_1 + bx_2 + cx_3}{a + b + c}, \frac{ay_1 + by_2 + cy_3}{a + b + c} \right)$$

#### Orthocenter

The intersection  $H$  of the three altitudes  $AH_A$ ,  $BH_B$ , and  $CH_C$  of a triangle is called the orthocenter.

For a triangle with Cartesian vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$ ,  $(x_3, y_3)$ , the Cartesian coordinates of the circumcenter are given by:

$$H(x_0, y_0) = \left( \frac{x_1 t_A + x_2 t_B + x_3 t_C}{t_A + t_B + t_C}, \frac{y_1 t_A + y_2 t_B + y_3 t_C}{t_A + t_B + t_C} \right)$$

where  $t_A, t_B, t_C$  is equal to  $\tan A, \tan B, \tan C$  respectively.

#### Circumcenter

The circumcenter is the center  $O$  of a triangle's circumscribed circle. It can be found as the intersection of the perpendicular bisectors.

For a triangle with Cartesian vertices  $(x_1, y_1)$ ,  $(x_2, y_2)$ ,  $(x_3, y_3)$ , the Cartesian coordinates of the circumcenter are given by:

$$O(x_0, y_0) = (O_x, O_y)$$

$$\text{where } O_x = \frac{(y_2 - y_3)(x_1^2 + (y_1 - y_2)(y_1 - y_3)) + x_2^2(y_1 - y_2) + x_3^2(y_1 - y_2)}{2(x_1(y_2 - y_3) + x_2(y_1 - y_3) + x_3(y_1 - y_2))}$$

$$\text{and } O_y = \frac{2(x_2 - x_3)(-x_1^2 + x_2^2 - y_2^2 + y_3^2) + 2(x_1 - x_2)(x_2^2 - x_3^2 + y_2^2 - y_3^2)}{4(x_1(y_2 - y_3) + x_2(y_1 - y_3) + x_3(y_1 - y_2))}$$

#### Excenter

An excenter, denoted  $J_i$ , is the center of an excircle of a triangle. An excircle is a circle tangent to the extensions of two sides and the third side. It is also known as an escribed circle.

## Trigonometry Gram Sheet

### 11.3 Area of the Triangle

#### Base and Altitude

$$A_{\Delta} = \frac{1}{2}bh$$

where  $b$  is the base and  $h$  is the altitude.

#### SAS Triangle

$$A_{\Delta} = \frac{1}{2}ab \sin \gamma = \frac{1}{2}bc \sin \alpha = \frac{1}{2}ac \sin \beta$$

#### SSS Triangle

Given three sides  $a$ ,  $b$ , and  $c$ , the area of the triangle can be determined by Heron's formula:

$$A_{\Delta} = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \frac{1}{4} \sqrt{4a^2b^2 - (a^2 + b^2 - c^2)^2}$$

where  $s = \frac{a+b+c}{2}$  is the semiperimeter. The area can also be determined by:

$$A_{\Delta} = \frac{abc}{4R}$$

where  $R$  is the circumradius.

#### AAS Triangle

$$A_{\Delta} = \frac{a^2 \sin \beta \sin \gamma}{2 \sin \alpha} = \frac{b^2 \sin \alpha \sin \gamma}{2 \sin \beta} = \frac{c^2 \sin \alpha \sin \beta}{2 \sin \gamma}$$

where the missing angle can be easily determined through  $\alpha + \beta + \gamma = 180^\circ$ .

#### ASA Triangle

$$A_{\Delta} = \frac{a^2}{2(\cot \beta + \cot \gamma)} = \frac{b^2}{2(\cot \alpha + \cot \gamma)} = \frac{c^2}{2(\cot \alpha + \cot \beta)}$$

#### Three Vertices

Given the vertices of the triangle  $(x_i, y_i) \forall i \in \{1, 2, 3\}$ , the shoelace formula can be applied for  $n = 3$  as follows:

$$A_{\Delta} = \frac{1}{2} \left| \sum_{i=1}^2 x_i y_{i+1} + x_2 y_1 - \sum_{i=1}^2 x_{i+1} y_i - x_1 y_3 \right|$$

$$= \frac{1}{2} |x_1 y_2 + x_2 y_3 + x_3 y_1 - x_2 y_1 - x_3 y_2 - x_1 y_3|$$

However, if one of the vertices is at the origin, then the area of the triangle can be simplified:

$$A_{\Delta} = \frac{1}{2} \left\| \begin{pmatrix} x_2 & x_3 \\ y_2 & y_3 \end{pmatrix} \right\| = \frac{1}{2} |x_2 y_3 - x_3 y_2|$$

where  $\|\cdot\|$  denotes the absolute value of the determinant.