

Assignment 4: Data Wrangling

Allison Barbaro

Spring 2023

OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Wrangling

Directions

1. Rename this file `<FirstLast>_A04_DataWrangling.Rmd` (replacing `<FirstLast>` with your first and last name).
2. Change “Student Name” on line 3 (above) with your name.
3. Work through the steps, **creating code and output** that fulfill each instruction.
4. Be sure to **answer the questions** in this assignment document.
5. When you have completed the assignment, **Knit** the text and code into a single PDF file.

The completed exercise is due on Friday, Feb 20th @ 5:00pm.

Set up your session

- 1a. Load the `tidyverse`, `lubridate`, and `here` packages into your session.
 - 1b. Check your working directory.
 - 1c. Read in all four raw data files associated with the EPA Air dataset, being sure to set string columns to be read in as factors. See the README file for the EPA air datasets for more information (especially if you have not worked with air quality data previously).
2. Apply the `glimpse()` function to reveal the dimensions, column names, and structure of each dataset.

```
# 1a - loaded all necessary packages
library(tidyverse)
```

```
## -- Attaching packages ----- tidyverse 1.3.2 --
## v ggplot2 3.4.0      v purrr   1.0.0
## v tibble  3.1.8      v dplyr  1.1.0
## v tidyr   1.2.1      v stringr 1.5.0
## v readr   2.1.3      v forcats 0.5.2
## -- Conflicts ----- tidyverse_conflicts() --
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()    masks stats::lag()
```

```
library(lubridate)
```

```
## Loading required package: timechange
##
## Attaching package: 'lubridate'
##
## The following objects are masked from 'package:base':
##
##     date, intersect, setdiff, union
```

```
library(here)
```

```
## here() starts at /home/guest/R/EDA-Spring2023
```

```
# 1b - checked workspace
getwd()
```

```
## [1] "/home/guest/R/EDA-Spring2023"
```

```
# 1c - read in all raw data needed
ozone.2018 <- read.csv(file = here("Data/Raw/EPAair_03_NC2018_raw.csv"), stringsAsFactors = TRUE)

ozone.2019 <- read.csv(file = here("Data/Raw/EPAair_03_NC2019_raw.csv"), stringsAsFactors = TRUE)

PM25.2018 <- read.csv(file = here("Data/Raw/EPAair_PM25_NC2018_raw.csv"), stringsAsFactors = TRUE)

PM25.2019 <- read.csv(file = here("Data/Raw/EPAair_PM25_NC2019_raw.csv"), stringsAsFactors = TRUE)

# 2 - gathered initial info about each dataset
glimpse(ozone.2018)
```

```
## Rows: 9,737
## Columns: 20
## $ Date                <fct> 03/01/2018, 03/02/2018, 03/03/201~
## $ Source              <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS~
## $ Site.ID             <int> 370030005, 370030005, 370030005, ~
## $ POC                 <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Max.8.hour.Ozone.Concentration <dbl> 0.043, 0.046, 0.047, 0.049, 0.047~
## $ UNITS               <fct> ppm, ppm, ppm, ppm, ppm, ppm, ppm~
## $ DAILY_AQI_VALUE     <int> 40, 43, 44, 45, 44, 28, 33, 41, 4~
## $ Site.Name           <fct> Taylorsville Liledoun, Taylorsvil~
## $ DAILY_OBS_COUNT     <int> 17, 17, 17, 17, 17, 17, 17, 17, 1~
## $ PERCENT_COMPLETE    <dbl> 100, 100, 100, 100, 100, 100, 100~
## $ AQS_PARAMETER_CODE  <int> 44201, 44201, 44201, 44201, 44201~
## $ AQS_PARAMETER_DESC  <fct> Ozone, Ozone, Ozone, Ozone, Ozone~
## $ CBSA_CODE           <int> 25860, 25860, 25860, 25860, 25860~
## $ CBSA_NAME           <fct> "Hickory-Lenoir-Morganton, NC", "~
## $ STATE_CODE          <int> 37, 37, 37, 37, 37, 37, 37, 37, 3~
## $ STATE              <fct> North Carolina, North Carolina, N~
## $ COUNTY_CODE         <int> 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, ~
## $ COUNTY             <fct> Alexander, Alexander, Alexander, ~
## $ SITE_LATITUDE       <dbl> 35.9138, 35.9138, 35.9138, 35.913~
## $ SITE_LONGITUDE      <dbl> -81.191, -81.191, -81.191, -81.19~
```

```
glimpse(ozone.2019)
```

```
## Rows: 10,592
## Columns: 20
## $ Date                <fct> 01/01/2019, 01/02/2019, 01/03/201~
## $ Source              <fct> AirNow, AirNow, AirNow, AirNow, A~
## $ Site.ID             <int> 370030005, 370030005, 370030005, ~
## $ POC                 <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Max.8.hour.Ozone.Concentration <dbl> 0.029, 0.018, 0.016, 0.022, 0.037~
## $ UNITS               <fct> ppm, ppm, ppm, ppm, ppm, ppm, ppm~
## $ DAILY_AQI_VALUE     <int> 27, 17, 15, 20, 34, 34, 27, 35, 3~
## $ Site.Name           <fct> Taylorsville Liledoun, Taylorsvil~
## $ DAILY_OBS_COUNT     <int> 24, 24, 24, 24, 24, 24, 24, 24, 2~
## $ PERCENT_COMPLETE    <dbl> 100, 100, 100, 100, 100, 100, 100~
## $ AQS_PARAMETER_CODE  <int> 44201, 44201, 44201, 44201, 44201~
## $ AQS_PARAMETER_DESC  <fct> Ozone, Ozone, Ozone, Ozone, Ozone~
## $ CBSA_CODE           <int> 25860, 25860, 25860, 25860, 25860~
## $ CBSA_NAME           <fct> "Hickory-Lenoir-Morganton, NC", "~
## $ STATE_CODE          <int> 37, 37, 37, 37, 37, 37, 37, 37, 3~
## $ STATE               <fct> North Carolina, North Carolina, N~
## $ COUNTY_CODE         <int> 3, 3, 3, 3, 3, 3, 3, 3, 3, 3, ~
## $ COUNTY              <fct> Alexander, Alexander, Alexander, ~
## $ SITE_LATITUDE       <dbl> 35.9138, 35.9138, 35.9138, 35.913~
## $ SITE_LONGITUDE      <dbl> -81.191, -81.191, -81.191, -81.19~
```

```
glimpse(PM25.2018)
```

```
## Rows: 8,983
## Columns: 20
## $ Date                <fct> 01/02/2018, 01/05/2018, 01/08/2018, 01/~
## $ Source              <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS, AQS,~
## $ Site.ID             <int> 370110002, 370110002, 370110002, 370110~
## $ POC                 <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Mean.PM2.5.Concentration <dbl> 2.9, 3.7, 5.3, 0.8, 2.5, 4.5, 1.8, 2.5,~
## $ UNITS               <fct> ug/m3 LC, ug/m3 LC, ug/m3 LC, ug/m3 LC,~
## $ DAILY_AQI_VALUE     <int> 12, 15, 22, 3, 10, 19, 8, 10, 18, 7, 24~
## $ Site.Name           <fct> Linville Falls, Linville Falls, Linvill~
## $ DAILY_OBS_COUNT     <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ PERCENT_COMPLETE    <dbl> 100, 100, 100, 100, 100, 100, 100, 100,~
## $ AQS_PARAMETER_CODE  <int> 88502, 88502, 88502, 88502, 88502, 8850~
## $ AQS_PARAMETER_DESC  <fct> Acceptable PM2.5 AQI & Speciation Mass,~
## $ CBSA_CODE           <int> NA, NA, NA, NA, NA, NA, NA, NA, NA, NA,~
## $ CBSA_NAME           <fct> "", "", "", "", "", "", "", "", "", "",~
## $ STATE_CODE          <int> 37, 37, 37, 37, 37, 37, 37, 37, 37, 37,~
## $ STATE               <fct> North Carolina, North Carolina, North C~
## $ COUNTY_CODE         <int> 11, 11, 11, 11, 11, 11, 11, 11, 11, 11,~
## $ COUNTY              <fct> Avery, Avery, Avery, Avery, Avery, Aver~
## $ SITE_LATITUDE       <dbl> 35.97235, 35.97235, 35.97235, 35.97235,~
## $ SITE_LONGITUDE      <dbl> -81.93307, -81.93307, -81.93307, -81.93~
```

```
glimpse(PM25.2019)
```

```
## Rows: 8,581
## Columns: 20
## $ Date          <fct> 01/03/2019, 01/06/2019, 01/09/2019, 01/~
## $ Source        <fct> AQS, AQS, AQS, AQS, AQS, AQS, AQS, AQS,~
## $ Site.ID       <int> 370110002, 370110002, 370110002, 370110~
## $ POC           <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ Daily.Mean.PM2.5.Concentration <dbl> 1.6, 1.0, 1.3, 6.3, 2.6, 1.2, 1.5, 1.5,~
## $ UNITS         <fct> ug/m3 LC, ug/m3 LC, ug/m3 LC, ug/m3 LC,~
## $ DAILY_AQI_VALUE <int> 7, 4, 5, 26, 11, 5, 6, 6, 15, 7, 14, 20~
## $ Site.Name     <fct> Linville Falls, Linville Falls, Linvill~
## $ DAILY_OBS_COUNT <int> 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, ~
## $ PERCENT_COMPLETE <dbl> 100, 100, 100, 100, 100, 100, 100, 100, 100,~
## $ AQS_PARAMETER_CODE <int> 88502, 88502, 88502, 88502, 88502, 8850~
## $ AQS_PARAMETER_DESC <fct> Acceptable PM2.5 AQI & Speciation Mass,~
## $ CBSA_CODE      <int> NA, NA, NA, NA, NA, NA, NA, NA, NA, NA,~
## $ CBSA_NAME      <fct> "", "", "", "", "", "", "", "", "", "",~
## $ STATE_CODE     <int> 37, 37, 37, 37, 37, 37, 37, 37, 37, 37,~
## $ STATE          <fct> North Carolina, North Carolina, North C~
## $ COUNTY_CODE    <int> 11, 11, 11, 11, 11, 11, 11, 11, 11, 11,~
## $ COUNTY         <fct> Avery, Avery, Avery, Avery, Avery, Aver~
## $ SITE_LATITUDE  <dbl> 35.97235, 35.97235, 35.97235, 35.97235,~
## $ SITE_LONGITUDE <dbl> -81.93307, -81.93307, -81.93307, -81.93~
```

Wrangle individual datasets to create processed files.

3. Change date columns to be date objects.
4. Select the following columns: Date, DAILY_AQI_VALUE, Site.Name, AQS_PARAMETER_DESC, COUNTY, SITE_LATITUDE, SITE_LONGITUDE
5. For the PM2.5 datasets, fill all cells in AQS_PARAMETER_DESC with “PM2.5” (all cells in this column should be identical).
6. Save all four processed datasets in the Processed folder. Use the same file names as the raw files but replace “raw” with “processed”.

```
# 3 - fixed date columns of all datasets
ozone.2018$Date <- mdy(ozone.2018$Date)
ozone.2019$Date <- mdy(ozone.2019$Date)
PM25.2018$Date <- mdy(PM25.2018$Date)
PM25.2019$Date <- mdy(PM25.2019$Date)

# 4 - selecting relevant data
ozone.2018.processed <- ozone.2018 %>%
  select(c("Date", "DAILY_AQI_VALUE", "Site.Name", "AQS_PARAMETER_DESC", "COUNTY",
           "SITE_LATITUDE", "SITE_LONGITUDE"))

ozone.2019.processed <- ozone.2019 %>%
  select(c("Date", "DAILY_AQI_VALUE", "Site.Name", "AQS_PARAMETER_DESC", "COUNTY",
           "SITE_LATITUDE", "SITE_LONGITUDE"))

PM25.2018.processed <- PM25.2018 %>%
  select(c("Date", "DAILY_AQI_VALUE", "Site.Name", "AQS_PARAMETER_DESC", "COUNTY",
           "SITE_LATITUDE", "SITE_LONGITUDE"))
```

```

PM25.2019.processed <- PM25.2019 %>%
  select(c("Date", "DAILY_AQI_VALUE", "Site.Name", "AQ5_PARAMETER_DESC", "COUNTY",
           "SITE_LATITUDE", "SITE_LONGITUDE"))

# 5 - filled in the AQS Parameter Descr cells for PM25 datasets
PM25.2018.processed.1 <- PM25.2018.processed %>%
  mutate(AQS_PARAMETER_DESC = "PM2.5")

PM25.2019.processed.1 <- PM25.2019.processed %>%
  mutate(AQS_PARAMETER_DESC = "PM2.5")

# 6 - saved new datasets
write.csv(ozone.2018.processed, "Data/Processed/EPAair_03_NC2018_processed.csv")

write.csv(ozone.2019.processed, "Data/Processed/EPAair_03_NC2019_processed.csv")

write.csv(PM25.2018.processed.1, "Data/Processed/EPAair_PM25_NC2018_processed.csv")

write.csv(PM25.2019.processed.1, "Data/Processed/EPAair_PM25_NC2019_processed.csv")

```

Combine datasets

7. Combine the four datasets with `rbind`. Make sure your column names are identical prior to running this code.
8. Wrangle your new dataset with a pipe function (`%>%`) so that it fills the following conditions:
 - Include all sites that the four data frames have in common: “Linville Falls”, “Durham Armory”, “Leggett”, “Hattie Avenue”, “Clemmons Middle”, “Mendenhall School”, “Frying Pan Mountain”, “West Johnston Co.”, “Garinger High School”, “Castle Hayne”, “Pitt Agri. Center”, “Bryson City”, “Millbrook School” (the function `intersect` can figure out common factor levels - but it will include sites with missing site information...)
 - Some sites have multiple measurements per day. Use the split-apply-combine strategy to generate daily means: group by date, site name, AQS parameter, and county. Take the mean of the AQI value, latitude, and longitude.
 - Add columns for “Month” and “Year” by parsing your “Date” column (hint: `lubridate` package)
 - Hint: the dimensions of this dataset should be 14,752 x 9.
9. Spread your datasets such that AQI values for ozone and PM2.5 are in separate columns. Each location on a specific date should now occupy only one row.
10. Call up the dimensions of your new tidy dataset.
11. Save your processed dataset with the following file name: “EPAair_O3_PM25_NC1819_Processed.csv”

```

# 7 - combined datasets
ozone.PM25.allyears <- rbind(ozone.2018.processed, ozone.2019.processed, PM25.2018.processed.1,
                             PM25.2019.processed.1)

# 8

```

```
new.combined.dataset <- ozone.PM25.allyears %>%
  filter(Site.Name %in% c("Linville Falls", "Durham Armory", "Leggett", "Hattie Avenue",
    "Clemmons Middle", "Mendenhall School", "Frying Pan Mountain", "West Johnston Co.",
    "Garinger High School", "Castle Hayne", "Pitt Agri. Center", "Bryson City",
    "Millbrook School")) %>%
  group_by(Date, Site.Name, AQS_PARAMETER_DESC, COUNTY) %>%
  summarise(meanAQI = mean(DAILY_AQI_VALUE), meanlat = mean(SITE_LATITUDE), meanlong = mean(SITE_LONGITUDE))
  separate(Date, c("Y", "m", "d"))
```

'summarise()' has grouped output by 'Date', 'Site.Name', 'AQS_PARAMETER_DESC'.
You can override using the '.groups' argument.

```
# 9
combined.spread <- new.combined.dataset %>%
  pivot_wider(names_from = AQS_PARAMETER_DESC, values_from = meanAQI)

# 10
dim(combined.spread)
```

```
## [1] 8976    9
```

```
# 11
write.csv(combined.spread, "Data/Processed/EPAair_03_PM25_NC1819_Processed.csv")
```

Generate summary tables

12. Use the split-apply-combine strategy to generate a summary data frame. Data should be grouped by site, month, and year. Generate the mean AQI values for ozone and PM2.5 for each group. Then, add a pipe to remove instances where mean **ozone** values are not available (use the function `drop_na` in your pipe). It's ok to have missing mean PM2.5 values in this result.

13. Call up the dimensions of the summary dataset.

```
# 12
summary.03.PM25 <- combined.spread %>%
  group_by(Site.Name, m, Y) %>%
  summarise(meanO3AQI = mean(Ozone), meanpm25AQI = mean(PM2.5)) %>%
  drop_na(meanO3AQI)
```

'summarise()' has grouped output by 'Site.Name', 'm'. You can override using
the '.groups' argument.

```
# 13
dim(summary.03.PM25)
```

```
## [1] 182    5
```

14. Why did we use the function `drop_na` rather than `na.omit`?

Answer: We used 'drop_na' because we were only focused on "NAs" in one column. 'na.omit' is typically used for removing "NAs" from entire dataframes.