

The centre of the even hybrid family quantum GL_n

the statement for general n , and the case $n = 1$ proved*

Tamás Hausel

Abstract

A family quantum group carries, alongside the deformation parameter, independent toral parameters $t_1, \dots, t_n$. Its *even hybrid integral form* is the part of it that preserves every integral Verma lattice; the question this paper is about is what the centre of that form looks like. The answer proposed here is that the Harish–Chandra projection carries it isomorphically onto the Weyl-invariant part of a *Newton lattice* — the functions on the multiplicative grid that take integral values at every weight. Section 2 states this for GL_n . Sections 3 and 4 state and prove it for $n = 1$, where the root system is empty, the Weyl group is trivial, and the whole content is a statement about integer-valued Laurent polynomials on a geometric grid: a bilateral form of the quantum Pólya theorem of Harman and Hopkins [5]. Section 5 records what was searched for in the literature and what was found.

1 Introduction

Two things distinguish the family quantum group from the usual one. The toral parameters $t_1, \dots, t_n$ are free, so the algebra is a family over a torus rather than a single object; and one takes not the whole algebra but the *hybrid integral form* — the elements preserving every integral Verma lattice — and inside that, the torally even part. The centre of that form is the object of study.

The shape of the answer is the same in every rank, and it is worth saying before any notation. A central element is determined by what it does to a highest-weight vector, weight by weight; that assignment is a function on a grid of weights; and the condition of being *integral* — of landing in A_t at every weight rather than merely in the fraction field — cuts out a lattice of such functions. *The theorem is that the centre is exactly that lattice of integer-valued functions, cut down by the Weyl symmetry.* Nothing is lost and nothing extra appears.

The lattice in question has a long classical ancestry. That the integer-valued polynomials on $\mathbb{Z}$ are free on the binomial coefficients is Pólya’s basis theorem (we reach the original through its quotation as [5, Prop. 1.1]; the standard modern reference for the subject is Cahen–Chabert [1]). On a *geometric* progression — the grid that appears here — the corresponding basis is due to Gramain [3, Prop. 2.2], for q a natural number; the q -analogue on the additive grid, with q an indeterminate, is Harman–Hopkins [5]. What the present setting adds is recorded in Section 5.

*The statement for $n = 1$ was fixed and signed before any proof was attempted, and the proof was checked against the signed text; this document was assembled from those two records by the Almanac Editor of Project Sandbox — for the pilot edition, a software agent. The full record, including per-claim warrants and machine verification, is the accompanying almanac edition.

Section 2 states the result for GL_n in four clauses: a PBW basis, an intrinsic characterisation of the integral form, the Harish–Chandra isomorphism onto the invariant Newton lattice, and a coefficient-even refinement. **This general statement is not proved here.** It is the programme’s flagship, and it is stated so the rank-one case can be seen for what it is: not a curiosity, but the first instance, with the root system empty.

Sections 3 and 4 are the case $n = 1$, stated and then proved in full. With $\Phi = \emptyset$ and $W = \{1\}$ the algebra is a torus, the Harish–Chandra projection is the identity, and the three clauses reduce to one substantial assertion: the even Newton lattice is spanned over A_t by the shifted divided classes $\check{Y}^u \nu_r(\check{Y})$.

The rank-one case is small on purpose. It was carried from statement to proof to machine-checked formalisation so that the cost and the failure modes of doing so could be measured; nine of the twelve results in the accompanying edition carry a proof-kernel certificate. That apparatus is not the subject of this paper and is not described here.

2 The statement for GL_n

What follows is the programme’s flagship statement. It is stated, not proved. Its individual ingredients are classical: the triangular decomposition and the freeness of the toral part go back to Lusztig [7, §3.2]; quantum Verma modules and their contravariant forms are treated in De Concini–Procesi [2, §§17.1–17.4]; and over the fraction field the Harish–Chandra map is an isomorphism onto the invariants by De Concini–Procesi [2, §18.3] and Jantzen [6, Thm. 6.25 and §6.26] — Jantzen’s §6.6 already placing the image inside the *even* toral invariants. Integral forms assembled, as here, from a divided-power negative part and a toral lattice appear in Habiro–Lê [4, §8G]. What is new in the statement is the family parameter, the hybrid (Verma-preserving) characterisation, and above all the assertion that over A_t the centre is the full *integer-valued* lattice — not the monomial ring the generic statement would suggest.

Fix $n \geq 1$ and put

$$q = v^2, \quad \mathbb{A}_t = \mathbb{Z}[v, v^{-1}][t_1^{\pm 1}, \dots, t_n^{\pm 1}], \quad \mathbb{F}_t = \mathbb{Q}(v)[t_1^{\pm 1}, \dots, t_n^{\pm 1}],$$

$$\mathbb{A}_t^{\text{ev}} = \mathbb{Z}[q, q^{-1}][t_1^{\pm 1}, \dots, t_n^{\pm 1}], \quad \mathbb{F}_t^{\text{ev}} = \mathbb{Q}(q)[t_1^{\pm 1}, \dots, t_n^{\pm 1}],$$

with v an indeterminate and $t_1, \dots, t_n$ algebraically independent. Let $\varepsilon_1, \dots, \varepsilon_n$ be the standard basis of $\mathbb{Z}^n$, equipped with $(\varepsilon_a, \varepsilon_b) = \delta_{ab}$; put $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ ($1 \leq i < n$) and $\beta_{ab} = \varepsilon_a - \varepsilon_b$ ($1 \leq a < b \leq n$), and let $\Phi^+ = \{\beta_{ab} : 1 \leq a < b \leq n\} \subset \mathbb{Z}^n$ be the set of positive roots, listed from left to right in the order

$$(1, 2), (1, 3), \dots, (1, n), (2, 3), \dots, (n-1, n).$$

Arrays are written $\mathbf{a} = (a_{ab})_{a < b} \in \mathbb{Z}_{\geq 0}^{\Phi^+}$, with $\mathbf{0}$ the zero array. For Z the centre of an algebra we write $Z(-)$. Quantum integers and factorials are

$$[r]_v = \frac{v^r - v^{-r}}{v - v^{-1}}, \quad [r]_v! = \prod_{j=1}^r [j]_v, \quad [0]_v! = 1.$$

Definition 1 (Reduced double of family quantum GL_n). $\tilde{U}_{\mathbb{F}_t}$ is the $\mathbb{F}_t$ -algebra with invertible pairwise commuting generators K_a^+, K_a^- ($1 \leq a \leq n$) and generators e_i, F_i ($1 \leq i < n$), subject to

$$K_a^+ K_a^- = t_a, \quad K_a^{\pm} e_i (K_a^{\pm})^{-1} = v^{\pm(\varepsilon_a, \alpha_i)} e_i, \quad K_a^{\pm} F_i (K_a^{\pm})^{-1} = v^{\mp(\varepsilon_a, \alpha_i)} F_i,$$

$$\begin{aligned}
e_i F_j - F_j e_i &= \delta_{ij} (K_i^+ K_{i+1}^- - K_i^- K_{i+1}^+), \\
e_i e_j &= e_j e_i, \quad F_i F_j = F_j F_i \quad (|i-j| > 1), \\
e_i^2 e_j - (v + v^{-1}) e_i e_j e_i + e_j e_i^2 &= 0, \quad F_i^2 F_j - (v + v^{-1}) F_i F_j F_i + F_j F_i^2 = 0 \quad (|i-j| = 1).
\end{aligned}$$

Define interval root vectors by $e_{a,a+1} = e_a$, $F_{a,a+1} = F_a$ and, for $b > a + 1$,

$$e_{ab} = \frac{e_{a,b-1} e_{b-1,b} - v^{-1} e_{b-1,b} e_{a,b-1}}{v - v^{-1}}, \quad F_{ab} = F_{b-1,b} F_{a,b-1} - v F_{a,b-1} F_{b-1,b}.$$

For $\mathbf{a} \in \mathbb{Z}_{\geq 0}^+$ the divided and ordinary PBW words are

$$F^{(\mathbf{a})} = \prod_{a < b}^{\leftarrow} \frac{F_{ab}^{a_{ab}}}{[a_{ab} v]!}, \quad e^{\mathbf{a}} = \prod_{a < b}^{\rightarrow} e_{ab}^{a_{ab}},$$

where the arrows read the displayed root list backwards and forwards. Put $\nu(\mathbf{a}) = \sum_{a < b} a_{ab} \beta_{ab} \in \mathbb{Z}^n$, and for $\mu = \sum_a \mu_a \varepsilon_a \in \mathbb{Z}^n$ put $K_\mu^+ = \prod_a (K_a^+)^{\mu_a}$.

Definition 2 (Torally even part). $\tilde{U}_{\mathbb{F}_t}$ carries the $\mathbb{Z}^n/2\mathbb{Z}^n$ -grading determined on generators by

$$\deg K_a^+ = \varepsilon_a, \quad \deg K_a^- = -\varepsilon_a, \quad \deg K_\mu^+ = \mu, \quad \deg e_i = 0, \quad \deg F_i = -\alpha_i, \quad \deg t_a = 0;$$

every defining relation is homogeneous modulo $2\mathbb{Z}^n$ (in the mixed relation both toral terms have degree $-\alpha_i \equiv \alpha_i$, the degree of its left side), so the grading is well defined, and the degree-zero part is a subalgebra. The interval vectors are homogeneous with $\deg e_{ab} = 0$ and $\deg F_{ab} = -\beta_{ab}$, so $\deg F^{(\mathbf{a})} = -\nu(\mathbf{a})$. $U_{\mathbb{F}_t}$ is the degree-zero part of $\tilde{U}_{\mathbb{F}_t}$.

Definition 3 (Newton lattice; the hybrid form). Put

$$Y_a = q^{n-a} t_a^{-1} (K_a^+)^2 \in \tilde{U}_{\mathbb{F}_t}, \quad \text{equivalently} \quad (K_a^+)^2 = q^{a-n} t_a Y_a.$$

Granting the basis in clause (i) below, the Y_a are algebraically independent invertible elements of $U_{\mathbb{F}_t}$, so $\mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}]$ is a Laurent polynomial ring; for $\lambda \in \mathbb{Z}^n$ let

$$\text{ev}_\lambda : \mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}] \longrightarrow \mathbb{F}_t, \quad \text{ev}_\lambda(Y_a) = q^{\lambda_a + n - a} t_a^{-1},$$

be the $\mathbb{F}_t$ -algebra map. The Newton lattice is

$$\mathcal{N}_{\mathbb{A}_t} = \{ f \in \mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}] : \text{ev}_\lambda(f) \in \mathbb{A}_t \text{ for every } \lambda \in \mathbb{Z}^n \},$$

and the even hybrid form is the $\mathbb{A}_t$ -submodule

$$U_{\mathbb{A}_t} = \sum_{\mathbf{a}, \mathbf{b} \in \mathbb{Z}_{\geq 0}^+} F^{(\mathbf{a})} K_{\nu(\mathbf{a})}^+ \mathcal{N}_{\mathbb{A}_t} e^{\mathbf{b}} \subset U_{\mathbb{F}_t},$$

each summand torally even since $\deg(F^{(\mathbf{a})} K_{\nu(\mathbf{a})}^+) = 0$; only finite sums occur. The symmetric group S_n fixes $\mathbb{F}_t$ and permutes the variables $Y_1, \dots, Y_n$; the invariant Newton lattice is $(\mathcal{N}_{\mathbb{A}_t})^{S_n} = \{ f \in \mathcal{N}_{\mathbb{A}_t} : wf = f \text{ for every } w \in S_n \}$.

Definition 4 (Integral Verma lattices; Harish–Chandra projection). Let $\tilde{U}_{\mathbb{F}_t}^{\geq 0} \subset \tilde{U}_{\mathbb{F}_t}$ be the subalgebra generated by all $K_a^\pm$ and all e_i . For $\lambda \in \mathbb{Z}^n$ let $\chi_\lambda : \tilde{U}_{\mathbb{F}_t}^{\geq 0} \rightarrow \mathbb{F}_t$ be the $\mathbb{F}_t$ -algebra map with

$$\chi_\lambda(K_a^+) = v^{\lambda_a}, \quad \chi_\lambda(K_a^-) = t_a v^{-\lambda_a}, \quad \chi_\lambda(e_i) = 0,$$

writing $\mathbb{F}_t^{\chi_\lambda}$ for $\mathbb{F}_t$ viewed as a $\tilde{U}_{\mathbb{F}_t}^{\geq 0}$ -module through χ_λ , let $M_{\mathbb{F}_t}(\lambda) = \tilde{U}_{\mathbb{F}_t} \otimes_{\tilde{U}_{\mathbb{F}_t}^{\geq 0}} \mathbb{F}_t^{\chi_\lambda}$ be the induced module, with highest vector $m_\lambda = 1 \otimes 1$; its integral lattice is

$$M_{\mathbb{A}_t}(\lambda) = \sum_{\mathbf{a} \in \mathbb{Z}_{\geq 0}^{\Phi^+}} \mathbb{A}_t F^{(\mathbf{a})} K_{\nu(\mathbf{a})}^+ m_\lambda.$$

Granting the basis in clause (i) below, every element of $U_{\mathbb{F}_t}$ has a unique PBW coefficient on the row $\mathbf{a} = \mathbf{b} = \mathbf{0}$, and that coefficient lies in $\mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}]$; the Harish–Chandra projection is the $\mathbb{F}_t$ -linear map

$$\text{HC} : U_{\mathbb{F}_t} \longrightarrow \mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}]$$

retaining that coefficient.

Statement. (i) The PBW sums are direct: the monomials $F^{(\mathbf{a})}(K_1^+)^{\eta_1} \dots (K_n^+)^{\eta_n} e^{\mathbf{b}}$, for $\mathbf{a}, \mathbf{b} \in \mathbb{Z}_{\geq 0}^{\Phi^+}$ and $\eta \in \mathbb{Z}^n$, form an $\mathbb{F}_t$ -basis of $\tilde{U}_{\mathbb{F}_t}$; the degree-zero part decomposes as

$$U_{\mathbb{F}_t} = \bigoplus_{\mathbf{a}, \mathbf{b} \in \mathbb{Z}_{\geq 0}^{\Phi^+}} F^{(\mathbf{a})} K_{\nu(\mathbf{a})}^+ \mathbb{F}_t[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}] e^{\mathbf{b}};$$

the sum defining $U_{\mathbb{A}_t}$ is direct, $U_{\mathbb{A}_t}$ is an $\mathbb{A}_t$ -subalgebra of $U_{\mathbb{F}_t}$ with $\mathbb{F}_t \otimes_{\mathbb{A}_t} U_{\mathbb{A}_t} \simeq U_{\mathbb{F}_t}$, and each sum defining $M_{\mathbb{A}_t}(\lambda)$ is direct.

(ii) The even hybrid form is the common preserver of the integral Verma lattices:

$$U_{\mathbb{A}_t} = \{ x \in U_{\mathbb{F}_t} : x M_{\mathbb{A}_t}(\lambda) \subseteq M_{\mathbb{A}_t}(\lambda) \text{ for every } \lambda \in \mathbb{Z}^n \}.$$

(iii) The intrinsic centre satisfies $Z(U_{\mathbb{A}_t}) = Z(\tilde{U}_{\mathbb{F}_t}) \cap U_{\mathbb{A}_t}$, and Harish–Chandra projection restricts to an isomorphism of $\mathbb{A}_t$ -algebras

$$\text{HC} : Z(U_{\mathbb{A}_t}) \xrightarrow{\sim} (\mathcal{N}_{\mathbb{A}_t})^{S_n}.$$

(iv) With

$$\mathcal{N}_{\mathbb{A}_t^{\text{ev}}} = \{ f \in \mathbb{F}_t^{\text{ev}}[Y_1^{\pm 1}, \dots, Y_n^{\pm 1}] : \text{ev}_\lambda(f) \in \mathbb{A}_t^{\text{ev}} \text{ for every } \lambda \in \mathbb{Z}^n \}$$

and $\mathfrak{Z}_{\mathbb{A}_t^{\text{ev}}} = \{ C \in Z(U_{\mathbb{A}_t}) : \text{HC}(C) \in (\mathcal{N}_{\mathbb{A}_t^{\text{ev}}})^{S_n} \}$ the coefficient-even central form, HC restricts to an isomorphism of $\mathbb{A}_t^{\text{ev}}$ -algebras $\mathfrak{Z}_{\mathbb{A}_t^{\text{ev}}} \xrightarrow{\sim} (\mathcal{N}_{\mathbb{A}_t^{\text{ev}}})^{S_n}$, and multiplication induces $\mathbb{A}_t \otimes_{\mathbb{A}_t^{\text{ev}}} \mathfrak{Z}_{\mathbb{A}_t^{\text{ev}}} \xrightarrow{\sim} Z(U_{\mathbb{A}_t})$.

3 The case $n = 1$

Here $\Phi = \emptyset$, $W = \{1\}$, and the cocharacter lattice is $\Lambda = \mathbb{Z}$. The four clauses above collapse: there are no root vectors, so the PBW basis is the toral monomials alone; the Harish–Chandra projection is the identity; and the Weyl invariance is vacuous. What remains is clause (iii), and it is not vacuous at all.

Let $G = GL_1$: the root system is $\Phi = \emptyset$, the Weyl group is $W = \{1\}$, the cocharacter lattice is $\Lambda = \mathbb{Z}$, and $X_\Lambda = \mathbb{Z}$. Let

$$\mathbb{A}_t = \mathbb{Z}[v^{\pm 1}, t^{\pm 1}], \quad \mathbb{F}_t = \mathbb{Q}(v)[t^{\pm 1}], \quad q = v^2,$$

with v, t independent indeterminates.

Definition 5 (Hybrid family quantum GL_1). U is the F_t -algebra with invertible pairwise commuting generators $K_{n,+}, K_{n,-}$ ($n \in \mathbb{Z}$) subject to

$$K_{0,\pm} = 1, \quad K_{m+n,\pm} = K_{m,\pm} K_{n,\pm}, \quad K_{n,+} K_{n,-} = t^n.$$

There are no generators e_i, F_i , since $\Phi = \emptyset$. Writing $T^\pm := K_{1,\pm}$, the monomials $(T^+)^w$ ($w \in \mathbb{Z}$) form a F_t -basis of U .

Definition 6 (Even Newton lattice). A toral element is an element of $F_t[(T^+)^{\pm 1}]$; it is even when all its toral exponents are even — equivalently, it is a Laurent polynomial in $\check{Y} := (T^+)^2$. For $\chi \in X_\Lambda$ let ev_χ be the A_t -linear map on even toral elements determined by $\text{ev}_\chi(\check{Y}) = q^\chi$. The even Newton lattice is the A_t -submodule

$$\mathcal{N}^{\text{ev}} = \{ f \in F_t[\check{Y}^{\pm 1}] : \text{ev}_\chi(f) \in A_t \text{ for every } \chi \in \mathbb{Z} \}.$$

Definition 7 (Even hybrid family integral form). Since $\Phi^+ = \emptyset$, the even hybrid family integral form is the A_t -submodule

$$U^{\text{hyb, ev}} = \mathcal{N}^{\text{ev}} \subset U,$$

and its central part is $Z(U^{\text{hyb, ev}}) = U^{\text{hyb, ev}} \cap Z(U \otimes_{F_t} \text{Frac}(A_t))$.

Definition 8 (Harish–Chandra projection; shifted Weyl action). Every element of U is toral, so the Harish–Chandra projection HC is the identity map on U . With

$$Y_n = t^{-n} K_{n,+}^2 \quad (n \in \mathbb{Z}),$$

every even toral element is a Laurent polynomial in the Y_n , and the shifted Weyl action of W is the F_t -algebra action $w(Y_n) = Y_{wn}$; set $(\mathcal{N}^{\text{ev}})^W = \{ f \in \mathcal{N}^{\text{ev}} : wf = f \text{ for all } w \in W \}$.

Definition 9 (Newton divided classes). For $r \geq 0$ set

$$\nu_r(X) = \prod_{s=0}^{r-1} \frac{X - q^s}{q^r - q^s} \in \mathbb{Q}(v)[X], \quad \nu_0(X) = 1.$$

Statement. For $G = GL_1$, with $\Lambda = \mathbb{Z}$:

- (i) $U^{\text{hyb, ev}}$ is an A_t -subalgebra of U ;
- (ii) the Harish–Chandra projection restricts to an isomorphism of A_t -algebras

$$\text{HC} : Z(U^{\text{hyb, ev}}) \xrightarrow{\sim} (\mathcal{N}^{\text{ev}})^W;$$

$$(iii) \mathcal{N}^{\text{ev}} = \sum_{u \in \mathbb{Z}} \sum_{r \geq 0} A_t \check{Y}^u \nu_r(\check{Y}).$$

4 Proof of the case $n = 1$

Throughout, v and t are independent indeterminates and

$$A_t = \mathbb{Z}[v^{\pm 1}, t^{\pm 1}], \quad F_t = \mathbb{Q}(v)[t^{\pm 1}], \quad q = v^2.$$

Thus $A_t \subseteq F_t$ is a subring, and $\mathbb{Z}[q^{\pm 1}] = \mathbb{Z}[v^{\pm 2}] \subseteq \mathbb{Z}[v^{\pm 1}] \subseteq A_t$.

The algebra U . U is the F_t -algebra with invertible pairwise-commuting generators $K_{n,+}, K_{n,-}$ ($n \in \mathbb{Z}$) subject to $K_{0,\pm} = 1$, $K_{m+n,\pm} = K_{m,\pm} K_{n,\pm}$ and $K_{n,+} K_{n,-} = t^n$.

Writing $T^\pm = K_{1,\pm}$, the monomials $(T^+)^w$ ($w \in \mathbb{Z}$) form a F_t -basis; U is commutative. A *toral* element is an element of $F_t[(T^+)^\pm]$; it is *even* when it is a Laurent polynomial in

$$\check{Y} := (T^+)^2.$$

Even toral elements thus form the Laurent-polynomial algebra $F_t[\check{Y}^\pm]$.

Evaluations. For $\chi \in \mathbb{Z}$, let $\text{ev}_\chi: F_t[\check{Y}^\pm] \rightarrow F_t$ be the F_t -algebra homomorphism substituting the unit $q^\chi \in A_t$ for $\check{Y}$, i.e.

$$\text{ev}_\chi\left(\sum_u c_u \check{Y}^u\right) = \sum_u c_u q^{u\chi} \quad (c_u \in F_t).$$

It is the unique A_t -linear ring homomorphism with $\text{ev}_\chi(\check{Y}) = q^\chi$, and it fixes F_t (hence A_t) pointwise. The *even Newton lattice* is

$$\mathcal{N}^{\text{ev}} = \{f \in F_t[\check{Y}^\pm] : \text{ev}_\chi(f) \in A_t \text{ for every } \chi \in \mathbb{Z}\}.$$

Because $\Phi^+ = \emptyset$ at GL_1 , the *even hybrid family integral form* is $U^{\text{hyb, ev}} = \mathcal{N}^{\text{ev}} \subseteq U$, with central part $Z(U^{\text{hyb, ev}}) = U^{\text{hyb, ev}} \cap Z(U \otimes_{F_t} \text{Frac}(A_t))$. Every element of U is toral, so the Harish–Chandra projection HC is the identity map on U ; the Weyl group is $W = \{1\}$, acting trivially, so $(\mathcal{N}^{\text{ev}})^W = \mathcal{N}^{\text{ev}}$.

Newton divided classes. For $r \geq 0$,

$$\nu_r(X) = \prod_{s=0}^{r-1} \frac{X - q^s}{q^r - q^s} \in \mathbb{Q}(v)[X], \quad \nu_0(X) = 1.$$

Each ν_r has degree exactly r (its leading coefficient $\prod_{s=0}^{r-1} (q^r - q^s)^{-1}$ is a nonzero element of $\mathbb{Q}(v)$), so $\{\nu_0, \dots, \nu_d\}$ is a F_t -basis of the module $F_t[\check{Y}]_{\leq d}$ of even toral polynomials of degree $\leq d$.

We write $\mathbb{Z}[q^\pm]$ -*integral* for “lies in $\mathbb{Z}[q^\pm]$ ”. For $m \in \mathbb{Z}$ and $r \geq 0$ put

$$B_r(m) := \nu_r(q^m) = \prod_{s=0}^{r-1} \frac{q^m - q^s}{q^r - q^s}, \quad B_0(m) = 1.$$

4.1 Clause (i): $U^{\text{hyb, ev}}$ is an A_t -subalgebra

Proposition 1. $U^{\text{hyb, ev}} = \mathcal{N}^{\text{ev}}$ is an A_t -subalgebra of U . (This is the bilateral, t -spectator analogue of the fact that quantum integer-valued polynomials form a ring [5, §1].)

Proof. For each $\chi \in \mathbb{Z}$ the map $\text{ev}_\chi: F_t[\check{Y}^\pm] \rightarrow F_t$ is a ring homomorphism, and $A_t \subseteq F_t$ is a subring; hence the preimage $\text{ev}_\chi^{-1}(A_t)$ is a subring of $F_t[\check{Y}^\pm]$. It contains A_t , because ev_χ fixes A_t pointwise, so $\text{ev}_\chi^{-1}(A_t)$ is an A_t -subalgebra. By definition

$$\mathcal{N}^{\text{ev}} = \bigcap_{\chi \in \mathbb{Z}} \text{ev}_\chi^{-1}(A_t)$$

is an intersection of A_t -subalgebras of $F_t[\check{Y}^\pm]$, hence itself an A_t -subalgebra (it contains 1 since $\text{ev}_\chi(1) = 1 \in A_t$, and is closed under sums, products and A_t -scalars). As $F_t[\check{Y}^\pm] \subseteq U$, this is an A_t -subalgebra of U . $\square$

4.2 Clause (ii): the Harish–Chandra isomorphism is the identity

Proposition 2. *The Harish–Chandra projection restricts to an isomorphism of A_t -algebras $\text{HC}: Z(U^{\text{hyb, ev}}) \xrightarrow{\sim} (\mathcal{N}^{\text{ev}})^W$, and this isomorphism is the identity map of $\mathcal{N}^{\text{ev}}$. (The rank-one, $\Phi^+ = \emptyset$ degeneration of the even hybrid Harish–Chandra isomorphism established at GL_2 in [8]; the classical ancestor is the Harish–Chandra isomorphism itself, whose quantum-group form the GL_2 record cites.)*

Proof. The algebra U is commutative, being spanned by monomials in the pairwise commuting invertible generators $K_{n, \pm}$. Base change is compatible with commutativity, so $U \otimes_{F_t} \text{Frac}(A_t)$ is a commutative ring and equals its own centre. Therefore

$$Z(U^{\text{hyb, ev}}) = U^{\text{hyb, ev}} \cap Z(U \otimes_{F_t} \text{Frac}(A_t)) = U^{\text{hyb, ev}} \cap (U \otimes_{F_t} \text{Frac}(A_t)) = U^{\text{hyb, ev}} = \mathcal{N}^{\text{ev}},$$

the middle equality because $U \hookrightarrow U \otimes_{F_t} \text{Frac}(A_t)$ is injective — $F_t \hookrightarrow \text{Frac}(A_t) = \mathbb{Q}(v, t)$ and U is free over F_t — so $U^{\text{hyb, ev}} = \mathcal{N}^{\text{ev}} \subseteq U$ is literally a subset of the base-changed algebra. Every element of U is toral, so $\text{HC} = \text{id}_U$, and its restriction to $Z(U^{\text{hyb, ev}}) = \mathcal{N}^{\text{ev}}$ is $\text{id}_{\mathcal{N}^{\text{ev}}}$. Finally $W = \{1\}$ acts trivially, so $(\mathcal{N}^{\text{ev}})^W = \mathcal{N}^{\text{ev}}$. Hence

$$\text{HC}: Z(U^{\text{hyb, ev}}) = \mathcal{N}^{\text{ev}} \xrightarrow{\text{id}} \mathcal{N}^{\text{ev}} = (\mathcal{N}^{\text{ev}})^W$$

is the identity map of the A_t -algebra $\mathcal{N}^{\text{ev}}$ (an A_t -algebra by clause (i)), which is manifestly an isomorphism of A_t -algebras. $\square$

There is no content here beyond commutativity: at GL_1 the rational centre is everything, HC is the identity, and Weyl invariance is vacuous.

4.3 Clause (iii): the Newton presentation

Write

$$S := \sum_{u \in \mathbb{Z}} \sum_{r \geq 0} A_t \check{Y}^u \nu_r(\check{Y}) \subseteq F_t[\check{Y}^{\pm 1}]$$

for the A_t -submodule generated by the shifted divided classes $\check{Y}^u \nu_r(\check{Y})$.

Theorem 1 (Even Newton presentation for GL_1). $\mathcal{N}^{\text{ev}} = S = \sum_{u \in \mathbb{Z}} \sum_{r \geq 0} A_t \check{Y}^u \nu_r(\check{Y})$. (The bilateral quantum Pólya theorem of Harman–Hopkins [5, §§1,4], transported to the multiplicative grid; the precise dictionary is the Attribution remark below.)

The proof occupies the rest of this section: the grid lemma, the forward inclusion $S \subseteq \mathcal{N}^{\text{ev}}$, the windowed reverse inclusion for polynomials, and the Laurent-clearing step.

4.3.1 The grid values are $\mathbb{Z}[q^{\pm 1}]$ -integral

Lemma 1. *For all $m \in \mathbb{Z}$ and $r \geq 0$ one has $B_r(m) = \nu_r(q^m) \in \mathbb{Z}[q^{\pm 1}]$ (the Gaussian-binomial value pattern of [5, §1], extended to negative nodes as their §4 bilateral localization requires). Explicitly:*

- (a) if $0 \leq m < r$ then $B_r(m) = 0$;
- (b) if $m \geq r \geq 0$ then $B_r(m) = \begin{bmatrix} m \\ r \end{bmatrix}_q \in \mathbb{Z}[q]$;
- (c) if $m = -n$ with $n \geq 1$ then

$$B_r(-n) = (-1)^r q^{-nr - \binom{r}{2}} \begin{bmatrix} n + r - 1 \\ r \end{bmatrix}_q \in \mathbb{Z}[q^{\pm 1}].$$

Proof. (a) If $0 \leq m < r$, the factor of the product at $s = m$ has numerator $q^m - q^m = 0$, so $B_r(m) = 0$.

(b) Assume $m \geq r \geq 0$. Pulling q^s out of each numerator and denominator factor,

$$B_r(m) = \prod_{s=0}^{r-1} \frac{q^s(q^{m-s} - 1)}{q^s(q^{r-s} - 1)} = \prod_{s=0}^{r-1} \frac{q^{m-s} - 1}{q^{r-s} - 1}.$$

Re-indexing by $i = r - s$ (so i runs from 1 to r),

$$B_r(m) = \prod_{i=1}^r \frac{q^{(m-r)+i} - 1}{q^i - 1} = \left[\begin{matrix} m \\ r \end{matrix} \right]_q,$$

the Gaussian binomial coefficient. It lies in $\mathbb{Z}[q]$: this is standard from the q -Pascal recurrence $\left[\begin{matrix} m \\ r \end{matrix} \right]_q = \left[\begin{matrix} m-1 \\ r-1 \end{matrix} \right]_q + q^r \left[\begin{matrix} m-1 \\ r \end{matrix} \right]_q$ with $\left[\begin{matrix} m \\ 0 \end{matrix} \right]_q = 1$, by induction on m .

(c) Let $m = -n$, $n \geq 1$. In each factor write the numerator as $q^{-n} - q^s = -q^{-n}(q^{n+s} - 1)$ and the denominator as $q^r - q^s = q^s(q^{r-s} - 1)$. Then

$$B_r(-n) = \prod_{s=0}^{r-1} \frac{-q^{-n}(q^{n+s} - 1)}{q^s(q^{r-s} - 1)} = (-1)^r q^{-nr} q^{-\sum_{s=0}^{r-1} s} \prod_{s=0}^{r-1} \frac{q^{n+s} - 1}{q^{r-s} - 1}.$$

Here $\sum_{s=0}^{r-1} s = \binom{r}{2}$. For the remaining product, the denominators give $\prod_{s=0}^{r-1} (q^{r-s} - 1) = \prod_{i=1}^r (q^i - 1)$ and the numerators give $\prod_{s=0}^{r-1} (q^{n+s} - 1) = \prod_{j=n}^{n+r-1} (q^j - 1)$, so

$$\prod_{s=0}^{r-1} \frac{q^{n+s} - 1}{q^{r-s} - 1} = \frac{\prod_{j=n}^{n+r-1} (q^j - 1)}{\prod_{i=1}^r (q^i - 1)} = \frac{[n+r-1]_q! / [n-1]_q!}{[r]_q!} = \left[\begin{matrix} n+r-1 \\ r \end{matrix} \right]_q,$$

where $[k]_q = (q^k - 1)/(q - 1)$ and the common powers of $(q - 1)$ cancel. The Gaussian binomial $\left[\begin{matrix} n+r-1 \\ r \end{matrix} \right]_q \in \mathbb{Z}[q]$ by part (b). Collecting the prefactor gives

$$B_r(-n) = (-1)^r q^{-nr - \binom{r}{2}} \left[\begin{matrix} n+r-1 \\ r \end{matrix} \right]_q \in \mathbb{Z}[q^{\pm 1}]. \quad \square$$

4.3.2 Forward inclusion $S \subseteq \mathcal{N}^{\text{ev}}$

Lemma 2. *For every $u \in \mathbb{Z}$ and $r \geq 0$, $\check{Y}^u \nu_r(\check{Y}) \in \mathcal{N}^{\text{ev}}$. Hence $S \subseteq \mathcal{N}^{\text{ev}}$. (The transported forward half of [5, §1]: q -binomials are quantum integer-valued.)*

Proof. Fix u, r and $\chi \in \mathbb{Z}$. Since ev_χ is a ring homomorphism with $\text{ev}_\chi(\check{Y}) = q^\chi$,

$$\text{ev}_\chi(\check{Y}^u \nu_r(\check{Y})) = q^{u\chi} \nu_r(q^\chi) = q^{u\chi} B_r(\chi).$$

By Lemma 1, $B_r(\chi) \in \mathbb{Z}[q^{\pm 1}] \subseteq A_t$, and $q^{u\chi}$ is a unit of A_t ; so the product lies in A_t . As χ was arbitrary, $\check{Y}^u \nu_r(\check{Y}) \in \mathcal{N}^{\text{ev}}$. The lattice $\mathcal{N}^{\text{ev}}$ is an A_t -module (indeed an A_t -algebra, clause (i)), so it contains every A_t -combination of these generators, i.e. $S \subseteq \mathcal{N}^{\text{ev}}$. $\square$

4.3.3 Reverse inclusion for polynomials: windowed Newton interpolation

Lemma 3. *Let $g \in \mathcal{N}^{\text{ev}}$ be an even toral polynomial, $g \in F_t[\check{Y}]$, of degree $\leq d$. Then $g = \sum_{r=0}^d c_r \nu_r(\check{Y})$ with all $c_r \in A_t$. In particular $g \in \sum_{r \geq 0} A_t \nu_r(\check{Y}) \subseteq S$. (Pólya's windowed-interpolation argument as q -deformed in the proof of [5, §1], run on the multiplicative grid with t a spectator.)*

Proof. Because $\{\nu_0, \dots, \nu_d\}$ is a F_t -basis of $F_t[\check{Y}]_{\leq d}$, there are unique $c_0, \dots, c_d \in F_t$ with

$$g = \sum_{r=0}^d c_r \nu_r(\check{Y}).$$

Evaluate on the window $\chi = m$ for $m = 0, 1, \dots, d$. As ev_m is F_t -linear and $\text{ev}_m(\nu_r(\check{Y})) = B_r(m)$,

$$\text{ev}_m(g) = \sum_{r=0}^d B_r(m) c_r = \sum_{r=0}^d V[m, r] c_r, \quad V[m, r] := B_r(m) \quad (0 \leq m, r \leq d).$$

By Lemma 1, every entry $V[m, r] \in \mathbb{Z}[q^{\pm 1}]$; moreover V is *lower unitriangular*: $V[m, r] = B_r(m) = 0$ for $m < r$ (part (a)), and $V[m, m] = B_m(m) = \prod_{s=0}^{m-1} \frac{q^m - q^s}{q^m - q^s} = 1$. Hence $\det V = 1$, and the inverse V^{-1} is again lower unitriangular with entries in $\mathbb{Z}[q^{\pm 1}]$ (back-substitution never leaves $\mathbb{Z}[q^{\pm 1}]$, and by uniqueness this inverse coincides with the inverse taken over the larger ring F_t).

Writing $\mathbf{c} = (c_0, \dots, c_d)^\top$ and $\mathbf{v} = (\text{ev}_0(g), \dots, \text{ev}_d(g))^\top$, we have $\mathbf{v} = V\mathbf{c}$ in F_t^{d+1} , hence

$$\mathbf{c} = V^{-1}\mathbf{v}.$$

Since $g \in \mathcal{N}^{\text{ev}}$, each $\text{ev}_m(g) \in A_t$, so $\mathbf{v} \in A_t^{d+1}$; and V^{-1} has entries in $\mathbb{Z}[q^{\pm 1}] \subseteq A_t$. Therefore $\mathbf{c} = V^{-1}\mathbf{v} \in A_t^{d+1}$, i.e. every $c_r \in A_t$.

The t -spectator argument. The map ev_m substitutes only $\check{Y} \mapsto q^m = v^{2m}$; it does not touch v or t . Thus t never enters the matrix V , whose entries lie in $\mathbb{Z}[q^{\pm 1}]$. Concretely, expand each coordinate along the free $\mathbb{Z}[v^{\pm 1}]$ -basis of t -powers: $c_r = \sum_k c_{r,k} t^k$ with $c_{r,k} \in \mathbb{Q}(v)$, so that $\text{ev}_m(g) = \sum_k (\sum_r B_r(m) c_{r,k}) t^k$. Membership $\text{ev}_m(g) \in A_t = \mathbb{Z}[v^{\pm 1}, t^{\pm 1}]$ means that for each fixed k the t^k -coefficient $\sum_r B_r(m) c_{r,k}$ lies in $\mathbb{Z}[v^{\pm 1}]$ for all $m \in \{0, \dots, d\}$. For each fixed k this is precisely the unitriangular system above, over $\mathbb{Z}[q^{\pm 1}]$ with right-hand sides in $\mathbb{Z}[v^{\pm 1}]$; it gives $c_{r,k} \in \mathbb{Z}[v^{\pm 1}]$. Hence $c_r = \sum_k c_{r,k} t^k \in \mathbb{Z}[v^{\pm 1}, t^{\pm 1}] = A_t$, as claimed. This is the same conclusion reached above; we spell out the coefficient-wise reading to make the spectator role of t explicit.

Base change $\mathbb{Z}[q^{\pm 1}] \rightarrow \mathbb{Z}[v^{\pm 1}] \rightarrow A_t$. Two rings appear: the matrix V and its inverse live over $\mathbb{Z}[q^{\pm 1}] = \mathbb{Z}[v^{\pm 2}]$, whereas the coordinates c_r and the values $\text{ev}_m(g)$ live over $F_t = \mathbb{Q}(v)[t^{\pm 1}]$ — honest v (odd powers) may occur, so the c_r need not lie in $\mathbb{Q}(q)$. This causes no loss: the extension $\mathbb{Z}[q^{\pm 1}] \hookrightarrow \mathbb{Z}[v^{\pm 1}]$ is free with basis $\{1, v\}$, hence flat, and multiplication by a $\mathbb{Z}[q^{\pm 1}]$ -entry preserves each of the two grades (even/odd powers of v). The reverse computation $\mathbf{c} = V^{-1}\mathbf{v}$ therefore runs identically in each grade, so integrality of the values over $\mathbb{Z}[v^{\pm 1}, t^{\pm 1}]$ forces integrality of the coordinates over the same ring. Equivalently, and most directly: V^{-1} has entries in $\mathbb{Z}[q^{\pm 1}] \subseteq A_t$ and $\mathbf{v} \in A_t^{d+1}$, so $\mathbf{c} = V^{-1}\mathbf{v} \in A_t^{d+1}$. $\square$

4.3.4 Laurent clearing and conclusion

Proof of Theorem 1. The inclusion $S \subseteq \mathcal{N}^{\text{ev}}$ is Lemma 2. For $\mathcal{N}^{\text{ev}} \subseteq S$, let $f \in \mathcal{N}^{\text{ev}} \subseteq F_t[\check{Y}^{\pm 1}]$ be arbitrary. Choose $N \geq 0$ so large that $g := \check{Y}^N f \in F_t[\check{Y}]$ is an honest polynomial (take N at least the largest magnitude of a negative $\check{Y}$ -exponent occurring in f). For every $\chi \in \mathbb{Z}$, using that ev_χ is a ring homomorphism,

$$\text{ev}_\chi(g) = \text{ev}_\chi(\check{Y}^N) \text{ev}_\chi(f) = q^{N\chi} \text{ev}_\chi(f).$$

Here $q^{N\chi}$ is a unit of A_t and $\text{ev}_\chi(f) \in A_t$, so $\text{ev}_\chi(g) \in A_t$; thus $g \in \mathcal{N}^{\text{ev}} \cap F_t[\check{Y}]$. By Lemma 3, $g = \sum_r c_r \nu_r(\check{Y})$ with $c_r \in A_t$. Multiplying back by the unit $\check{Y}^{-N}$,

$$f = \check{Y}^{-N} g = \sum_{r \geq 0} c_r \check{Y}^{-N} \nu_r(\check{Y}) \in \sum_{u \in \mathbb{Z}} \sum_{r \geq 0} A_t \check{Y}^u \nu_r(\check{Y}) = S,$$

each summand being an A_t -multiple of the generator $\check{Y}^u \nu_r(\check{Y})$ with $u = -N$. Hence $\mathcal{N}^{\text{ev}} \subseteq S$, and with Lemma 2, $\mathcal{N}^{\text{ev}} = S$. $\square$

5 Precedents

This section records what was looked for in the literature, what was found, and what was not. It is included because a claim of novelty is only as good as the search behind it; one search returned nothing, which is a result, and is stated as one.

Clause (iii): the Pólya ancestry, in three layers

The classical layer is Pólya's basis theorem: the integer-valued polynomials on $\mathbb{Z}$ are free on the binomial coefficients. We hold it through its quotation as [5, Prop. 1.1]; the source states a *basis*, from which the strict containment $\mathbb{Z}[x] \subsetneq \text{Int}(\mathbb{Z})$ is immediate — but the strictness itself is not Pólya's statement, and we do not attribute it to him. The standard reference for the whole subject is Cahen–Chabert [1].

The multiplicative layer is Gramain [3, Prop. 2.2]: a basis for the functions taking integral values on a *geometric* progression. One distinction matters and is easy to miss: Gramain's q is a natural number ≥ 2 and integrality means values in $\mathbb{Z}$, whereas here q is an indeterminate and integrality means values in A_t . His grid is ours; his coefficient ring is not. (Cahen–Chabert treat the geometric progression once, in Chapter II, Exercise 15, attributing it to Gramain on the page; the exercise as printed carries two misprints — the second node of the product and the sign of an exponent — identified in the course of this search and recorded in the edition's errata cards.)

The q -analogue layer is Harman–Hopkins [5, §1, Props. 1.1–1.2]: one variable, additive grid, coefficient ring $\mathbb{Z}[q^{\pm 1}]$, no family parameter. Both the forward inclusion and the windowed reverse interpolation of Section 4 follow their §1. What is added here is the passage to the multiplicative grid $\check{Y} = q^x$ and to bilateral exponents $u \in \mathbb{Z}$, together with the Laurent clearing that makes the two-sided statement follow from the one-sided one; the substitution $\check{Y} = 1 + (q - 1)x$, which their §4 closing remark connects to the Cartan part of Lusztig's integral form, carries their grid node $x = [n]_q$ to $\check{Y} = q^n$ and their basis term to $\nu_k(\check{Y})$ term by term.

Clause (ii), and the general statement

Clause (ii) rests on the corresponding GL_2 statement proved earlier in this programme [8]. For the GL_n statement of Section 2 the generic precedents are as cited there: Lusztig [7, §3.2], De Concini–Procesi [2, §§17–18], Jantzen [6, §6], Habiro–Lê [4, §8G]. None of these treats the family parameter or the integer-valued lattice; over the fraction field the centre is the invariant *monomial* ring, and the content of the flagship is precisely that over A_t it is the strictly larger Newton lattice.

What was searched, and how

The citation cycle over the manuscript was run on 2026-08-09 and re-run over the whole library on 2026-08-13. Every result of Section 4 carries a citation; no result claims to

have no predecessor. One methodological finding is worth repeating, because it changed an answer: the first query run — the exact phrase *Harish–Chandra isomorphism* against the statement shelf — returned two entries, both irrelevant, where the shelf in fact held twelve. **A hit count over a phrase reports the shape of the query, not the shape of the shelf**, and a “none located” finding is worth only as much as the surface it was run against.

What was not found

For the coefficient-field bridge — the assertion that enlarging the ambient coefficient ring from F_t to $\text{Frac}(A_t)$ adds no new elements to the Newton lattice — **no predecessor was located**. The queries were *base change*, *faithfully flat*, *extension of scalars* and *localization*; every hit was unrelated. The nearest published statement, Vandermonde coefficient descent in Cahen–Chabert [1, Prop. I.3.1], was located on the later pass and is the closest thing to a predecessor this claim has. Harman–Hopkins [5, §4, Prop. 4.3] was assessed as a candidate and **rejected**, and the reason is worth stating because the two look alike: their proposition says the object *grows* under localization and identifies the enlargement exactly, while the claim here is that it does *not* grow. “Bigger, and here is how much bigger” is not evidence for “not bigger at all”, however much both statements wear the word *localization*; their localization also inverts the deformation parameter, where this one enlarges the coefficient field of the toral parameter. Adjacent, and adjacency is not ancestry.

Remark 1. *Clause (iii) is the quantum Pólya theorem of Harman and Hopkins, Quantum integer-valued polynomials (arXiv:1601.06110), in bilateral Laurent form. Their §1 Proposition (Prop. §1.2, **prop:qbinom**), the q -analogue of Pólya’s 1919 theorem, states that the ring $\mathcal{R}_q^+ = \{P \in \mathbb{Q}(q)[x] : P([n]_q) \in \mathbb{Z}[q] \ \forall n \in \mathbb{N}\}$ is freely generated as a $\mathbb{Z}[q]$ -module by*

the q -binomial coefficient polynomials $\begin{bmatrix} x \\ k \end{bmatrix}_q$ (the q -deformed Newton basis on the additive

*grid $x = [n]_q$, with denominator $q^{\binom{k}{2}}[k]_q!$ — not the classical Pólya $\binom{x}{k}$); their §4 Proposition (Prop. §4.3, **prop:localization**), $\mathcal{R}_q = \mathcal{R}_q^+ \otimes_{\mathbb{Z}[q]} \mathbb{Z}[q^{\pm 1}]$, is the bilateral localization. In the current notation their parameter q coincides with our q ($= v^2$), so their displayed formulas read verbatim on our side.*

The node-preserving affine substitution $\check{Y} = 1 + (q-1)x$ is exactly their §4 z -variable; it carries the grid node $x = [n]_q$ to $\check{Y} = q^n$ and carries each basis term $\begin{bmatrix} x \\ k \end{bmatrix}_q$ onto $\nu_k(\check{Y})$ term

by term — the factors $(q-1)$ cancel in every Newton ratio. (They remark further that on adjoining square roots $K^2 = z$, $v^2 = q$ — which is precisely the passage to our T^+ and our v — the ring $\mathcal{R}_q[K, v]$ is equivalent to the Cartan part of Lusztig’s integral form of $U_v(\mathfrak{sl}_2)$. Nothing below rests on that remark.) The values-in- $\mathbb{Z}[q^{\pm 1}]$ side matches values-in- A_t with t a spectator, since evaluation is t -coefficient-wise and $\mathbb{Q}(q) \cap \mathbb{Z}[v^{\pm 1}] = \mathbb{Z}[q^{\pm 1}]$.

The two-variable (GL_2) form of this equivalence is established in [8]: there the grid criterion agrees with membership in the A_t -span of the product Newton generators, and passes to the full Laurent even lattice. The GL_1 statement above is the rank-one specialization of that transport.

References

- [1] P.-J. Cahen and J.-L. Chabert, *Integer-Valued Polynomials*, Mathematical Surveys and Monographs **48**, American Mathematical Society, 1997.

- [2] C. De Concini and C. Procesi, *Quantum groups*, in: D-modules, Representation Theory, and Quantum Groups, Lecture Notes in Mathematics **1565**, Springer, 1993.
- [3] F. Gramain, *Fonctions entières d'une ou plusieurs variables complexes prenant des valeurs entières sur une progression géométrique*, in: Cinquante Ans de Polynômes / Fifty Years of Polynomials, Lecture Notes in Mathematics **1415**, Springer, 1990, pp. 123–137.
- [4] K. Habiro and T. T. Q. Lê, *Unified quantum invariants for integral homology spheres associated with simple Lie algebras*, Geom. Topol. **20** (2016), 2687–2835.
- [5] N. Harman and S. Hopkins, *Quantum integer-valued polynomials*, arXiv:1601.06110 (2016).
- [6] J. C. Jantzen, *Lectures on Quantum Groups*, Graduate Studies in Mathematics **6**, American Mathematical Society, 1996.
- [7] G. Lusztig, *Introduction to Quantum Groups*, Progress in Mathematics **110**, Birkhäuser, 1993.
- [8] Project Sandbox, *The centre of the even hybrid family quantum GL_2* (closure record of the signed goal, revision A1+R5), 2026.