Kinetics of *Escherichia coli* β-galactosidase with ONPG and its inhibition by IPTG: how initial-rate estimation and model choice shape K_m, V_{max} and K_i

Experiment 2: Studying Enzyme Kinetics using Spectrophotometry

Methods in Molecular and Cellular Biology, B241

Student: Almokhtar Aljarodi

Laboratory partner: Abdulrahman Khardali

Laboratory sessions: 7 September 2026 (Part II) and 11 September 2026 (Part III); report compiled 11 October 2026

Summary

Enzyme kinetics condenses an enzyme's response to substrate, inhibitor and its own amount into two constants, K_m and V_{max}, but the numbers obtained depend on how initial velocities are extracted from the measurements. *Escherichia coli* β-galactosidase acting on *o*-nitrophenyl-β-D-galactopyranoside (ONPG) was followed by stopping timed aliquots with Na_2CO_3 and reading the *o*-nitrophenol formed at 415 nm (ε = 5.22 mM^{-1} cm^{-1}, linear to 0.43 mM). A course-provided reference dataset was analysed for the Results; our own measurements were analysed by the same rules and compared in the Discussion. Initial velocity rose approximately in proportion to enzyme (R^2 = 0.97). Over 0.069 to 2.20 mM ONPG the velocity approached saturation: K_m = 0.31 mM and V_{max} = 0.48 µmol min^{-1} by Lineweaver–Burk, 0.36 mM and 0.49 µmol min^{-1} by nonlinear fitting. With 0.458 mM IPTG the apparent K_m rose 5.6-fold, inhibition grew as substrate fell, and a competitive model fitted both series better than a noncompetitive one; K_i was 0.10 mM by the course method and 0.10 to 0.50 mM across estimators, because five of six uninhibited velocities rest on an assumed zero-time absorbance. The observations are consistent with competitive inhibition by IPTG; the constants are model-dependent estimates, not measured quantities.

Objectives

To test whether the initial velocity of β-galactosidase is proportional to enzyme concentration, to determine K_m and V_{max} for ONPG hydrolysis, and to establish whether IPTG inhibits the enzyme competitively from its effect on the apparent K_m and V_{max} and the derived K_i, while examining how the choice of initial-rate window and of fitting method changes each of these estimates.

Introduction

Enzymes accelerate reactions by binding substrates at active sites, and the dependence of reaction velocity on substrate concentration is the most direct window on how they do it. Michaelis and Menten described this dependence by postulating a reversible enzyme–substrate complex that breaks down to product, which gives $V_0 = V_{max}[S]/(K_m + [S])$ (1, 2). $V_{max} = k_{cat}[E]_T$ is the limiting velocity when every active site is occupied and is therefore proportional to the amount of enzyme; K_m is the substrate concentration at half-maximal velocity and sets how much substrate the enzyme needs to approach it. Lineweaver and Burk rearranged the equation into a straight line, $1/V_0 = (K_m/V_{max})(1/[S]) + 1/V_{max}$, whose intercepts give the two constants (3). The transformation is the course's required method and the classical teaching representation, but it stretches the low-substrate points, so small errors in the smallest velocities dominate the fitted line (4); a direct fit of the untransformed velocities weights every point equally. This is one of two questions I wanted to examine: how far the constants depend on the mathematics applied to the same measurements. The other is more basic: V_0 is defined at t = 0 but must be estimated from a few stopped samples, and which samples are used, whether a line or a first-interval average is taken, and whether the unmeasured starting absorbance is assumed to be zero are decisions that each change K_m.

β-Galactosidase (EC 3.2.1.23), the *lacZ* product of *Escherichia coli,* is a homotetramer of 1023-residue subunits that hydrolyses the β-1,4 glycosidic bond of lactose and of synthetic β-galactosides (5, 6). The chromogenic substrate ONPG is cleaved to galactose and *o*-nitrophenol. At alkaline pH *o*-nitrophenol exists as the yellow *o*-nitrophenolate anion, whose absorbance near 415 to 420 nm follows the Beer–Lambert law, $A = \varepsilon cl$. Adding 1 M Na_2CO_3 to a timed aliquot both stops the reaction, by raising the pH far above the enzyme's working range, and develops the colour, so one reading gives the product formed up to that moment (7). A series of stopped samples builds a progress curve whose initial slope is V_0, the velocity before substrate depletion, product accumulation or enzyme instability slow the reaction. The stopped method needs only a fixed-wavelength spectrophotometer; its coarse 2-min time resolution is the origin of the analytical choices examined here.

Inhibitors reveal themselves by how they change K_m and V_{max}. A competitive inhibitor binds free enzyme at the active site and is displaced by substrate: it leaves V_{max} unchanged but raises the

apparent Michaelis constant to $K_{m,app} = K_m(1 + [I]/K_i)$, where K_i is the inhibitor's dissociation constant. A noncompetitive inhibitor inactivates a fraction of the enzyme whatever the substrate concentration, lowering V_{max} and leaving K_m unchanged (2). Isopropyl-β-D-thiogalactopyranoside (IPTG) is the thiogalactoside used as a gratuitous inducer of the *lac* operon because it induces the enzyme without being hydrolysed by it (8, 9). The same stability makes it a dead-end ligand of β-galactosidase: it occupies the galactose subsite of the active site, but its thioglycosidic bond is not cleaved, so it is expected to behave as a competitive inhibitor with K_i of the order of 0.1 mM (6, 10). The inducer of *lacZ* thus competes with the substrates of the enzyme it induces, the logic that underlies competitive drugs such as statins and many antivirals. Distinguishing the two mechanisms from data is less simple than textbook figures suggest: they differ only near saturation, and a competitive model fitted with a shared V_{max} cannot itself show that V_{max} was unchanged.

Here the ONP spectrum and standard curve were established, V_0 was measured against enzyme concentration at saturating ONPG and over a 32-fold range of ONPG with and without IPTG, and the analysis asks whether the velocities scale with enzyme, saturate with substrate and respond to IPTG as competition predicts, and how much each constant moves when the initial-rate rule or the fitting method changes.

Experimental Procedure

All procedures followed the Experiment 2 laboratory manual (11) except for the modifications listed below. Z buffer (0.06 M Na_2HPO_4, 0.04 M NaH_2PO_4, 0.01 M KCl, 0.001 M $MgSO_4$, 0.05 M β-mercaptoethanol, pH 7.0), ONPG (1.0 mg mL^{-1}, 3.3 mM), IPTG (1.31 mg mL^{-1}, 5.5 mM), 1 mM ONP and 1 M Na_2CO_3 were provided, together with β-galactosidase standardised by the teaching staff to give A_{415} = 1.20 to 1.30 after 10 min in the standard reaction. Reactions (12.0 mL, room temperature) were assembled as in the manual's tables with enzyme added last; 2.0 mL aliquots were stopped with 1.0 mL Na_2CO_3 at 2, 4, 6, 8 and 10 min and read at 415 nm against a Z-buffer blank in a GENESYS 10S UV-Vis spectrophotometer (1 cm path). Part I standards and the Part I blank contained Na_2CO_3 as specified in the manual.

Datasets and modifications. Two datasets were analysed: a course-provided reference dataset supplied by the laboratory instructor (Professor Samah Gadhoum; e-mail of 13 September 2026) and an experimental dataset collected by our laboratory group. The reference measurements (Part I spectrum, ONP standard curve and enzyme-dilution series; Part III substrate series with and without IPTG, tubes A to F) were used for the primary kinetic analysis in the Results, as instructed. Our experimental measurements (Part II tubes A to E on 7 September; Part III tubes A to E without IPTG and A to D with IPTG on 11 September) were analysed with the identical procedure and are examined in the Discussion; they are not combined with the reference measurements in any curve.

Deviations from the manual in our own sessions: Part I was performed by the teaching staff; in Part III, tube F was not performed and the IPTG tube E was discarded before measurement, so our inhibited series has four tubes. Our 70 readings (14 printouts) were transcribed and verified against the photographs (Table R1); every digit is kept, including the 10-min reading of inhibited tube B (1.106), which was excluded from that tube's regression; the whole tube was excluded from our inhibited fits (Discussion). The reference standard-curve table lists seven readings where the manual describes a blank and seven standards; the instructor confirmed that the first reading is the zero-concentration point (mapping in the Results).

Data analysis. Concentrations were calculated from the stated stock concentrations and the manual's volumes ([ONPG] = 3.3 mM × V/12.0 mL; [IPTG] = 5.5 mM × 1.0/12.0 = 0.458 mM; ONP standards 1 mM × V/3.0 mL). The manual asks for V_0 from the initial linear portion of each progress curve, with the points used identified. One rule was applied to all 31 curves: the least-squares line $A_{415} = a + bt$ was fitted to the 2, 4 and 6 min points and extended one point at a time while the next increment lay within 20% (or 0.010 A, whichever larger) of the increment predicted by the current line; $V_0 = b$. When the second and third increments both fell below the first by more than this tolerance, no linear window could be fitted and V_0 was taken as the average rate over the first interval, A_{415}(2 min)/2 min, which assumes $A_{415} = 0$ at t = 0 because no reading exists there (flagged † throughout). This origin-assumed estimate is an average rate over 0 to 2 min, not a measured instantaneous initial velocity, and may lie below or above the true V_0. Four alternative estimators (regression only; the 0 to 2 min average rate for every curve; all five points; a baseline-offset scenario with the t = 0 absorbance set to the median intercept of the linear curves, an assumption rather than a measurement) test its sensitivity. The extinction coefficient was the slope of A_{415} against [ONP] over the linear region chosen by the same increment rule. V_0 in ΔA_{415} min^{-1} was converted to μmol ONP min^{-1} per assay as $V_0/\varepsilon \times 1.5$ (the 2 mL aliquot was diluted with 1 mL stop reagent) × 12.0 mL (reaction volume); the manual's division by ε gives the cuvette rate in mM min^{-1}, and the two factors convert it to an amount per minute. K_m and V_{max} were obtained from the Lineweaver–Burk regression as required (V_{max} = 1/intercept, K_m = slope × V_{max}) and checked by unweighted nonlinear least squares on the velocity scale; K_i was calculated from $K_{m,app} = K_m(1 + [I]/K_i)$. Competitive, pure noncompetitive and mixed models were also fitted to both series simultaneously. All calculations were performed in R 4.3.3 with tidyverse (12); scripts, data and outputs are public (Data and Code Availability). Each condition was measured once, so the uncertainties quoted are regression standard errors, not replicate variability, and the alternative estimators are different answers to the same data, not a confidence interval.

Results

All values in this section are from the reference dataset: Part I to III measurements provided by the course (Prof. Samah Gadhoum, e-mail of 13 September 2026) for use in the Results. Our own measurements (experimental dataset, Table R1) are analysed in the Discussion.

1. Absorbance spectrum of o-nitrophenol. The spectrum of 0.333 mM ONP (manual tube 4: [ONP] = 1 mM × 1.0 mL/3.0 mL) peaked at 415 nm (A = 1.723; Table 1, Figure 1A); 410 to 420 nm lay within 2% of the maximum, so 415 nm was used for all kinetic measurements.

Table 1. Absorbance spectrum of 0.333 mM ONP in Z buffer/1 M Na_2CO_3 (2:1), read against the Z buffer/Na_2CO_3 blank (reference Part I).

Wavelength (nm)	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465
Absorbance	1.212	1.344	1.463	1.562	1.643	1.697	1.723	1.720	1.686	1.625	1.540	1.432	1.310	1.178	1.042	0.905	0.778

2. Standard curve of o-nitrophenol. The reference table reported seven readings (Table 2): the first (0) is the zero-concentration preparation (2.0 mL Z buffer + 1.0 mL Na_2CO_3), as the instructor confirmed, and the other six are manual standards 1 to 6 in order, anchored by the spectrum tube (standard 4, 1.723 at 415 nm, identical to the fifth reading). Standard 7 (0.667 mM) has no reported reading; whether it was unprepared or unreported is unknown, and no value was assumed. Absorbance was linear for the five standards from 0.083 to 0.433 mM, $A_{415} = 5.222[ONP] - 0.022$ ($R^2 = 0.9988$; Figure 1B), giving $\varepsilon = 5.22 \pm 0.10$ A mM^{-1} cm^{-1} (5.15 through the origin). The 0.533 mM standard (A = 3.24) lay 17% above the line and was excluded; including it would give $\varepsilon = 6.01$ and lower every V_{max} by 13%, leaving K_m and K_i unchanged.




Figure 1. ONP absorbs maximally at 415 nm and obeys the Beer–Lambert law to 0.43 mM. (A) Spectrum of 0.333 mM ONP; points, readings; line, spline; dashed line, maximum; shaded, within 2% of it. (B) Standard curve; filled points and line, calibration over the shaded range (equation in the text); ×, zero-concentration point (not fitted); open point, the excluded 0.533 mM standard, 17% above the line. Reference Part I data.

Table 2. ONP standard curve at 415 nm (reference Part I): reported readings, their assignment to the manual's preparations, [ONP] in the 3.0 mL cuvette (1 mM × V/3.0 mL) and use in the fit.

Reported tube	Manual preparation	ONP 1 mM (mL)	[ONP] in cuvette (mM)	A415	A415 / [ONP] (per mM)	Fitted
1	Blank	0	0	0	–	no (zero point)
2	Standard 1	0.25	0.0833	0.428	5.14	yes
3	Standard 2	0.5	0.1667	0.856	5.14	yes
4	Standard 3	0.75	0.25	1.24	4.96	yes
5	Standard 4	1	0.3333	1.723	5.17	yes
6	Standard 5	1.3	0.4333	2.256	5.21	yes
7	Standard 6	1.6	0.5333	3.236	6.07	no (+17%)

3. Enzyme dilution. At 2.20 mM ONPG, product formation was linear over an initial window in every tube (Figure 2A; Table 3). V_0 fell from 0.136 ΔA_{415} min^{-1} with undiluted enzyme to 0.0229 at 0.15 of that amount (5.9-fold for a 6.7-fold dilution). Sample calculation, tube A: 0.136/5.22 = 0.0260 mM min^{-1} in the cuvette, × 1.5 = 0.0391 mM min^{-1} in the reaction, × 12.0 mL = 0.469 µmol min^{-1}; relative rate of tube B = 0.1041/0.136 = 0.77. The relative rates (Table 3) were approximately proportional to enzyme (V_0 = 0.481 × E µmol min^{-1} through the origin, R^2 = 0.972; free intercept -0.0006, p = 0.98; Figure 2B), although the rate per unit enzyme ranged from 0.71 (tube D) to 1.21 (tube C), and tubes D and E gave nearly equal rates despite 1.7-fold more enzyme in D.




Figure 2. Initial velocity rose approximately in proportion to the amount of β-galactosidase. (A) Progress curves (A_{415} of the stopped samples), 2.20 mM ONPG, 1.00 to 0.15 relative enzyme (tubes A to E); lines, fits to the initial linear window (filled, used; open, excluded; dotted, extrapolation); V_0 values in Table 3. (B) V_0 against relative enzyme (letters, tubes); line through the origin, V_0 = 0.481 × E µmol min^{-1} (R^2 = 0.97); a free-intercept fit (intercept -0.0006, p = 0.98) coincides with it. n = 1 per tube; reference Part II data.

Table 3. Enzyme dilution (reference Part II): A_{415} of the stopped samples, points used, V_0 (ΔA_{415} min^{-1}; µmol ONP min^{-1} per assay, × 3.447), rate relative to tube A and rate per unit enzyme.

Tube	Enzyme (mL)	Rel. enzyme	2 min	4 min	6 min	8 min	10 min	Points used (min)	V0 (dA/min)	V0 (umol/min)	Rel. rate	Rate/enzyme
A	0.2	1	0.332	0.602	0.876	0.965	1.407	2-6	0.136	0.469	1	1
B	0.15	0.75	0.245	0.487	0.67	0.89	1.084	2-10	0.1041	0.359	0.77	1.02
C	0.1	0.5	0.171	0.321	0.499	0.592	0.736	2-6	0.082	0.283	0.6	1.21
D	0.05	0.25	0.07	0.11	0.167	0.229	0.262	2-6	0.0243	0.084	0.18	0.71
E	0.03	0.15	0.063	0.11	0.158	0.198	0.248	2-10	0.0229	0.079	0.17	1.12

4. Michaelis–Menten kinetics without inhibitor. Only the 2.20 mM tube remained linear to 10 min (V_0 = 0.132 ΔA_{415} min^{-1}); in tubes B to F the later increments fell below the first, so V_0 is the origin-assumed 0 to 2 min average rate (†; Figure 3A, Table 4). V_0 rose 5.6-fold over the 32-fold substrate range and flattened above 1 mM (Figure 4A). Sample calculation, tube B: [S] = 3.3 mM × 4.0/12.0 = 1.10 mM; V_0 = 0.101/5.22 × 1.5 × 12.0 = 0.348 µmol min^{-1}; 1/[S] = 0.909 mM^{-1}, $1/V_0$ = 2.87 min µmol^{-1}. Lineweaver–Burk regression (Figure 4B): $1/V_0$ = 0.658(1/[S]) + 2.097 (R^2 = 0.928), so V_{max} = 1/2.097 = 0.477 µmol min^{-1} and K_m = 0.658 × 0.477 = 0.31 mM; the nonlinear fit gave K_m = 0.36 ± 0.13 mM and V_{max} = 0.494 ± 0.059 µmol min^{-1} (Table 5).




Figure 3. Product formation slowed as substrate fell and when IPTG was present. Progress curves for 2.20 to 0.069 mM ONPG (tubes A to F) without inhibitor (A) and with 0.458 mM IPTG (B; smaller absorbance scale). Symbols as in Figure 2A; dashed from ×, origin-assumed 0 to 2 min average rate (A_{415} = 0 assumed at t = 0; tubes B to F in A, tube A in B). V_0 values in Table 4; reference Part III data.

Table 4. Substrate series without and with 0.458 mM IPTG (reference Part III): A_{415} of stopped samples, V_0 (ΔA_{415} min^{-1}; µmol min^{-1} per assay) and Lineweaver–Burk reciprocals (1/[S], mM^{-1}; $1/V_0$, min µmol^{-1}); †, origin-assumed 0 to 2 min average rate; others, regression.

Series	Tube	[S] (mM)	2 min	4 min	6 min	8 min	10 min	V0 (dA/min)	V0 (umol/min)	1/[S]	1/V0
no inhibitor	A	2.2	0.282	0.546	0.815	1.094	1.328	0.132	0.455	0.45	2.2
no inhibitor	B	1.1	0.202	0.383	0.516	0.628	0.723	0.101†	0.348†	0.91	2.87
no inhibitor	C	0.55	0.156	0.278	0.357	0.425	0.468	0.078†	0.269†	1.82	3.72
no inhibitor	D	0.275	0.11	0.172	0.211	0.24	0.301	0.055†	0.190†	3.64	5.28
no inhibitor	E	0.1375	0.116	0.148	0.166	0.171	0.18	0.058†	0.200†	7.27	5
no inhibitor	F	0.06875	0.047	0.067	0.077	0.084	0.09	0.0235†	0.081†	14.55	12.35

Series	Tube	[S] (mM)	2 min	4 min	6 min	8 min	10 min	V0 (dA/min)	V0 (umol/min)	1/[S]	1/V0
+ 0.458 mM IPTG	A	2.2	0.198	0.48	0.586	0.784	0.966	0.099†	0.341†	0.45	2.93
+ 0.458 mM IPTG	B	1.1	0.065	0.127	0.196	0.239	0.294	0.03275	0.113	0.91	8.86
+ 0.458 mM IPTG	C	0.55	0.045	0.086	0.128	0.154	0.187	0.02075	0.072	1.82	13.98
+ 0.458 mM IPTG	D	0.275	0.029	0.051	0.068	0.085	0.099	0.0087	0.030	3.64	33.35
+ 0.458 mM IPTG	E	0.1375	0.02	0.035	0.047	0.06	0.069	0.00615	0.021	7.27	47.18
+ 0.458 mM IPTG	F	0.06875	0.011	0.019	0.025	0.031	0.036	0.0031	0.011	14.55	93.59

5. Michaelis–Menten kinetics with IPTG. With 0.458 mM IPTG every tube was slower (Figure 3B), by 25% at 2.20 mM ONPG and 87% at 0.069 mM, so inhibition grew as substrate fell (Figure 4A). The Lineweaver–Burk line steepened 9.5-fold, $1/V_0 = 6.24(1/[S]) + 3.55$ ($R^2 = 0.988$; Figure 4B), giving $K_{m,app}$ = 1.76 mM (5.6-fold increase) and $V_{max,app}$ = 0.282 µmol min^{-1} from an extrapolated intercept (3.55 ± 2.33). $K_i = [I]/(K_{m,app}/K_m - 1) = 0.458/(1.76/0.31 - 1) = 0.100$ mM. A competitive model fitted to both series (shared V_{max} = 0.53 ± 0.07 µmol min^{-1}, K_m = 0.44 ± 0.17 mM, K_i = 0.107 ± 0.042 mM) left a residual sum of squares 1.9 times smaller than a noncompetitive model with the same number of parameters, and a mixed model reduced to it (Table 5).




Figure 4. β-Galactosidase approached saturation with ONPG, and IPTG raised the apparent K_m. (A) V_0 against [ONPG] without (circles) and with 0.458 mM IPTG (triangles), from Figure 3; solid, competitive model fitted to both series (shared V_{max} assumed); dashed, Lineweaver–Burk hyperbolae; †, origin-assumed. (B) Lineweaver–Burk plot; x-intercepts $-1/K_m$ = -3.19, $-1/K_{m,app}$ = -0.57 mM^{-1}. Reference Part III data.

Table 5. Kinetic parameters (reference Part III; K_m, K_i in mM; V_{max} in µmol min^{-1} per assay; ± SE of the fit) by Lineweaver–Burk (course method) and nonlinear fits; app., apparent (with IPTG); n.d., no plateau, not determinable; global fits share V_{max} (competitive) or K_m (noncompetitive) between the series; RSS, residual sum of squares.

Condition	Method	Km	Vmax	Ki	Fit
No inhibitor	Lineweaver-Burk	0.314	0.477	–	R2 0.928
No inhibitor	Nonlinear MM	0.363 ± 0.128	0.494 ± 0.059	–	R2 0.919
+ 0.458 mM IPTG	Lineweaver-Burk	1.757 (app.)	0.282 (app.)	0.100	R2 0.988
+ 0.458 mM IPTG	Nonlinear MM	n.d.	n.d.	–	R2 0.962
Both series	Global, competitive	0.435 ± 0.166	0.529 ± 0.072	0.107 ± 0.042	RSS 0.0194
Both series	Global, noncompetitive	0.744 ± 0.382	0.626 ± 0.139	0.433 ± 0.181	RSS 0.0371

Discussion

The reference data behave as the Michaelis–Menten model predicts: V_0 rose approximately in proportion to enzyme at saturating substrate (R^2 = 0.97), velocity saturated with substrate, and IPTG raised the apparent K_m 5.6-fold while inhibiting most at low substrate. Much of the numerical answer was decided before any fit. Five of six uninhibited curves had no linear window, so their V_0 is a 0 to 2 min average with A_{415} = 0 assumed at t = 0; other defensible rules for the same readings move K_m from 0.31 to 0.55 mM and K_i from 0.10 to 0.50 mM (Table 5). Lineweaver–Burk weights the low-substrate points most, where the estimates are weakest (4); the nonlinear fit gave K_m = 0.36 ± 0.13 mM. A K_m reported without its estimation rule is incomplete.

The inhibition data fit a competitive model with a 1.9-fold lower residual sum of squares than a noncompetitive model of equal size. The fit does not show that V_{max} was unchanged: the model shares V_{max} by construction, and the inhibited series never approached saturation, so $V_{max,app}$ (0.28 µmol min^{-1}) is an extrapolation. The observations are therefore consistent with competitive inhibition, with $K_i \approx 0.1$ mM by the course method (0.11 mM by the global fit), of the same order as the 0.08 mM reported for purified enzyme (TES, pH 7.0, 25 °C) (10). K_m (0.31 to 0.36 mM) is two to three times the published 0.12 mM (10); origin-assumed rates understate V_0 if the rate was already falling, and room temperature, 2-min sampling and an uncertain effective ONPG concentration act the same way.

Our own measurements, analysed by the same rules, gave the same picture with a lower K_m, 0.16 mM by Lineweaver–Burk and 0.14 ± 0.03 mM by nonlinear fit (five tubes), because four of five uninhibited curves had linear windows. With IPTG, $K_{m,app}$ rose 8.8-fold (nonlinear, three tubes; $V_{max,app}$ ratio 1.01) and K_i = 0.059 mM, or 0.024 mM by Lineweaver–Burk. Inhibited tube B ran 1.57-fold faster than tube A at half the substrate and faster than its uninhibited control, which no inhibition model at fixed enzyme allows. A late or mistimed stop, a longer reaction time before the first aliquot, a pipetting difference in enzyme or ONPG, or omitted or diluted IPTG would each produce this; the record cannot distinguish them. It is kept in Table R1 and excluded from the fits (included, $K_{m,app}$ = 38 mM, intercept near zero). Tube E was discarded before measurement, so three inhibited points remain and our K_i is the less certain. Improvements: 30-s sampling in the first 2 min, ONPG from 0.03 to above 2 mM, duplicates, and verified ONPG and IPTG stocks.

Conclusion. Velocity was proportional to enzyme, saturated with ONPG (K_m = 0.31 mM, V_{max} = 0.477 µmol min^{-1} by Lineweaver–Burk) and was inhibited by IPTG as competition for the active site predicts ($K_i \approx 0.1$ mM). Observed: the rise of the apparent K_m and the growth of inhibition at low substrate. Inferred: the constants, the unchanged V_{max} and K_i, whose uncertainty is set by the models' assumptions, above all the estimate of V_0 from 2-min samples.

Data and Code Availability

The raw data (our 14 printouts and their transcription; the reference values as received), the R scripts, processed tables, figures and report sources are in the public repository https://github.com/almokhtar8-stack/exp2; Rscript R/run_all.R regenerates every number, table and figure. Instructor correspondence and course documents are not redistributed.

References


1. Michaelis, L., and Menten, M. L. (1913) Die Kinetik der Invertinwirkung, Biochem. Z. 49, 333-369.
2. Johnson, K. A., and Goody, R. S. (2011) The original Michaelis constant: translation of the 1913 Michaelis-Menten paper, Biochemistry 50, 8264-8269. DOI 10.1021/bi201284u
3. Lineweaver, H., and Burk, D. (1934) The determination of enzyme dissociation constants, J. Am. Chem. Soc. 56, 658-666. DOI 10.1021/ja01318a036
4. Dowd, J. E., and Riggs, D. S. (1965) A comparison of estimates of Michaelis-Menten kinetic constants from various linear transformations, J. Biol. Chem. 240, 863-869. DOI 10.1016/S0021-9258(17)45254-9
5. Jacobson, R. H., Zhang, X.-J., DuBose, R. F., and Matthews, B. W. (1994) Three-dimensional structure of β-galactosidase from E. coli, Nature 369, 761-766. DOI 10.1038/369761a0
6. Juers, D. H., Heightman, T. D., Vasella, A., McCarter, J. D., Mackenzie, L., Withers, S. G., and Matthews, B. W. (2001) A structural view of the action of Escherichia coli (lacZ) β-galactosidase, Biochemistry 40, 14781-14794. DOI 10.1021/bi011727i
7. Miller, J. H. (1972) Experiments in Molecular Genetics, pp 352-355, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
8. Jacob, F., and Monod, J. (1961) Genetic regulatory mechanisms in the synthesis of proteins, J. Mol. Biol. 3, 318-356. DOI 10.1016/S0022-2836(61)80072-7
9. Herzenberg, L. A. (1959) Studies on the induction of β-galactosidase in a cryptic strain of Escherichia coli, Biochim. Biophys. Acta 31, 525-538. DOI 10.1016/0006-3002(59)90029-0
10. Juers, D. H., Rob, B., Dugdale, M. L., Rahimzadeh, N., Giang, C., Lee, M., Matthews, B. W., and Huber, R. E. (2009) Direct and indirect roles of His-418 in metal binding and in the activity of β-galactosidase (E. coli), Protein Sci. 18, 1281-1292. DOI 10.1002/pro.140
11. Biological and Environmental Science and Engineering Division, KAUST. (2026) Methods in Molecular and Cellular Biology B241 Laboratory Manual, Experiment 2: Studying Enzyme Kinetics using Spectrophotometry, pp 1-19.

12. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L. D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache, S. M., Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., and Yutani, H. (2019) Welcome to the tidyverse, J. Open Source Softw. 4, 1686. DOI 10.21105/joss.01686

VII. Questions – Answer on this form and submit with your lab report

Michaelis-Menten Kinetics

1. Sketch by hand a typical enzyme kinetic curve of [S] versus V. The maximum [S] should be about 4 times the Km. Draw lines at [S]=Km and V=Vmax. Make sure you get the slope at the origin ([S]=0) correct. Draw the asymptote (tangent line) at [S]=0.



The curve is the hyperbola V = Vmax[S]/(Km + [S]), drawn to [S] = 4 Km. It leaves the origin with slope Vmax/Km (the first-order region, V ≈ (Vmax/Km)[S]), passes V = Vmax/2 at [S] = Km, and bends towards the asymptote V = Vmax, reaching 0.8 Vmax at [S] = 4 Km; the tangent at the origin meets the line V = Vmax at [S] = Km.

2. What effect does altering Km have on Vmax?

None. Vmax = kcat[E]T depends on the turnover number and the amount of enzyme; Km only moves the curve along the [S] axis (a mutation may change both constants, but that is two changes, not a consequence of one).

3. What effect does altering [E] have on Km? Vmax?

Km is unchanged, because it is a ratio of rate constants for the enzyme–substrate pair; Vmax changes in proportion to [E], as Figure 2 shows at 2.20 mM ONPG (5.9-fold for a 6.7-fold dilution, approximately proportional).

4. At what concentration of [S] is the initial velocity proportional to [E]?

At every [S], provided [E] is much smaller than [S] so that the steady state holds: V_0 = kcat[E][S]/(Km + [S]) is linear in [E] for any fixed [S]. The proportionality constant depends on [S], so enzyme is assayed at saturating substrate ([S] much greater than Km), where $V_0 \approx$ Vmax = kcat[E] and small errors in [S] do not matter; that is why Part II used 2.20 mM ONPG.

Competitive Inhibition

5. What effect does increasing [I] have on the apparent Vmax and Km?

Apparent Vmax is unchanged, because sufficient substrate displaces the inhibitor; apparent Km increases linearly with [I], Km,app = Km(1 + [I]/Ki). Here 0.458 mM IPTG raised the apparent Km 5.6-fold.

6. What is the effect on the Km (apparent) when [I] = Ki; when [I] = 2Ki?

Km,app = 2 Km when [I] = Ki and 3 Km when [I] = 2 Ki. (The observed 5.6-fold increase corresponds to [I]/Ki = 5.6 − 1 = 4.6, i.e. Ki = 0.458/4.6 = 0.100 mM.)

Non-competitive Inhibition

7. What effect does increasing [I] have on the apparent Vmax and Km?

Apparent Vmax falls as Vmax/(1 + [I]/Ki), because the inhibitor removes active enzyme regardless of [S]; apparent Km is unchanged, because the inhibitor binds E and ES equally and does not alter substrate affinity.

8. What is the effect on Vmax when [I] = Ki; [I] = 9 Ki?

Vmax,app = Vmax/2 when [I] = Ki and Vmax/10 when [I] = 9 Ki.

9. Can you distinguish noncompetitive from competitive inhibition using just the activity at low substrate concentrations? (i.e. the slope of V versus [S] at the origin?) Why or why not? How much does the slope change?

No. At low [S] the slope is Vmax,app/Km,app. For competitive inhibition it is Vmax/[Km(1 + [I]/Ki)]; for noncompetitive inhibition it is [Vmax/(1 + [I]/Ki)]/Km. Both equal (Vmax/Km)/(1 + [I]/Ki), so the slope falls by the same factor 1/(1 + [I]/Ki) in both cases (halved at [I] = Ki; to about 18% at the [I]/Ki ≈ 4.6 of this experiment). The two mechanisms differ only near saturation, where the competitive curve recovers Vmax and the noncompetitive one does not.

10. Given a series of enzyme preparations with differing activities, how can you distinguish between the presence of an inhibitor and varying amounts of the active enzyme? Explain, considering both competitive and non-competitive inhibitors.

Measure full V_0 versus [S] curves rather than single activities. Different amounts of active enzyme change Vmax in proportion and leave Km identical (Lineweaver–Burk lines share the x-intercept). A competitive inhibitor leaves Vmax and raises Km (lines share the y-intercept), which cannot be mimicked by enzyme amount. A noncompetitive inhibitor lowers Vmax at constant Km and so mimics less enzyme kinetically; to separate the two, dilute the preparation and re-assay (a reversible inhibitor's effect weakens as [I] falls, so specific activity rises, whereas a true enzyme difference gives constant specific activity), mix preparations (activities of inhibitor-free preparations add; an inhibitor makes the mixture sub-additive), remove small molecules by dialysis or gel filtration, or measure enzyme independently (protein assay, immunoblot or active-site titration with a tight-binding ligand such as PETG).


Raw Data

Our own instrument printouts (GENESYS 10S UV-Vis; 14 photographs, 70 readings, transcribed and verified value by value; Table R1) are appended below; the photographs reproduce the original printouts, which the course requires to be signed by a TA. The course-provided reference measurements used in the Results are reproduced as received in Tables 1 to 4 and in the repository (data/raw/reference/, transcribed from the course's files and cross-checked against them); our printout data are in data/raw/experimental/ and the photographs in report/raw_data_printouts/.

Table R1. Transcribed absorbance values from our printouts (experimental dataset; A_{415} of stopped samples; NA = sample discarded before measurement; Part III + IPTG tube B 10 min = 1.106 retained as read). Printout labels: Day 1 = Part II (7 Sep 2026); Day 2 Blue = Part III without IPTG; Day 2 Black = Part III with IPTG (11 Sep 2026).

Dataset	Printout label	Condition	Tube	2 min	4 min	6 min	8 min	10 min
Experimental Part II	Day 1	Enzyme dilution (Part II)	A	0.276	0.544	0.817	1.137	1.328
Experimental Part II	Day 1	Enzyme dilution (Part II)	B	0.180	0.354	0.533	0.694	0.860
Experimental Part II	Day 1	Enzyme dilution (Part II)	C	0.133	0.255	0.375	0.499	0.583
Experimental Part II	Day 1	Enzyme dilution (Part II)	D	0.065	0.121	0.176	0.235	0.283
Experimental Part II	Day 1	Enzyme dilution (Part II)	E	0.047	0.090	0.127	0.169	0.205
Experimental Part III (-IPTG)	Day 2 Blue	ONPG series, no inhibitor	A	0.259	0.480	0.696	0.938	1.061
Experimental Part III (-IPTG)	Day 2 Blue	ONPG series, no inhibitor	B	0.224	0.436	0.655	0.864	1.065
Experimental Part III (-IPTG)	Day 2 Blue	ONPG series, no inhibitor	C	0.200	0.388	0.549	0.742	0.880
Experimental Part III (-IPTG)	Day 2 Blue	ONPG series, no inhibitor	D	0.202	0.386	0.551	0.674	0.764
Experimental Part III (-IPTG)	Day 2 Blue	ONPG series, no inhibitor	E	0.111	0.201	0.249	0.301	0.346
Experimental Part III (+IPTG)	Day 2 Black	ONPG series + IPTG	A	0.163	0.320	0.476	0.628	0.780
Experimental Part III (+IPTG)	Day 2 Black	ONPG series + IPTG	B	0.269	0.506	0.748	0.995	1.106
Experimental Part III (+IPTG)	Day 2 Black	ONPG series + IPTG	C	0.080	0.158	0.244	0.304	0.362
Experimental Part III (+IPTG)	Day 2 Black	ONPG series + IPTG	D	0.041	0.082	0.116	0.151	0.188
Experimental Part III (+IPTG)	Day 2 Black	ONPG series + IPTG	E	–	–	–	–	–


Day 1 (7 Sep 2026, Part II, tubes A to E):



TEST SETUP
GENESYS 10S UV-Vis v4.006 2L6S357212

Advanced A-%T-C	9:45pm 7Sep26
Test Name	_Default_600
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	On
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.276
2	0.544
3	0.817
4	1.137
5	1.328




B

TEST SETUP
GENESYS 10S UV-Vis v4.006 2L6S357212

Advanced A-%T-C	9:54pm 7Sep26
Test Name	_Default_600
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	On
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.180
2	0.354
3	0.533
4	0.694
5	0.860




C

TEST SETUP
GENESYS 10S UV-Vis v4.006 2L6S357212

Advanced A-%T-C	9:58pm 7Sep26
Test Name	_Default_600
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	On
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.133
2	0.255
3	0.375
4	0.499
5	0.583



D

TEST SETUP
GENESYS 10S UV-Vis v4.006 2L6S357212

Advanced A-%T-C	10:04pm 7Sep26
Test Name	_Default_600
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	On
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.065
2	0.121
3	0.176
4	0.235
5	0.283




E

TEST SETUP
GENESYS 10S UV-Vis v4.006 2L6S357212

Advanced A-%T-C	10:07pm 7Sep26
Test Name	_Default_600
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	On
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.047
2	0.090
3	0.127
4	0.169
5	0.205

Day 2 (11 Sep 2026, Part III without IPTG, blue labels, tubes A to E):




A

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-%T-C	3:32am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.259
2	0.480
3	0.696
4	0.938
5	1.061




B

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-%T-C	3:36am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.224
2	0.436
3	0.655
4	0.864
5	1.065




C

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-%T-C	3:39am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.200
2	0.388
3	0.549
4	0.742
5	0.880




D

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-%T-C	3:46am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.202
2	0.386
3	0.551
4	0.674
5	0.764




E

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-%T-C	3:51am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.111
2	0.201
3	0.249
4	0.301
5	0.346


Day 2 (11 Sep 2026, Part III with IPTG, black labels, tubes A to D; tube E discarded before measurement):


A

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-XT-C	4:29am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.163
2	0.320
3	0.476
4	0.628
5	0.780


B

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-XT-C	4:25am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off


ID#	Abs 415.0nm
1	0.269
2	0.506
3	0.748
4	0.995
5	1.106


C

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-XT-C	4:33am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.080
2	0.158
3	0.244
4	0.304
5	0.362


D

TEST SETUP
GENESYS 10S UV-Vis v4.005 2L6R296203

Advanced A-XT-C	4:20am 11Sep26
Test Name	_Default_[Saved]
Measurement Mode	Absorbance
Wavelength	415.0nm
Ref. Wavelength Correction	Off
Delay Time (min:sec)	0:00
Sample Positioner	Auto 6
Number of Samples	5
Cell Correction	Off
ID# (0=OFF)	1
Low/High Limits	-9999/9999
Statistics	Off
Auto Print	Off
Auto Save Data	Off

ID#	Abs 415.0nm
1	0.041
2	0.082
3	0.116
4	0.151
5	0.188