

=====

FOR YOUR EYES ONLY

SUBJECT STRAIN LINE B1-F3-HY (Hypogeal 3)

=====

1. EXECUTIVE SUMMARY & IDENTIFICATION

This strain is an obligate calciphile parasitoid. It releases "drones" which attach directly to a host's central nervous system, allowing it to control the host and forming a localized hive-mind network controlled by the broodmother via pheromones. When an area has been sufficiently secured, the broodmother lays her eggs in the abdomen of her victims. After 3-8 hours, the larvae devour the victims from the inside out. Each larvae fixes itself to the cervical spine of its host.

BEHAVIOUR

This strain cannot survive in open or brightly lit environments, exhibiting photophobia. The parasite nests in subterranean areas, such as tunnels, crawlspaces, ventilation, and plumbing.

The parasite typically occupies a building undetected, only turning humans one at a time by secreting a substance

Histological analysis of the parasite's outer membrane reveals an arachnid-like chitinous respiratory structure. The organism is therefore likely sensitive to aerosolized synthetic pyrethroids and neuro-paralytic compounds, such as Raid. Upon killing the broodmother, all drones typically leave their human hosts within 24 hours.

EMERGENCY PROTOCOL

NEUTRALISATION:

1. FIND THE CAN OF BUG SPRAY.
2. PROCEED IMMEDIATELY TO THE NEST.
3. SHAKE CAN AND FOLLOW INSTRUCTIONS.

=====

END OF FILE

=====