



AL-MURABIT ISLAMIC
CHARITABLE FOUNDATION,
NIGERIA

**AL-MURABIT ISLAMIC CHARITABLE
PUBLICATIONS (AMICF PUBLICATIONS)**



**FEBRUARY 3, 2026
AMICF
NIGERIA**

Article Type: Research Article

Article History:

Received: 29th January, 2026

Accepted: 3rd February, 2026

Published: 10th February 2026

ISSN: To be assigned

DOI: To be assigned

Antioxidant Potential of *Moringa oleifera*: A Comprehensive Review of
Phytochemical Constituents, Mechanisms, and Therapeutic Applications

Author: Sulyman Habeeb

Affiliation: Department of Energy and Applied Chemistry, Usmanu Danfodiyo
University, Sokoto, Nigeria

Abstract

Moringa oleifera Lam., commonly known as the “miracle tree,” has emerged as a significant source of natural antioxidants with substantial potential for nutraceutical and pharmaceutical applications. This review synthesizes current knowledge on the phytochemical composition, antioxidant mechanisms, and factors influencing the antioxidant capacity of *M. oleifera* leaves and seeds. The plant's antioxidant activity is primarily attributed to polyphenols, flavonoids, phenolic acids, and vitamins, which function through radical scavenging, metal chelation, and modulation of endogenous antioxidant enzymes. Comparative analyses consistently demonstrate superior antioxidant potential in leaves compared to seeds, attributed to differential phytochemical profiles linked to their distinct physiological roles. Extraction parameters, environmental conditions, and post-harvest processing significantly influence antioxidant yield. This review highlights the therapeutic implications of *M. oleifera* antioxidants in mitigating oxidative stress-related pathologies and identifies research gaps requiring further investigation to optimize its industrial and clinical applications.

Keywords: *Moringa oleifera*, phytochemicals, polyphenols, oxidative stress, free radicals, functional foods, nutraceuticals

1. INTRODUCTION

The global burden of chronic diseases associated with oxidative stress—including cardiovascular diseases, cancer, diabetes, and neurodegenerative disorders has intensified research into natural antioxidant sources [1,2]. Synthetic antioxidants, while effective, raise concerns regarding safety and toxicity, driving interest in plant-derived alternatives [3,4].

Moringa oleifera Lam. (Moringaceae), a fast-growing deciduous tree native to India and now widespread across tropical and subtropical regions, represents one of the most promising candidates in this regard [5,6]. Recognized as the “miracle tree,” “drumstick tree,” or

“Mother’s Best Friend,” *M. oleifera* has been integral to traditional medicine systems for centuries [7,8].

Every part of *M. oleifera* leaves, seeds, pods, flowers, bark, and roots possesses nutritional and medicinal value [9]. However, leaves and seeds have garnered particular scientific attention due to their concentrated bioactive constituents and ease of utilization [10,11]. This review critically examines the antioxidant potential of these plant parts, exploring their phytochemical basis, mechanistic actions, and practical applications in health promotion and disease prevention.

2. PHYTOCHEMICAL COMPOSITION OF M. OLEIFERA

2.1 Leaf Constituents

M. oleifera leaves represent a nutritional powerhouse, containing proteins with all essential amino acids, vitamins A, C, and E, minerals (calcium, potassium, iron), and diverse secondary metabolites [12,13]. Phytochemical screening consistently identifies:

Flavonoids: Quercetin, kaempferol, and their glycosides dominate, contributing significantly to radical scavenging capacity [14,15]

Phenolic acids: Chlorogenic acid, gallic acid, and caffeic acid provide reducing power and metal chelation ability [16]

Vitamins: Ascorbic acid and tocopherols act as primary antioxidants [17]

Alkaloids and glucosinolates: Isothiocyanates derived from glucosinolates exhibit indirect antioxidant effects through phase II enzyme induction [18]

Qualitative analyses confirm the presence of saponins, tannins, and phenolic compounds, with total phenolic contents ranging 58.00–78.67 mg GAE/g and flavonoid contents ~31.73 mg QE/g in methanolic extracts [19,20].

2.2 Seed Constituents

Seeds contain bioactive peptides, oil rich in oleic acid (~70%), and phenolic compounds, albeit at lower concentrations than leaves [21,22]. Key constituents include:

Phenolic acids: Gentisic acid, gallic acid, and chlorogenic acid [23]

Flavonoids: Quercetin and kaempferol at reduced levels (~20.73 mg QE/g) [19]

Tocopherols: α -, γ -, and δ -tocopherols providing lipid-soluble antioxidant protection [24]


Bioactive peptides: Exhibiting direct radical scavenging and metal chelation properties [25]

The functional distinction between leaves and seeds photosynthetic versus reproductive/storage organs explains their divergent phytochemical profiles [26].

3. MECHANISMS OF ANTIOXIDANT ACTION

3.1 Direct Radical Scavenging

Plant antioxidants neutralize reactive oxygen species (ROS) and reactive nitrogen species (RNS) through hydrogen atom donation or single electron transfer, converting radicals to stable products [27,28]. The DPPH assay, measuring the reduction of purple DPPH• to yellow DPPH-H, exemplifies this mechanism:



Phenolic compounds, with their aromatic hydroxyl groups, are particularly effective hydrogen donors [29].

3.2 Metal Ion Chelation

Transition metals (Fe^{2+} , Cu^+) catalyze ROS formation through Fenton and Haber-Weiss reactions. Phenolic compounds chelate these

ions, preventing hydroxyl radical generation [30]: $\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \text{OH}^\bullet + \text{OH}^-$

3.3 Endogenous Antioxidant Modulation

M. oleifera phytochemicals upregulate antioxidant enzyme expression (superoxide dismutase, catalase, glutathione peroxidase) and enhance glutathione synthesis, bolstering cellular defense systems [31,32].

3.4 Synergistic Interactions

Hydrophilic antioxidants from leaves (phenolic acids, ascorbic acid) and lipophilic components from seeds (tocopherols, carotenoids) may produce synergistic effects, protecting both aqueous and lipid cellular compartments [33,34].

4. COMPARATIVE ANTIOXIDANT ACTIVITY:
LEAVES VERSUS SEEDS

consistently demonstrate superior antioxidant activity in leaf extracts (Table 1).

Multiple studies employing DPPH, FRAP, ABTS, and ORAC assays

Table 1. Comparative Antioxidant Parameters of *M. oleifera* Leaf and Seed Extracts

Parameter	Leaves	Seeds	Reference
Total Phenolic Content	58.00–78.67 mg GAE/g	17.67 mg GAE/g	[19]
Total Flavonoid Content	31.73 mg QE/g	20.73 mg QE/g	[19]
DPPH IC ₅₀	35.42 µg/mL	91.13 µg/mL	[19]
DPPH Inhibition (0.4 mg/mL)	82.67%	77.4%	[35]
TPC (mg GAE/g extract)	30.4	29.8	[35]

The 2.5-fold lower IC₅₀ for leaves indicates substantially higher radical scavenging efficiency. This superiority is attributed to:

Higher polyphenol density
required for photoprotection in
photosynthetic tissues [36]

Greater flavonoid diversity,
particularly quercetin and
kaempferol glycosides [14]

Elevated ascorbic acid content, a
potent water-soluble antioxidant
[17]

However, seeds offer advantages
in specific applications: their
lipophilic antioxidants provide
lipid peroxidation inhibition, and
their coagulant proteins enable
water purification with concurrent
antioxidant delivery [37,38].

5. FACTORS INFLUENCING ANTIOXIDANT CAPACITY

5.1 Extraction Parameters

Solvent polarity critically
determines extraction efficiency.
Polar solvents (methanol, ethanol,
ethyl acetate) preferentially extract
phenolic compounds, while

nonpolar solvents (hexane) isolate
lipid-soluble antioxidants [39,40].
Optimization of solvent
concentration, temperature, and
extraction time maximizes yield
[41].

5.2 Environmental Conditions

Altitude, temperature, UV
exposure, and soil composition
affect secondary metabolite
synthesis. Highland cultivation
increases flavonoid and phenolic
accumulation due to stress-induced
biosynthesis [42,43].

5.3 Post-Harvest Processing

Drying temperature, storage
duration, and light exposure
significantly impact antioxidant
stability. Heat-sensitive
compounds (ascorbic acid,
carotenoids) degrade at elevated
temperatures [44,45]. Freeze-
drying and dark storage preserve
antioxidant activity [46].

5.4 Plant Maturity

Young leaves exhibit higher antioxidant activity and phenolic content than mature leaves due to

active metabolic processes during growth [47].

6. THERAPEUTIC APPLICATIONS AND HEALTH IMPLICATIONS

6.1 Chronic Disease Prevention

M. oleifera antioxidants demonstrate protective effects against:

Cardiovascular disease: Inhibition of LDL oxidation, reduction of atherosclerotic plaque formation [48,49]

Diabetes: Attenuation of hyperglycemia-induced oxidative stress, protection of pancreatic β -cells [50,51]

Cancer: Prevention of DNA damage, induction of apoptosis in malignant cells [52,53]

Neurodegeneration: Reduction of neuronal oxidative damage in

Alzheimer's and Parkinson's models [54,55]

6.2 Nutraceutical and Food Industry Applications

M. oleifera leaf powder is incorporated into functional foods, beverages, and dietary supplements [56,57]. Seed oil ("Ben oil") finds application in cosmetic formulations due to its oxidative stability [58].

6.3 Water Treatment

Seed extracts' dual function as natural coagulants and antioxidants enables sustainable water purification with added health benefits [59,60].

7. CHALLENGES AND FUTURE DIRECTIONS

Despite promising findings, several challenges remain:

Standardization: Variability in cultivation conditions and processing methods necessitates quality control protocols [61]

Bioavailability: Limited data on absorption, metabolism, and tissue distribution of *M. oleifera* antioxidants in humans [62]

Clinical evidence: Most studies are preclinical; randomized controlled trials are needed to validate health claims [63]

Toxicological assessment: Long-term safety data for high-dose supplementation are incomplete [64]

Future research should prioritize:
Identification and quantification of individual bioactive compounds
Structure-activity relationship studies

Formulation development for enhanced bioavailability

Sustainable cultivation practices for consistent phytochemical profiles

8. CONCLUSION

Moringa oleifera represents a valuable source of natural antioxidants, with leaves demonstrating superior activity compared to seeds due to their enriched polyphenol and flavonoid content. The diverse mechanisms of antioxidant action radical scavenging, metal chelation, and enzyme modulation underpin its

therapeutic potential against oxidative stress-related diseases. Optimization of extraction and processing parameters, coupled with rigorous clinical validation, will facilitate the integration of *M. oleifera* into evidence-based nutraceutical and pharmaceutical applications, contributing to global health and sustainable development goals.

REFERENCES

- [1] Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010;4(8):118-126.
- [2] Halliwell B, Gutteridge JMC. Free radicals in biology and medicine. 5th ed. Oxford University Press; 2015.
- [3] Pisoschi AM, Pop A. The role of antioxidants in the chemistry of oxidative stress. *Eur J Med Chem.* 2015;97:55-74.
- [4] Nimse SB, Pal D. Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Adv.* 2015;5(35):27986-28006.
- [5] Fahey JW. *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. *Trees Life J.* 2005;1(5):1-33.
- [6] Iqbal S, Bhanger MI. Effect of season and production location on antioxidant activity of *Moringa oleifera* leaves grown in Pakistan. *J Food Compost Anal.* 2006;19(6-7):544-551.
- [7] Shindano J, Chitundu M. Acceptance of *Moringa* leaf powder as a nutrition supplement in Zambia. *J Nutr Educ Behav.* 2008;40(3):194-199.
- [8] Fuglie LJ. The miracle tree: *Moringa oleifera*: Natural nutrition for the tropics. CTA; 2001.
- [9] Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother Res.* 2007;21(1):17-25.
- [10] Makkar HPS, Becker K. Nutritional value and

antinutritional components of whole and ethanol extracted Moringa oleifera leaves. Anim Feed Sci Technol. 1996;63(1-4):211-228.

[11] Bennett RN, Mellon FA, Foidl N, et al. Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multi-purpose trees Moringa oleifera L. (horseradish tree) and Moringa stenopetala L. J Agric Food Chem. 2003;51(12):3546-3553.

[12] Gopalakrishnan L, Doriya K, Kumar DS. Moringa oleifera: A review on nutritive importance and its medicinal application. Food Sci Hum Wellness. 2016;5(2):49-56.

[13] Saini RK, Sivanesan I, Keum YS. Phytochemicals of Moringa oleifera: A review of their nutritional, therapeutic and industrial significance. 3 Biotech. 2016;6(2):203.

[14] Manguro LOA, Lemmen P. Phenolics of Moringa oleifera leaves. Nat Prod Res. 2007;21(1):56-68.

[15] Vongsak B, Sithisarn P, Mangmool S, Thongpraditchote S, Wongkrajang Y, Gritsanapan W. Maximizing total phenolics, total flavonoids contents and antioxidant activity of Moringa oleifera leaf extract by the appropriate extraction method. Ind Crops Prod. 2013;44:566-571.

[16] Sánchez-Moreno C, Larrauri JA, Saura-Calixto F. Free radical scavenging capacity and inhibition of lipid oxidation of wines, grape juices and related polyphenolic constituents. Food Res Int. 1999;32(6):407-412.

[17] Nambiar VS, Guin P, Parnami S, Daniel M. Impact of antioxidants from drumstick leaves on antioxidant enzyme status in adult,

postmenopausal women. *Int J Food Sci Nutr*. 2014;65(8):963-967.

[18] Waterman C, Cheng DM, Rojas-Silva P, et al. Stable, water extractable isothiocyanates from *Moringa oleifera* leaves attenuate inflammation in vitro.

Phytochemistry. 2014;103:114-122.

[19] Unuigbo CA, Okeri HA, Erharuyi O, Oghenero EE, Obamedo DA. Phytochemical and antioxidant evaluation of *Moringa oleifera* leaf and seed. *J Pharm Bioresour*. 2014;11(2):51-57.

[20] Kamaluddin IDK, Ramdhani PT, Bima IH, Irwan AA, Matoo RF. Comparative analysis of antioxidants in *Moringa* leaves and soursop leaves. *J Biol Trop*. 2024;24(2):81-88.

[21] Lalas S, Tsaknis J. Characterization of *Moringa oleifera* seed oil variety

'Periyakulam 1'. *J Food Compos Anal*. 2002;15(1):65-77.

[22] Abdulkarim SM, Long K, Lai OM, Muhammad SKS, Ghazali HM. Frying quality and stability of high-oleic *Moringa oleifera* seed oil in comparison with other vegetable oils. *Food Chem*. 2007;105(4):1382-1389.

[23] Sreelatha S, Padma PR. Antioxidant activity and total phenolic content of *Moringa oleifera* leaves in two stages of maturity. *Plant Foods Hum Nutr*. 2009;64(4):303-311.

[24] Tsaknis J, Lalas S, Gergis V, Dourtoglou V, Spiliotis V. Characterization of *Moringa oleifera* variety Mbololo seed oil of Kenya. *J Agric Food Chem*. 1999;47(11):4495-4499.

[25] Al-Asmari AK, Athar MT, Kadasah SG. An updated phytopharmacological review on medicinal plant of Arab region:

- Moringa peregrina (Forssk.) Fiori.
Biomed Res Int.
2015;2015:951021.
- [26] Vergara-Jimenez M,
Almatrafi MM, Fernandez ML.
Bioactive components in Moringa
oleifera leaves protect against
chronic disease. Antioxidants.
2017;6(4):91.
- [27] Prior RL, Wu X, Schaich K.
Standardized methods for the
determination of antioxidant
capacity and phenolics in foods and
dietary supplements. J Agric Food
Chem. 2005;53(10):4290-4302.
- [28] Huang D, Ou B, Prior RL. The
chemistry behind antioxidant
capacity assays. J Agric Food
Chem. 2005;53(6):1841-1856.
- [29] Rice-Evans CA, Miller NJ,
Paganga G. Structure-antioxidant
activity relationships of flavonoids
and phenolic acids. Free Radic Biol
Med. 1996;20(7):933-956.
- [30] Perron NR, Brumaghim JL. A
review of the antioxidant
mechanisms of polyphenol
compounds related to iron binding.
Cell Biochem Biophys.
2009;53(2):75-100.
- [31] Fard MT, Arulselvan P,
Karthivashan G, Adam SK,
Fakurazi S. Bioactive extract from
Moringa oleifera inhibits the pro-
inflammatory mediators in
lipopolysaccharide stimulated
macrophages. Pharmacogn Mag.
2015;11(Suppl 1):S556-S563.
- [32] Gupta R, Mathur M, Bajaj
VK, Katariya P, Yadav S, Kamal
R, Gupta RS. Evaluation of
antidiabetic and antioxidant
activity of Moringa oleifera in
experimental diabetes. J Diabetes.
2012;4(2):164-171.
- [33] Anwar F, Zafar SN, Rashid U.
Characterization of Moringa
oleifera seed oil variety in

Pakistan. Grasas Aceites.

2006;57(4):397-407.

[34] Leone A, Spada A, Battezzati A, Schmidlin A, Bertoli S, Battezzati A. Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves: An overview. *Int J Mol Sci*. 2015;16(6):12791-12835.

[35] Habeeb S. Comparative study of antioxidant activity of *Moringa* leave and seed [B.Sc. Project]. Usmanu Danfodiyo University Sokoto; 2025.

[36] Chumark P, Khunawat P, Sanvarinda Y, et al. The in vitro and ex vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *J Ethnopharmacol*. 2008;116(3):439-446.

[37] Ghebremichael KA, Gunaratna KR, Henriksson H,

Brumer H, Dalhammar G. A simple purification and activity assay of the coagulant protein from *Moringa oleifera* seed. *Water Res*. 2005;39(11):2338-2344.

[38] Ndabigengesere A, Narasiah KS, Talbot BG. Active agents and mechanism of coagulation of turbid waters using *Moringa oleifera*. *Water Res*. 1995;29(2):703-710.

[39] Horax R, Hettiarachchy N, Kannan A, Chen P. Protein extraction optimisation, characterisation, and functionalities of protein isolate from bitter melon (*Momordica charantia*) seed. *Food Chem*. 2010;124(2):545-550.

[40] Erkan N, Ayranci G, Ayranci E. Antioxidant activities of rosemary (*Rosmarinus officinalis* L.) extract, blackseed (*Nigella sativa* L.) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food Chem*. 2008;110(1):76-82.

- [41] Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *J Agric Food Chem.* 2003;51(8):2144-2155.
- [42] Purwanti R. Comparison of vitamin C levels in *Moringa* leaves from different altitudes. *J Permata Indones.* 2022;13(1):62-67.
- [43] Amaglo NK, Bennett RN, Lo Curto RB, et al. Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. *Food Chem.* 2010;122(4):1047-1054.
- [44] Sreelatha S, Padma PR. Antioxidant activity and total phenolic content of *Moringa oleifera* leaves in two stages of maturity. *Plant Foods Hum Nutr.* 2009;64(4):303-311.
- [45] Yang R, Chang L, Hsu J, et al. Effects of different drying methods on the antioxidant properties of *Moringa oleifera* leaves. *J Food Process Preserv.* 2018;42(11):e13778.
- [46] Abdalrahman A, Jusoh WAW, Zaki NAM. Effect of drying methods on total antioxidant capacity of *Moringa oleifera* leaf powder. *IOP Conf Ser Mater Sci Eng.* 2018;358:012014.
- [47] Amaglo NK, Bennett RN, Lo Curto RB, et al. Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. *Food Chem.* 2010;122(4):1047-1054.
- [48] Chumark P, Khunawat P, Sanvarinda Y, et al. The in vitro and ex vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera*

Lam. leaves. J Ethnopharmacol. 2008;116(3):439-446.

[49] Randhir R, Lin YT, Shetty K. Stimulation of phenolics, antioxidant and antimicrobial activities in dark germinated mung bean sprouts in response to peptide and phytochemical elicitors. Process Biochem. 2004;39(5):637-646.

[50] Ndong M, Uehara M, Katsumata S, Suzuki K. Effects of oral administration of Moringa oleifera Lam on glucose tolerance in Goto-Kakizaki and Wistar rats. J Clin Biochem Nutr. 2007;40(3):229-233.

[51] Jaiswal D, Kumar Rai P, Kumar A, Mehta S, Watal G. Effect of Moringa oleifera Lam. leaves aqueous extract therapy on hyperglycemic rats. J Ethnopharmacol. 2009;123(3):392-396.

[52] Sreelatha S, Jeyachitra A, Padma PR. Antiproliferation and induction of apoptosis by Moringa oleifera leaf extract on human cancer cells. Food Chem Toxicol. 2011;49(6):1270-1275.

[53] Bose CK. Possible role of Moringa oleifera Lam. root in epithelial ovarian cancer. MedGenMed. 2007;9(1):26.

[54] Ekong MB, Akpanabiatu MI, Umoh IB, Eka OU, Edet EE, Akpaso MI. Antiperoxidative effect of Moringa oleifera leaf extract in rats. J Med Food. 2011;14(3):310-312.

[55] Ganguly R, Guha D. Alteration of brain monoamines & EEG wave pattern in rat model of Alzheimer's disease & protection by Moringa oleifera. Indian J Med Res. 2008;128(6):744-751.

[56] Soliva CR, Kreuzer M, Foidl N, Foidl G, Machmüller A, Hess HD. Feeding value of Moringa

oleifera leaves as a substitute for concentrate feed in dairy cows. Livest Res Rural Dev. 2005;17(9):89.

[57] Nambiar VS, Bhadalkar K, Daxini M. Drumstick leaves as source of vitamin A in ICDS-SFP. Indian J Pediatr. 2003;70(5):383-387.

[58] Anwar F, Rashid U. Physico-chemical characteristics of Moringa oleifera seeds and seed oil from a wild provenance of Pakistan. Pak J Bot. 2007;39(5):1443-1453.

[59] Ghebremichael KA, Gunaratna KR, Henriksson H, Brumer H, Dalhammar G. A simple purification and activity assay of the coagulant protein from Moringa oleifera seed. Water Res. 2005;39(11):2338-2344.

[60] Kansal SK, Kumari A. Potential of M. oleifera for the treatment of water and wastewater.

Chem Rev. 2014;114(10):4993-5010.

[61] Amaglo NK, Bennett RN, Lo Curto RB, et al. Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree Moringa oleifera L., grown in Ghana. Food Chem. 2010;122(4):1047-1054.

[62] Vergara-Jimenez M, Almatrafi MM, Fernandez ML. Bioactive components in Moringa oleifera leaves protect against chronic disease. Antioxidants. 2017;6(4):91.

[63] Stohs SJ, Hartman MJ. Review of the safety and efficacy of Moringa oleifera. Phytother Res. 2015;29(6):796-804.

[64] Oyeyinka AT, Oyeyinka SA. Moringa oleifera as a food fortificant: Recent trends and prospects. J Saudi Soc Agric Sci. 2018;17(2):171-180.

Conflict of Interest: The author declares no conflict of interest.