

SxIO

solid bedrock, intermediate siliceous rocks, impure

General parameters

Area	461.57 km ²
Percentage on total forest mapped area	9.5 %

Physics - mean values of all considered profiles (124)

Depth [cm]	Coarse fraction [%]	Field capacity [l/m ²]
0-15	25 ± 25	69 ± 41
15-30	45 ± 30	
30-60	55 ± 30	
60-100	70 ± 25	

Chemistry - stock of available profiles (5)

Ctot	Ntot	Ca	Mg	K	P
t/ha	t/ha	kg/ha	kg/ha	kg/ha	kg/ha
112.44	5.45	739.38	158.74	154.26	925.51

All stock values, 0-80 cm including humus layers (F,H), are short term available, except for phosphorus, which has long term availability

Chemistry - mean values of all considered profiles (31)

Depth [cm]	CEC [mmol/kg]	Base Saturation [%]	(Mg+Ca)/CEC	Ntot [%]	TOC [%]	C/N	pH _{CaCl2}
0-5	133.87	30.66	0.29	0.41	8.71	21.24	3.57
5-10	123.11	29.3	0.27	0.38	7.8	20.53	3.63
10-20	67.18	19.68	0.17	0.2	3.47	17.35	4.01
20-40	41.88	29.28	0.25	0.14	2.54	18.14	4.31
40-80	45.39	36.22	0.31	0.12	2.15	17.92	4.36





Figure 1: Profile's depth variation of the following median chemical properties, bounded by 25th and 75th percentiles: cation exchange capacity (mmol/kg), nitrogen (%), total organic carbon (%), pH and base saturation (%). The percentage values indicate how many profiles contribute to the median calculation at each depth step.

Biomass use

Effects of whole-tree harvesting



Intermediate negative effects

Compaction risk

Effects of the transit of heavy-duty machinery



Occasionally critical