

# FxI-

fluvial coarse deposits, intermediate siliceous rocks, pure

## General parameters

Area	28.55 km <sup>2</sup>
Percentage of total forest mapped area	0.59 %

## Physics - mean values of profiles (3)

Depth [cm]	Coarse fraction [%]	PAWC [dm <sup>3</sup> /m <sup>2</sup> ]
0-15	20 ± 10	132 ± 57
15-30	40 ± 5	
30-60	35 ± 10	
60-100	30 ± 20	


## Chemistry - mean stocks of profiles (0)

Ctot	Ntot	Ca	Mg	K	P
t/ha	t/ha	kg/ha	kg/ha	kg/ha	kg/ha

All stock values, 0-80 cm including humus layers (F, H), are short-term available, except for phosphorus, which gives long term availability

## Chemistry - mean values of profiles (6)

Depth [cm]	CEC [mmol/kg]	Base Saturation [%]	(Mg+Ca)/CEC	Ntot [%]	TOC [%]	C/N	pH <sub>CaCl2</sub>
0-5	154.62	56.06	0.54	0.3	5.88	19.6	4.23
5-10	154.62	56.06	0.54	0.3	5.88	19.6	4.23
10-20	56.17	55.2	0.53	0.13	2.16	16.62	4.3
20-40	51.09	59.65	0.57	0.09	1.41	15.67	4.53
40-80	56.91	60.54	0.58	0.08	0.9	11.25	4.65



Depth graph of median chemical properties. Shaded area: 25-75% percentiles; CEC: cation exchange capacity (mmol/kg); N: nitrogen (%); TOC: total organic carbon (%); pH<sub>CaCl2</sub>: ph value in CaCl<sub>2</sub> solution; BS: base saturation (%); right-hand y-axis= percentage of profiles used in the calculation

## Biomass use

Effects of whole-tree harvesting



Intermediate negative effects

## Compaction risk

Effects of the transit of heavy machinery



Occasionally critical