

SxS0

solid bedrock, felsic siliceous rocks, impure

General parameters

Area	132.56 km ²
Percentage of total forest mapped area	2.73 %

Physics - mean values of profiles (42)

Depth [cm]	Coarse fraction [%]	PAWC [dm ³ /m ²]
0-15	25 ± 25	76 ± 45
15-30	40 ± 30	
30-60	50 ± 30	
60-100	60 ± 20	



Chemistry - mean stocks of profiles (3)

Ctot	Ntot	Ca	Mg	K	P
t/ha	t/ha	kg/ha	kg/ha	kg/ha	kg/ha
170.98	7.94	312.24	115.37	151.86	922.73

All stock values, 0-80 cm including humus layers (F, H), are short-term available, except for phosphorus, which gives long term availability

Chemistry - mean values of profiles (9)


Depth [cm]	CEC [mmol/kg]	Base Saturation [%]	(Mg+Ca)/CEC	Ntot [%]	TOC [%]	C/N	pH _{CaCl2}
0-5	144.51	12.33	0.11	0.53	10.14	19.13	3.2
5-10	132.95	9.86	0.08	0.38	6.61	17.39	3.33
10-20	100.11	5.16	0.04	0.2	4.03	20.15	3.71
20-40	57.24	6.7	0.05	0.15	2.82	18.8	4.04
40-80	45.18	12.35	0.1	0.24	2.18	9.08	4.23



Depth graph of median chemical properties. Shaded area: 25-75% percentiles; CEC: cation exchange capacity (mmol/kg); N: nitrogen (%); TOC: total organic carbon (%); pH_{CaCl2}: ph value in CaCl₂ solution; BS: base saturation (%); right-hand y-axis= percentage of profiles used in the calculation

Biomass use


Effects of whole-tree harvesting



Strong negative effects

Compaction risk

Effects of the transit of heavy machinery



Minor negative effects