

## General parameters

Area	9.87 km <sup>2</sup>
Percentage on total forest mapped area	0.2 %

## Physics - mean values of all considered profiles (11)

Depth [cm]	Coarse fraction [%]	Field capacity [l/m <sup>2</sup> ]
0-15	30 ± 20	102 ± 37
15-30	45 ± 15	
30-60	50 ± 15	
60-100	65 ± 20	

## Chemistry - stock of available profiles (1)

Ctot	Ntot	Ca	Mg	K	P
t/ha	t/ha	kg/ha	kg/ha	kg/ha	kg/ha
66.38	4.99	4867.52	1132.83	128.19	2005.12

All stock values, 0-80 cm including humus layers (F,H), are short term available, except for phosphorus, which has long term availability

## Chemistry - mean values of all considered profiles (1)

Depth [cm]	CEC [mmol/kg]	Base Saturation [%]	(Mg+Ca)/CEC	Ntot [%]	TOC [%]	C/N	pH <sub>CaCl2</sub>
0-5	87.39	91.35	0.89	0.24	4.69	19.54	4.49
5-10	83.82	98.58	0.97	0.13	1.84	14.15	5.55
10-20	86.56	99.58	0.98	0.12	1.37	11.42	6.55
20-40	73.86	99.59	0.98	0.08	0.56	7	6.94
40-80	42.28	99.54	0.98	0.07	1.16	16.57	7.26





Figure 1: Profile's depth variation of the following median chemical properties, bounded by 25th and 75th percentiles: cation exchange capacity (mmol/kg), nitrogen (%), total organic carbon (%), pH and base saturation (%). The percentage values indicate how many profiles contribute to the median calculation at each depth step.

## Biomass use

Effects of whole-tree harvesting



Minor negative effects

## Compaction risk

Effects of the transit of heavy-duty machinery



Minor negative effects