

General parameters

| | |
|--|-----------------------|
| Area | 79.96 km ² |
| Percentage on total forest mapped area | 1.65 % |

Physics - mean values of all considered profiles (18)

| Depth [cm] | Coarse fraction [%] | Field capacity [l/m ²] |
|------------|---------------------|------------------------------------|
| 0-15 | 15 ± 15 | 136 ± 53 |
| 15-30 | 25 ± 20 | |
| 30-60 | 30 ± 25 | |
| 60-100 | 35 ± 25 | |

Chemistry - stock of available profiles (0)

| Ctot | Ntot | Ca | Mg | K | P |
|------|------|-------|-------|-------|-------|
| t/ha | t/ha | kg/ha | kg/ha | kg/ha | kg/ha |
| | | | | | |

All stock values, 0-80 cm including humus layers (F,H), are short term available, except for phosphorus, which has long term availability

Chemistry - mean values of all considered profiles (9)

| Depth [cm] | CEC [mmol/kg] | Base Saturation [%] | (Mg+Ca)/CEC | Ntot [%] | TOC [%] | C/N | pH _{CaCl2} |
|------------|---------------|---------------------|-------------|----------|---------|-------|---------------------|
| 0-5 | 174.42 | 40.04 | 0.39 | 0.46 | 6.46 | 14.04 | 3.91 |
| 5-10 | 174.42 | 40.04 | 0.39 | 0.46 | 6.46 | 14.04 | 3.91 |
| 10-20 | 72.52 | 17.23 | 0.16 | 0.17 | 2.79 | 16.41 | 3.96 |
| 20-40 | 63 | 36.68 | 0.34 | 0.17 | 2.3 | 13.53 | 4.55 |
| 40-80 | 83.8 | 39.09 | 0.37 | 0.22 | 2.71 | 12.32 | 4.62 |





Figure 1: Profile's depth variation of the following median chemical properties, bounded by 25th and 75th percentiles: cation exchange capacity (mmol/kg), nitrogen (%), total organic carbon (%), pH and base saturation (%). The percentage values indicate how many profiles contribute to the median calculation at each depth step.

Biomass use

Effects of whole tree harvesting



Intermediate negative effects

Compaction risk

Effects of heavy machines transit on the soil



Occasionally critical