

Supplementary Data

lifelines-hc: a Python package for Higher Criticism testing in two-sample survival analysis under non-proportional hazards

Alon Kipnis, Ben Galili, Zohar Yakhini

This document accompanies the Application Note *lifelines-hc: a Python package for Higher Criticism testing in two-sample survival analysis under non-proportional hazards*. It provides (i) the full analysis workflow and scripts used to reproduce the case studies, and (ii) complete per-method test statistics and p -values for all four datasets analysed in the main text.

1 Analysis scripts and data

All scripts live in the `AppNote/` directory of the `lifelines-hc` repository (<https://github.com/TODO/lifelines-hc/tree/main/AppNote>). They require Python ≥ 3.10 , `lifelines-hc`, and `lifelines` ≥ 0.29 .

Workflow

1. **Download trial IPD.** `01_get_kmdata.py` exports reconstructed individual patient data (IPD) from the `kmdata` R package [Guyot et al., 2012] for CheckMate 057 [Borghaei et al., 2015], COMET-1 [Smith et al., 2016], and AZURE [Coleman et al., 2011, 2014]. CSL liver-cirrhosis IPD are downloaded directly from the CRAN `timereg` package and the `SurvSet` repository [Martinussen and Scheike, 2006, Drysdale, 2022] (see `run_csl.py`).
2. **Run per-dataset analyses.** Each `run_*.py` script loads one dataset, calls `run_all_tests()` in `utils.py`, writes a results CSV to `results/`, and saves diagnostic figures to `figs/`.
3. **Build the composite figure.** `make_figure.py` assembles Figure 1 of the main text from the four domain scripts.

Script index

Reproduction commands

```
cd AppNote
python 01_get_kmdata.py --trials Checkmate057_1C COMET1_2A AZURE_2A
python run_immuno_oncology.py
python run_comet.py
python run_azure.py
python run_csl.py           # requires SurvSet or AppNote/data/CSL.csv
python make_figure.py
```

Each `run_*.py` script writes `results/<domain>_test_results.csv` containing one row per competing method (statistic and permutation- or asymptotic-based p -value).

Table 1: Analysis scripts for the four case studies.

Script	Dataset	Role
<code>01_get_kmdata.py</code>	CheckMate 057, COMET-1, AZURE	Download IPD via <code>kmdata</code> (Guyot algorithm).
<code>run_immuno_oncology.py</code>	CheckMate 057 PFS ($n = 582$)	Nivolumab vs. docetaxel; crossing curves.
<code>run_comet.py</code>	COMET-1 OS ($n = 1028$)	Cabozantinib vs. prednisone; transient early benefit.
<code>run_azure.py</code>	AZURE DFS ($n = 3359$)	Zoledronic acid vs. control; menopause-dependent effect.
<code>run_csl.py</code>	CSL OS ($n = 446$)	Prednisone vs. placebo; public raw IPD, multi-window effect.
<code>utils.py</code>	—	Shared <code>run_all_tests()</code> , plotting, and table utilities.
<code>make_figure.py</code>	All four	Composite Figure 1 panels (A–H).

2 Data sources

Table 2 lists the original clinical studies, endpoints, and how each dataset was obtained. For CheckMate 057, COMET-1, and AZURE, only published Kaplan–Meier curves were available; IPD were reverse-engineered with the algorithm of Guyot et al. [2012] and distributed in the `kmdata` package (<https://github.com/raredd/kmdata>). The CSL trial is the only case study with genuinely public raw IPD.

Table 2: Original studies and data acquisition for the four case studies.

Trial	Endpoint	n	Source and acquisition
CheckMate 057	PFS	582	Borghaei et al. [2015]; IPD reconstructed from published Kaplan–Meier curves via Guyot et al. [2012] (<code>kmdata</code>).
COMET-1	OS	1028	Smith et al. [2016]; IPD reconstructed from published Kaplan–Meier curves via Guyot et al. [2012] (<code>kmdata</code>).
AZURE	DFS	3359	Coleman et al. [2011, 2014]; IPD reconstructed from published Kaplan–Meier curves via Guyot et al. [2012] (<code>kmdata</code>).
CSL	OS	446	Copenhagen liver-cirrhosis trial (1962–1969); raw IPD from the CRAN <code>timereg</code> package [Martinussen and Scheike, 2006] and <code>SurvSet</code> [Drysdale, 2022].

Reconstructed IPD files are written to `AppNote/data/` as `Checkmate057_1C.csv`, `COMET1_2A.csv`, and `AZURE_2A.csv` by `01_get_kmdata.py`. CSL data are written to `AppNote/data/CSL.csv` by `run_csl.py`.

3 Shared analysis settings

Unless noted otherwise, all HC-based tests (HCHG, Fisher combination, MinP) use:

- equal-width time pooling with `n_intervals_to_pool` as listed per dataset below;
- two-sided interval hypergeometric p -values (`alternative='both'`);
- HC fraction $\gamma = 0.2$;
- label-permutation p -values with the random seed fixed at 42.

The log-rank test and four weighted log-rank variants (Gehan-Wilcoxon, Tarone-Ware, Peto-Prentice, Fleming-Harrington with $p = q = 1$) use the standard `lifelines.statistics.logrank_test` implementation with asymptotic p -values.

4 Per-method results tables

Tables 3–6 report the full comparison for each dataset. Asterisks mark $p < 0.05$. Permutation-based p -values for HCHG, Fisher, and MinP are discrete (multiples of $1/(n_{\text{perms}} + 1)$ with $n_{\text{perms}} = 2000$) and are rounded to three decimals; “ < 0.001 ” denotes the smallest attainable value, $1/2001$.

CheckMate 057 — progression-free survival ($n = 582$)

Original trial: [Borghaei et al. \[2015\]](#). IPD reconstructed from published Kaplan–Meier curves [[Guyot et al., 2012](#)]. Control: docetaxel ($n = 290$); treatment: nivolumab ($n = 292$). HC settings: $K = 60$ intervals, $n_{\text{perms}} = 2000$.

Table 3: CheckMate 057 PFS: all competing methods.

Method	Statistic	p -value
Log-rank	0.868	0.351
Gehan-Wilcoxon	4.169	0.041*
Tarone-Ware	0.844	0.358
Peto-Prentice	3.869	0.049*
Fleming-Harrington (1,1)	1.292	0.256
Higher Criticism (HC)	1.695	0.001*
Fisher combination	112.561	< 0.001 *
MinP (Bonferroni)	7.112	0.007*

COMET-1 — overall survival ($n = 1028$)

Original trial: [Smith et al. \[2016\]](#). IPD reconstructed from published Kaplan–Meier curves [[Guyot et al., 2012](#)]. Control: prednisone ($n = 346$); treatment: cabozantinib ($n = 682$). HC settings: $K = 80$ intervals, $n_{\text{perms}} = 2000$.

AZURE — disease-free survival ($n = 3359$)

Original trial: [Coleman et al. \[2011, 2014\]](#). IPD reconstructed from published Kaplan–Meier curves [[Guyot et al., 2012](#)]. Control: standard adjuvant therapy ($n = 1678$); treatment: zoledronic acid ($n = 1681$). HC settings: $K = 80$ intervals, $n_{\text{perms}} = 2000$.

Table 4: COMET-1 OS: all competing methods.

Method	Statistic	p -value
Log-rank	1.259	0.262
Gehan-Wilcoxon	1.693	0.193
Tarone-Ware	1.570	0.210
Peto-Prentice	1.602	0.206
Fleming-Harrington (1,1)	0.648	0.421
Higher Criticism (HC)	1.179	0.014*
Fisher combination	128.932	0.028*
MinP (Bonferroni)	4.858	0.242

Table 5: AZURE DFS: all competing methods.

Method	Statistic	p -value
Log-rank	1.054	0.305
Gehan-Wilcoxon	0.652	0.419
Tarone-Ware	0.691	0.406
Peto-Prentice	1.145	0.284
Fleming-Harrington (1,1)	0.277	0.598
Higher Criticism (HC)	1.513	0.005*
Fisher combination	157.060	0.004*
MinP (Bonferroni)	9.168	0.002*

CSL — overall survival ($n = 446$)

Original trial: Copenhagen Study group for Liver diseases (1962–1969); public IPD distributed via [Martinussen and Scheike \[2006\]](#) and [Drysdale \[2022\]](#). Control: placebo ($n = 220$); treatment: prednisone ($n = 226$). Follow-up times are in years. HC settings: $K = 50$ intervals, $n_{\text{perms}} = 2000$.

Table 6: CSL liver-cirrhosis OS: all competing methods.

Method	Statistic	p -value
Log-rank	0.757	0.384
Gehan-Wilcoxon	0.439	0.507
Tarone-Ware	0.774	0.379
Peto-Prentice	0.574	0.449
Fleming-Harrington (1,1)	2.234	0.135
Higher Criticism (HC)	1.257	0.002*
Fisher combination	77.413	0.034*
MinP (Bonferroni)	5.291	0.057

5 Cross-dataset summary

Table 7 condenses the log-rank and HCHG p -values reported in the main text. Full statistics for all eight methods appear in Section 4.

Table 7: Summary of log-rank and HCHG p -values across the four case studies (full tables in Section 4).

Dataset	Endpoint	n	Log-rank p	HCHG p
CheckMate 057	PFS	582	0.351	0.001
COMET-1	OS	1028	0.262	0.014
AZURE	DFS	3359	0.305	0.005
CSL	OS	446	0.384	0.002

Across all four datasets, HCHG achieves $p \leq 0.014$ while the log-rank test is non-significant throughout ($p \geq 0.26$). Among the weighted log-rank alternatives, only CheckMate 057 Gehan-Wilcoxon and Peto-Prentice reach $p < 0.05$ (at 0.041 and 0.049 respectively); every other weighted test returns $p \geq 0.13$.

6 Machine-readable results

The CSV files below contain the same values as Tables 3–6 (columns `method`, `statistic`, `p_value`):

- AppNote/results/immuno_test_results.csv
- AppNote/results/comet_test_results.csv
- AppNote/results/azure_test_results.csv
- AppNote/results/csl_test_results.csv

References

- Hossein Borghaei, Luis Paz-Ares, Leora Horn, David R. Spigel, Martin Steins, Neal E. Ready, Laura Q. Chow, Everett E. Vokes, Enriqueta Felip, Esther Holgado, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *New England Journal of Medicine*, 373(17):1627–1639, 2015. doi: 10.1056/NEJMoa1507643.
- Robert E. Coleman, Howard Marshall, David Cameron, David Dodwell, Robert Burkinshaw, Mary Keane, Michael Gil, Stephen J. Houston, Robert J. Grieve, Peter J. Barrett-Lee, et al. Breast-cancer adjuvant therapy with zoledronic acid. *New England Journal of Medicine*, 365(15): 1396–1405, 2011. doi: 10.1056/NEJMoa1105195.
- Robert E. Coleman, Marie Collinson, Eleanor Banks, Graham Clack, Dorothy Ritchie, Victoria Liversedge, David Cameron, David Dodwell, Michael Jones, Peter Barrett-Lee, et al. Benefits and risks of adjuvant treatment with zoledronic acid in stage II/III breast cancer. 10 year follow-up of the AZURE randomized clinical trial. *Journal of Bone Oncology*, 3(2):23–30, 2014. doi: 10.1016/j.jbo.2013.12.001.
- Erik Drysdale. SurvSet: An open-source time-to-event dataset repository. *arXiv preprint arXiv:2203.03094*, 2022. doi: 10.48550/arXiv.2203.03094.
- Patricia Guyot, A. E. Ades, Mario J. N. M. Ouwers, and Nicky J. Welton. Enhanced secondary analysis of survival data: reconstructing the data from published Kaplan-Meier survival curves. *BMC Medical Research Methodology*, 12:9, 2012. doi: 10.1186/1471-2288-12-9.

Torben Martinussen and Thomas H. Scheike. *Dynamic Regression Models for Survival Data*. Springer, New York, 2006. doi: 10.1007/0-387-33960-4. CSL liver cirrhosis trial (prednisone vs placebo, Copenhagen Study group for Liver diseases, 1962–1969); dataset `cs1` in the CRAN `timereg` package.

Matthew R. Smith, Christopher J. Sweeney, Paul G. Corn, Dana E. Rathkopf, David C. Smith, Maha Hussain, Daniel J. George, Celestia S. Higano, Andrea L. Harzstark, A. Oliver Sartor, et al. Cabozantinib in chemotherapy-pretreated metastatic castration-resistant prostate cancer: results of a phase III randomized controlled trial (COMET-1). *Journal of Clinical Oncology*, 34(25): 3005–3013, 2016. doi: 10.1200/JCO.2015.65.5597.