

Numerov Method for Bound States in the One-Dimensional Schrödinger Equation - Project Outline and Motivation -

Alon Sportes (ID: 204498745)

Contents

Table of Contents	1
List of Figures	3
1 What problem are we solving?	4
2 Which course topics are used?	4
2.1 Lecture 1: Numerical errors and computational project expectations	4
2.2 Lecture 3: Numerical differentiation and integration	4
2.3 Lecture 4: Root finding	5
2.4 Lecture 5: ODEs, boundary-value problems, eigenvalue problems, and shoot- ing	5
2.5 Lecture 6: PDEs and Schrödinger equation context	6
2.6 What is not used	6
3 Why Numerov?	6
4 Why shooting?	7
5 Why parity?	8
6 Why root finding and bisection?	9
7 What each code file does	10
7.1 src/potentials.py	10
7.2 src/numerov.py	10
7.3 src/shooting.py	11
7.4 src/analysis.py	11
7.5 src/rk4_compare.py	12
7.6 src/scattering.py	12
7.7 src/plotting.py	13
7.8 src/experiments.py	13
7.9 scripts/run_solver.py	14
7.10 tests/test_solver.py	14
8 What the project demonstrates	14
8.1 Level 1: Exact benchmarks	15
8.2 Level 2: Convergence	15
8.3 Level 3: New physics	15

9	What is apparent from the lectures and what is beyond them?	16
9.1	Clearly from the lectures	16
9.2	Partly from the lectures, but specialized to quantum mechanics	16
9.3	Beyond the slides	16
9.4	Not used from the lectures	17
10	The clean explanation for a reviewer	17

List of Figures

1 What problem are we solving?

The project solves the one-dimensional time-independent Schrödinger equation. In the dimensionless convention used in the code, it is written as

$$\psi''(x) = 2[V(x) - E]\psi(x). \quad (1)$$

The unknowns are:

- $\psi(x)$: the wavefunction,
- E : the energy eigenvalue,
- $V(x)$: the chosen potential.

This is not just an ordinary differential equation problem. It is a boundary-value eigenvalue problem. Only special values of E give physically valid bound states. This matches the proposal and the report structure exactly.

2 Which course topics are used?

2.1 Lecture 1: Numerical errors and computational project expectations

This project uses the main idea from Lecture 1: numerical computation is approximate, so numerical errors must be controlled and the method must be documented. The final project instructions also require code documentation and a short writeup explaining the problem, the method, the results, and the convergence and correctness checks.

In this project, this appears through:

- convergence studies,
- comparison with exact solutions,
- normalization checks,
- automated tests,
- README documentation,
- discussion of truncation error, round-off error, and numerical floors.

A very important example is the infinite square well convergence study. The solver originally produced very accurate energies, but the observed convergence order was not ideal. The issue was not the Numerov recurrence itself, but the startup values used to seed the two-step recurrence. After adding higher Taylor terms to the parity-based startup conditions, the square-well convergence slopes improved to approximately fourth order. A similar lesson appeared later in the quartic double-well study: the convergence behavior was polluted until the potential offset, startup expansion, and convergence methodology were made consistent. These are exactly the kinds of numerical-error diagnoses emphasized in the course.

2.2 Lecture 3: Numerical differentiation and integration

Numerical integration is used when normalizing the wavefunction:

$$\|\psi\| = \sqrt{\int |\psi(x)|^2 dx}. \quad (2)$$

In the code, this is done with `np.trapezoid`. The reason is physical: quantum wavefunctions must satisfy

$$\int |\psi(x)|^2 dx = 1. \quad (3)$$

Therefore, normalization is not optional. It makes the solution physically meaningful.

Numerical differentiation is also used in `derivative_at_right_edge()`. This is especially important for inward shooting. For even states, the condition at the origin is

$$\psi'(0) = 0. \quad (4)$$

If the derivative is computed with a low-order stencil, then the whole eigenvalue calculation can lose accuracy even if the Numerov recurrence itself is high order. This happened in the harmonic oscillator case. The code now uses a fourth-order backward stencil when enough points are available. This keeps the boundary-condition accuracy consistent with the rest of the method.

2.3 Lecture 4: Root finding

Root finding is used to determine the allowed energies. The project uses bisection: first detect a sign change, then repeatedly cut the interval in half.

In the code this appears in:

- `find_brackets()`,
- `bisect_energy()`,
- `find_inward_decay_brackets()`,
- `bisect_energy_inward_decay()`,
- `find_rk4_brackets()`,
- `bisect_rk4_energy()`.

Bisection is used because it is stable and easy to justify. Newton's method is faster, but it requires derivatives and can fail if the starting guess is poor. Bisection is slower, but it is robust. In the latest solver, the bisection routine was also adjusted so it does not stop too early only because the energy bracket became small; it now uses a safeguarded secant polishing step to improve the final root estimate.

The project also includes root-finding diagnostics. These are not strictly required, but they are useful because they show how the mismatch function crosses zero and how the bisection process isolates eigenvalues.

2.4 Lecture 5: ODEs, boundary-value problems, eigenvalue problems, and shooting

This is the core lecture for the project. The Schrödinger equation is a second-order ODE, but unlike an initial-value problem, the correct energy is not known in advance. Therefore, the method is:

1. guess an energy E ,
2. integrate the ODE,

3. check the boundary condition,
4. improve the energy guess.

That is the shooting method.

The project uses two shooting strategies:

- outward shooting from $x = 0$,
- inward shooting from $x_{\max}$ to $x = 0$.

Outward shooting is used for many symmetric finite-domain or effectively finite-domain problems. Inward shooting is used mainly for the harmonic oscillator. This is because the harmonic oscillator is defined on an infinite domain, and outward integration can pick up the unphysical growing exponential tail. Inward shooting starts from the decaying asymptotic side and imposes the parity condition at the origin.

2.5 Lecture 6: PDEs and Schrödinger equation context

Lecture 6 mentions the Schrödinger equation as one of the important PDEs in physics. This project uses the time-independent Schrödinger equation in one dimension, which becomes an ODE eigenvalue problem. Therefore, this project is not solving a time-dependent PDE. It is using an ODE method on the stationary quantum problem.

2.6 What is not used

The project does not use:

- finite-volume methods,
- upwind methods,
- CFL conditions,
- Riemann solvers,
- shock capturing,
- slope limiters,
- Euler fluid equations,
- MHD,
- Particle in Cell.

Those topics belong to the advection, hydrodynamics, Burgers equation, Euler equations, and PIC parts of the course. They are not part of this project.

3 Why Numerov?

The equation has the form

$$\psi''(x) = q(x)\psi(x), \tag{5}$$

where

$$q(x) = 2[V(x) - E]. \tag{6}$$

This form is exactly what Numerov is designed for: second-order ODEs without a first-derivative term.

In `src/numerov.py`:

- `q_from_energy()` builds $q(x)$,
- `numerov_outward()` performs the Numerov recurrence,
- `normalize_wavefunction()` rescales the final wavefunction so that total probability is 1,
- `derivative_at_right_edge()` estimates the derivative at the boundary when needed.

Numerov is used instead of Euler or a generic Runge-Kutta method because it is especially efficient for equations like the Schrödinger equation. It uses neighboring grid values and gives high accuracy for second-order linear ODEs. This is also consistent with Pang, where the one-dimensional Schrödinger equation is treated using wavefunction integration and eigenvalue search.

A key implementation detail is that Numerov is a two-step method. It needs ψ at the first two grid points. Therefore, the code must provide startup values. In `shooting.py`, `initial_conditions()` gives those startup values using Taylor expansion and parity.

For even states:

$$\psi(0) = 1, \quad \psi'(0) = 0. \quad (7)$$

For odd states:

$$\psi(0) = 0, \quad \psi'(0) = 1. \quad (8)$$

The latest code includes higher Taylor terms and the curvature of the local coefficient $q(x)$ near the origin:

- even states include the h^4 term,
- odd states include the h^5 term.

For the quartic double well, the startup expansion was improved further by including the relevant $q''(0)$ terms for both even and odd parity. This matters because the quartic potential has curvature structure near the origin that affects the first Numerov steps. Without these terms, one of the low-lying double-well states showed a degraded convergence slope. After the correction, the state recovered high-order convergence consistent with the Numerov method.

This matters because if the startup formula is too low order, it can pollute the observed convergence rate. This was seen first in the infinite square well: the energies were accurate, but the convergence slopes were not cleanly fourth order. It appeared again in the quartic double well, where the second excited state initially converged too slowly. After correcting the startup formula, the square-well states show slopes close to 4, and the double-well state that had been stuck near $p \approx 3.09$ improved to about $p \approx 4.55$.

4 Why shooting?

A bound state must behave correctly at both boundaries. However, Numerov integration needs starting values. Therefore, the algorithm is:

1. choose a trial energy E ,

2. choose initial conditions at $x = 0$, or from the decaying tail,
3. integrate with Numerov,
4. check whether the wavefunction satisfies the target boundary condition,
5. adjust E .

This is shooting.

In `src/shooting.py`, the outward shooting logic is built from:

- `initial_conditions()`,
- `half_domain_wavefunction()`,
- `boundary_mismatch()`,
- `find_brackets()`,
- `bisect_energy()`,
- `solve_state_from_bracket()`,
- `solve_symmetric_potential()`.

The important function is `boundary_mismatch()`. It answers the question: for this trial energy, did the wavefunction land correctly at the boundary? If not, the energy is wrong.

The latest solver also changes how the final mismatch is stored. Instead of reporting the raw, unnormalized boundary amplitude $\psi(x_{\max})$, the code now stores the normalized boundary leakage after the full wavefunction has been reconstructed and normalized. This makes the residual scale-invariant and physically more meaningful. In practice, this changed the low-state double-well residuals from raw values around 10^{-6} – 10^{-5} to normalized leakage values around 2×10^{-7} .

The inward shooting logic is built from:

- `inward_decay_initial_conditions()`,
- `inward_decay_half_domain_wavefunction()`,
- `inward_decay_boundary_mismatch()`,
- `find_inward_decay_brackets()`,
- `bisect_energy_inward_decay()`,
- `solve_symmetric_potential_inward_decay()`.

Inward shooting is used for confining potentials such as the harmonic oscillator. The physical solution decays at large x . If we integrate outward from the origin, the numerical solution can pick up the unphysical growing mode. Inward shooting starts in the forbidden region and integrates toward the origin, giving cleaner harmonic oscillator wavefunctions and more reliable parity enforcement.

5 Why parity?

The main bound-state potentials are symmetric:

$$V(-x) = V(x). \tag{9}$$

For symmetric potentials, quantum states are either even or odd.

Even states satisfy

$$\psi(-x) = \psi(x), \quad (10)$$

and odd states satisfy

$$\psi(-x) = -\psi(x). \quad (11)$$

Therefore, instead of integrating from $-x_{\max}$ to $x_{\max}$, the code integrates only from 0 to $x_{\max}$.

In code, `initial_conditions()` uses:

- even: $\psi(0) = 1, \psi'(0) = 0$,
- odd: $\psi(0) = 0, \psi'(0) = 1$.

Then `build_full_wavefunction()` reflects the half-domain solution to the left side.

This choice makes the algorithm simpler, faster, and more stable. It also lets the states be classified physically as even or odd.

Parity is also useful for the double well because tunneling splitting appears as pairs of even and odd states. The lowest two states are symmetric and antisymmetric combinations of wavefunctions localized in the two wells.

6 Why root finding and bisection?

For each trial energy, the boundary mismatch gives a number:

$$F(E) = \text{mismatch}(E). \quad (12)$$

Allowed eigenvalues satisfy

$$F(E) = 0. \quad (13)$$

So the eigenvalue problem becomes a root-finding problem.

The code first scans energies using `find_brackets()`. It looks for sign changes. If $F(E_1)$ and $F(E_2)$ have opposite signs, then a root is likely between them. Then `bisect_energy()` refines the energy until the interval is small enough.

This is exactly the root-finding logic from the course: bracket a root, then reduce the bracket until the required accuracy is reached.

The code also includes:

- `sample_boundary_mismatch()`,
- `bisection_history()`,
- `sample_inward_decay_mismatch()`,
- `bisection_history_inward_decay()`.

These are mainly for diagnostics and plots. They show that the eigenvalues are not magic numbers. They are roots of a clearly defined numerical function.

7 What each code file does

7.1 `src/potentials.py`

This file defines the physical systems:

- `harmonic_oscillator()`,
- `infinite_square_well_numeric()`,
- `finite_square_well()`,
- `quartic_double_well()`,
- `square_barrier()`,
- `double_square_barrier()`.

The purpose is to separate the physics definitions from the numerical solver. This is good design: the solver does not care which potential is chosen. It only needs $V(x)$. That makes the project modular.

The main potentials have clear roles:

- infinite square well: analytic validation,
- harmonic oscillator: analytic validation and inward shooting,
- finite square well: nontrivial finite bound-state system,
- quartic double well: tunneling and energy splitting, with the shifted potential using the analytic minimum $V_{\min} = -b^2/(4a)$,
- single barrier and double barrier: scattering extension.

A recent correction was made in `quartic_double_well()`: the zero shift now uses the analytic minimum

$$V_{\min} = -\frac{b^2}{4a} \quad (14)$$

instead of the sampled grid minimum. This is important because using the sampled minimum makes the potential itself depend slightly on the grid spacing h , which contaminates a grid-convergence study.

7.2 `src/numerov.py`

This is the numerical integration engine. Its main role is to take an x grid, a potential V , and a trial energy E , and produce a trial wavefunction ψ .

The most important functions are:

- `q_from_energy()`,
- `numerov_outward()`,
- `normalize_wavefunction()`,
- `derivative_at_right_edge()`.

Important details:

- `numerov_outward()` includes dynamic rescaling to avoid overflow when a non-eigenvalue trial solution grows exponentially.

- `normalize_wavefunction()` rescales before squaring to avoid overflow.
- `derivative_at_right_edge()` uses a fourth-order stencil when possible.

7.3 `src/shooting.py`

This is the eigenvalue solver. Its main role is to try energies, integrate with Numerov, check the boundary mismatch, find the correct eigenvalues, and return normalized eigenstates.

The most important functions are:

- `initial_conditions()`,
- `find_brackets()`,
- `bisect_energy()`,
- `solve_symmetric_potential()`,
- `solve_symmetric_potential_inward_decay()`.

Important latest details: `initial_conditions()` now includes higher Taylor terms so the first Numerov step does not reduce the convergence order. For the quartic double well, the startup was also extended with the $q''(0)$ contribution. This fixed the square-well convergence slope problem and restored the expected high-order convergence for the problematic double-well state. The bisection routine was also strengthened with a safeguarded secant polishing step, and `StateSolution.mismatch` now stores normalized boundary leakage rather than raw unnormalized wall amplitude.

7.4 `src/analysis.py`

This is the scientific validation layer. It contains:

- `exact_square_well_energies()`,
- `exact_harmonic_oscillator_energies()`,
- `relative_error()`,
- `estimate_convergence_slopes()`,
- `convergence_vs_grid()`,
- `convergence_vs_grid_successive()`,
- `convergence_vs_box_size()`,
- `convergence_vs_box_size_fixed_spacing()`,
- `splitting_vs_parameter()`.

This is what turns the code from “it runs” into “it was verified scientifically.”

For cases without analytic solutions, such as the quartic double well, the code now distinguishes between two convergence strategies. The box-size study is compared against a larger-box reference, while the grid-spacing study uses successive refinements, comparing each grid to the next finer one. This avoids pretending that there is an exact spectrum and avoids introducing an artificial reference floor into the h -convergence study.

7.5 `src/rk4_compare.py`

This is the method-comparison module. It solves the harmonic oscillator using RK4 shooting.

The harmonic oscillator is used because it has exact energies and a smooth potential, so it is a fair comparison.

RK4 works differently from Numerov. Numerov solves the second-order equation directly. RK4 rewrites it as a first-order system:

$$\psi' = \phi, \tag{15}$$

$$\phi' = 2[V(x) - E]\psi. \tag{16}$$

This means RK4 evolves both ψ and ψ' explicitly.

The comparison is useful because it shows the difference between a specialized solver and a general-purpose ODE solver. It also explains why the derivative boundary condition mattered so much. RK4 carries ψ' as a variable, while Numerov needs a finite-difference derivative estimate when enforcing some boundary conditions.

7.6 `src/scattering.py`

This is the scattering extension. It computes:

- transmission T ,
- reflection R ,
- conservation check $T + R$.

The method is:

1. assume the potential is zero far left and far right,
2. impose a transmitted right-moving wave on the far right,
3. integrate backward through the potential with complex Numerov,
4. decompose the left-side wave into incident and reflected parts.

The main functions are:

- `numerov_outward_complex()`,
- `integrate_from_right()`,
- `decompose_left_asymptotic()`,
- `solve_scattering()`,
- `sweep_scattering()`,
- `find_transmission_peaks()`.

This is not the core bound-state project, but it shows that the same Numerov framework can be extended to scattering and resonant tunneling.

7.7 src/plotting.py

This file generates report figures:

- `plot_potential_and_states()`,
- `plot_probability_densities()`,
- `plot_energy_comparison()`,
- `plot_error_curve()`,
- `plot_splitting_curve()`,
- `plot_root_finding_diagnostic()`,
- `plot_scattering_coefficients()`,
- `plot_scattering_potential_and_probability()`,
- `plot_numerov_vs_rk4_errors()`.

These plots let the reviewer visually see the states, convergence, tunneling behavior, scattering behavior, and method comparison. Recent plotting updates increased the displayed energy precision from 4 to 6 decimals and changed the root-diagnostic plots to show the raw mismatch on a symmetric logarithmic scale. This makes even and odd root scans easier to distinguish and avoids hiding structure through clipping or overly aggressive scaling.

7.8 src/experiments.py

This file connects the solver to the actual project cases. It runs:

- `run_square_well()`,
- `run_harmonic_oscillator()`,
- `run_double_well()`,
- `run_finite_square_well()`,
- `run_scattering()`.

This file is basically the scientific experiment plan in code.

Important latest double-well additions:

- the default double-well domain was increased from $x_{\max} = 3.0$ to $x_{\max} = 4.0$,
- grid-convergence and box-convergence studies are now separated properly,
- the grid-spacing study now uses successive refinements rather than a single fixed reference,
- the box-size study is still compared against a larger-box reference,
- the double-well plot legend now shows the explicit formula,
- root-finding diagnostics are now generated separately for the double well,
- redundant outputs such as the old `3_double_well_splitting.png` were removed,
- the retained splitting plot is `3_double_well_Numerov_splitting_vs_b.png`,
- each experiment now writes into its own subdirectory under `results/`, for example `results/3_double_well_Numerov/`.

7.9 scripts/run_solver.py

This is the entry point. It:

- clears old results,
- runs all experiments,
- runs tests,
- prints a completion message.

It is intentionally lightweight. Most logic lives in `src/`. The latest runner keeps a single canonical results root at `PROJECT_ROOT/results` and each experiment writes its CSV and PNG outputs into its own dedicated subdirectory there.

7.10 tests/test_solver.py

This file checks correctness. It tests:

- normalization,
- derivative stencil accuracy,
- RK4 bracket detection,
- square-well ground-state accuracy,
- square-well convergence order,
- harmonic oscillator energies,
- harmonic oscillator inward shooting,
- analytic quartic double-well shift,
- larger-box improvement for the double well,
- same-box successive refinement error decrease,
- low-state double-well convergence order,
- parity-specific mismatch scans,
- double-well splitting,
- scattering probability conservation,
- `run_solver` import behavior.

Important latest regression test: `test_square_well_convergence_order()` requires the first square-well states to show near-fourth-order convergence. This prevents the startup-order bug from quietly returning.

8 What the project demonstrates

The project demonstrates three levels of correctness.

8.1 Level 1: Exact benchmarks

The square well and harmonic oscillator have known exact energies. If the numerical method reproduces them, the solver is credible.

Square well:

- validates the basic outward Numerov shooting method,
- checks node structure and parity,
- checks fourth-order convergence after the improved startup values.

Harmonic oscillator:

- validates inward shooting,
- checks parity and node structure,
- checks domain truncation effects,
- supports the Numerov versus RK4 comparison.

8.2 Level 2: Convergence

The project varies:

- grid spacing h ,
- domain size $x_{\max}$.

The goal is to show that the energies stabilize. This addresses truncation error, resolution dependence, finite-domain effects, floating-point limits, and root-finding limits. For the quartic double well, the convergence studies now deliberately separate grid refinement from box-size effects: h -convergence uses successive refinements at fixed domain size, while $x_{\max}$ -convergence compares against a larger-domain reference.

A key interpretation point is that if a convergence slope is slightly above 4, this does not mean the method is truly higher than fourth order. It usually means the errors are very small and the log-log fit is being affected by numerical floors, root-finding tolerance, or floating-point limits.

8.3 Level 3: New physics

The double well has nearly degenerate even and odd pairs. This shows tunneling splitting, which is the strongest physics result in the project.

The double-well analysis now includes:

- states and densities,
- energy splitting versus parameter b ,
- root-finding diagnostics for even and odd parity,
- convergence versus h using successive refinements,
- convergence versus $x_{\max}$ using a larger-box reference,
- an analytic zero shift based on $V_{\min} = -b^2/(4a)$,
- normalized boundary leakage as the residual diagnostic.

9 What is apparent from the lectures and what is beyond them?

9.1 Clearly from the lectures

- ODEs,
- boundary-value problems,
- eigenvalue problems,
- shooting,
- root finding,
- bisection,
- numerical integration,
- numerical differentiation,
- convergence,
- numerical errors,
- documentation.

9.2 Partly from the lectures, but specialized to quantum mechanics

- Numerov method,
- wavefunction normalization,
- parity classification,
- bound-state interpretation,
- inward shooting from a decaying tail,
- tunneling splitting,
- scattering transmission and reflection.

The Numerov idea appears in Pang's Schrödinger equation treatment, but it is more specialized than the main lecture slides.

9.3 Beyond the slides

The most advanced or specialized parts are:

- fourth-order-consistent Numerov startup values,
- startup expansion terms involving $q''(0)$ for the quartic double well,
- high-order derivative stencil at the boundary,
- safeguarded secant polishing after bisection,
- scale-invariant normalized boundary leakage as a residual,
- inward shooting from the forbidden region,
- successive-refinement convergence for non-analytic potentials,

- high-resolution larger-box reference for domain-size studies,
- RK4 versus Numerov method comparison,
- complex Numerov scattering,
- double-barrier resonant tunneling.

9.4 Not used from the lectures

- advection schemes,
- finite-volume methods,
- limiters,
- Burgers equation,
- Euler equations,
- MHD,
- PIC,
- chaos,
- Monte Carlo,
- FFT,
- matrix eigenvalue solvers.

This is fine. The final project explicitly allows choosing a different physics problem or going beyond class material.

10 The clean explanation for a reviewer

Here is the project summary in reviewer language:

I am solving the one-dimensional time-independent Schrödinger equation as a boundary-value eigenvalue problem. I discretize space on a uniform grid and use the Numerov method to integrate the second-order ODE for a trial energy. Since only special energies satisfy the required boundary conditions, I use a shooting method. The boundary mismatch becomes a scalar function of energy, and I find its roots using bracketing and bisection. For symmetric potentials I use parity to reduce the calculation to the half domain, then reconstruct and normalize the full wavefunction. I validate the solver on the infinite square well and harmonic oscillator, study convergence with grid spacing and domain size, and then apply it to a finite square well and quartic double well to analyze nontrivial bound states and tunneling-induced splitting. For the double well, I separate grid and domain convergence, use an analytic potential shift, and report normalized boundary leakage as the residual diagnostic. I also compare Numerov with RK4 for the harmonic oscillator and include scattering through finite and double barriers as a secondary extension.

That is the full logic of the project.