

Numerov Method for Bound States in the One-Dimensional Schrödinger Equation

Alon Sportes (ID: 204498745)

Contents

Table of Contents	1
List of Figures	2
1 Introduction	3
2 Computational approach	3
2.1 The Schrödinger equation and Numerov method	3
2.2 Shooting and root finding	3
3 Results	4
3.1 Infinite square well	4
3.2 Harmonic oscillator	4
3.2.1 Numerov method on unbounded domains	4
3.2.2 Numerov versus Runge-Kutta	6
3.3 Double-well potential	6
3.4 Finite square well	9
3.5 Scattering and resonant tunneling	9
3.6 Convergence study	9
4 Discussion	10
5 Conclusion	10
References	11

List of Figures

1	Comparison between numerical and exact energy eigenvalues for the infinite square well. The agreement is excellent for the first four states.	5
2	Probability densities $ \psi(x) ^2$ for the first four eigenstates of the infinite square well. The correct number of nodes and boundary behavior are clearly reproduced.	5
3	Boundary mismatch functions for even (top) and odd (bottom) states. The zero crossings correspond to the physical eigenvalues found by the shooting method.	5
4	Convergence of the energy error as a function of grid spacing h . The slopes are close to fourth order, as expected for the Numerov method.	7
5	Wavefunctions for the first four states plotted together with their corresponding energies and the confining potential.	7
6	Comparison between numerical and exact energy eigenvalues for the harmonic oscillator. The agreement is excellent for the first four states.	7
7	Probability densities $ \psi(x) ^2$ for the first four harmonic oscillator eigenstates. The correct node structure is clearly reproduced.	7
8	Boundary mismatch functions for even (top) and odd (bottom) states. The zero crossings correspond to the physical eigenvalues.	8
9	Convergence of the energy error as a function of grid spacing h . The slopes are close to fourth order, as expected for the Numerov method.	8
10	Dependence of the energy error on the domain size $x_{\max}$. Errors decrease rapidly before reaching a plateau set by discretization and floating-point limits.	8
11	Wavefunctions for the first four harmonic oscillator states plotted together with the potential.	12
12	Comparison of Numerov and RK4 shooting for the harmonic oscillator. Both methods show fourth-order convergence, while Numerov gives a smaller maximum energy error at the same grid spacing.	12

1 Introduction

This project studies the one-dimensional time-independent Schrödinger equation using the Numerov method combined with a parity-based shooting procedure. The goals are to validate the method on systems with known analytic solutions, demonstrate numerical convergence and accuracy, and then apply it to more interesting bound-state potentials such as a double well and a finite square well. Extensions to scattering problems are also explored as a secondary component.

2 Computational approach

2.1 The Schrödinger equation and Numerov method

In dimensionless units, the Schrödinger equation is written as

$$\psi''(x) = 2[V(x) - E]\psi(x).$$

The wavefunction is integrated with the Numerov scheme on a uniform grid. For symmetric potentials, even and odd states are treated separately using parity conditions at $x = 0$. Eigenvalues are found by scanning for sign changes in the boundary mismatch and then refining them with bisection. The correctness of the numerical approach is verified through comparison with analytical solutions where available and through systematic convergence studies.

The overall numerical workflow is implemented in a modular way in the accompanying Python code. The core components include a Numerov integrator for propagating the wavefunction, a shooting module that enforces parity-based boundary conditions and constructs the mismatch function, and analysis utilities for convergence studies and parameter sweeps. This separation of concerns allows each numerical component to be validated independently and makes the implementation easier to interpret [1].

2.2 Shooting and root finding

The boundary-value eigenproblem is reformulated as a root-finding problem for a mismatch function $M(E)$. For a given trial energy E , the Schrödinger equation is integrated and a boundary mismatch is evaluated. For symmetric potentials we exploit parity and define

$$\text{even states: } M(E) = \psi'_E(0), \tag{1}$$

$$\text{odd states: } M(E) = \psi_E(0). \tag{2}$$

Eigenvalues correspond to the roots of $M(E) = 0$, which are located by bracketing sign changes and refined with bisection. To make the procedure transparent, we plot $M(E)$ versus E and mark the bisection iterates; zero crossings identify the allowed energies.

In the implementation, the mismatch function is evaluated numerically using the propagated wavefunction. For even states, the derivative at the origin is computed using a high-order finite-difference stencil to ensure that the overall accuracy of the method is not limited by the boundary evaluation. This detail is crucial for achieving consistent convergence behavior in the harmonic oscillator case.

3 Results

Each potential is used to validate, test, or demonstrate different aspects of the numerical method, ranging from analytic benchmarks to nontrivial quantum behavior.

3.1 Infinite square well

The infinite square well serves as a benchmark problem with known analytic spectrum

$$E_n = \frac{(n+1)^2 \pi^2}{8a^2}.$$

The potential is

$$V(x) = \begin{cases} 0 & |x| \leq a, \\ \infty & |x| > a \end{cases} \quad (\text{implemented numerically with a large finite } V_0).$$

The numerical eigenvalues agree with the analytic values to high precision, as shown in Fig. 1. Convergence with respect to grid spacing is demonstrated with log-log plots (Fig. 4), and the fitted slopes indicate power-law behavior $\Delta E \propto h^p$.

The probability densities for the first four states (Fig. 2) display the correct number of nodes and vanish at the boundaries, consistent with the analytic square-well eigenstates.

The shooting method is validated by plotting the boundary mismatch functions for both even and odd states (Fig. 3). The zero crossings correspond to the physical eigenvalues, confirming that the parity separation is working as intended and that the root finder is selecting the physical eigenvalues rather than spurious numerical solutions.

The convergence of the energy error as a function of grid spacing h (Fig. 4) is close to fourth order, as expected for the Numerov method. Higher excited states are more demanding to resolve.

Finally, the wavefunctions for the first four states are plotted together with their corresponding energies and the confining potential (Fig. 5), and the convergence study shows the expected high-order decrease in error.

3.2 Harmonic oscillator

3.2.1 Numerov method on unbounded domains

For the harmonic oscillator with $V(x) = \frac{1}{2}\omega^2 x^2$, the exact spectrum is

$$E_n = \omega \left(n + \frac{1}{2} \right), \quad n = 0, 1, 2, \dots$$

This provides a second analytic benchmark, but on an unbounded domain. Numerically, the domain is truncated to $[-x_{\max}, x_{\max}]$, and decaying boundary conditions are imposed.

The numerical eigenvalues agree with the exact linear spectrum to high precision, as shown in Fig. 6. This confirms that the inward-shooting formulation correctly handles the exponentially decaying asymptotic behavior.




Figure 1: Comparison between numerical and exact energy eigenvalues for the infinite square well. The agreement is excellent for the first four states.




Figure 2: Probability densities $|\psi(x)|^2$ for the first four eigenstates of the infinite square well. The correct number of nodes and boundary behavior are clearly reproduced.




Figure 3: Boundary mismatch functions for even (top) and odd (bottom) states. The zero crossings correspond to the physical eigenvalues found by the shooting method.

The probability densities (Fig. 7) show the expected structure: the ground state is Gaussian-like with no nodes, while higher states exhibit the correct number of nodes and increasing spatial extent. This confirms both physical correctness and stability of the integration on a truncated domain.

The shooting mismatch functions (Fig. 8) exhibit clear zero crossings at the eigenvalues. Even and odd states are separated using parity conditions at the origin, and the roots alternate as expected. Although the mismatch spans many orders of magnitude due to exponential behavior, the crossings remain well-defined, indicating a robust root-finding procedure.

Convergence with respect to grid spacing h (Fig. 9) shows near fourth-order behavior for most states, consistent with the Numerov method. Slight deviations (e.g. $p > 4$ for the ground state) are attributed to fitting in a regime close to machine precision rather than a true increase in order.

The dependence on domain size (Fig. 10) shows rapid error reduction as $x_{\max}$ increases, followed by a plateau. This reflects the transition from truncation error (small $x_{\max}$) to discretization and floating-point limits (large $x_{\max}$).

Finally, the combined state plot (Fig. 11) shows the wavefunctions together with the confining potential. The spatial extent increases with energy, and the states remain well-behaved across the domain.

3.2.2 Numerov versus Runge–Kutta

As an additional method comparison, the harmonic oscillator was also solved using a fourth-order Runge–Kutta (RK4) shooting method. In this formulation the second-order Schrödinger equation is rewritten as the first-order system

$$\psi' = \phi, \quad \phi' = 2[V(x) - E]\psi.$$

This makes RK4 a useful general-purpose comparison for Numerov, which instead exploits the special second-order structure of the Schrödinger equation directly.

Figure 12 compares the maximum energy error over the computed harmonic-oscillator states as a function of grid spacing. Both methods show approximately fourth-order convergence, as expected. Numerov is consistently more accurate for the same grid spacing, although the difference is moderate. This is consistent with the fact that both methods are already high-order and that the final accuracy is also affected by root-finding tolerance, domain truncation, and boundary-condition treatment.

This comparison also emphasizes an important numerical point: the order of the integration scheme alone is not sufficient. Boundary conditions and diagnostic quantities must be evaluated with comparable accuracy. After using a high-order boundary derivative for the Numerov inward-shooting condition, the Numerov and RK4 comparison behaves as expected.

3.3 Double-well potential

The quartic double-well potential is used to demonstrate nontrivial quantum behavior beyond analytic benchmarks and produces pairs of nearly degenerate states. The lowest even and odd states correspond to symmetric and antisymmetric combinations of localized wavefunctions. The




Figure 4: Convergence of the energy error as a function of grid spacing h . The slopes are close to fourth order, as expected for the Numerov method.




Figure 5: Wavefunctions for the first four states plotted together with their corresponding energies and the confining potential.




Figure 6: Comparison between numerical and exact energy eigenvalues for the harmonic oscillator. The agreement is excellent for the first four states.




Figure 7: Probability densities $|\psi(x)|^2$ for the first four harmonic oscillator eigenstates. The correct node structure is clearly reproduced.




Figure 8: Boundary mismatch functions for even (top) and odd (bottom) states. The zero crossings correspond to the physical eigenvalues.




Figure 9: Convergence of the energy error as a function of grid spacing h . The slopes are close to fourth order, as expected for the Numerov method.




Figure 10: Dependence of the energy error on the domain size $x_{\max}$. Errors decrease rapidly before reaching a plateau set by discretization and floating-point limits.

energy splitting between these states is analyzed as a function of the barrier height, illustrating quantum tunneling effects.

3.4 Finite square well

The finite square well demonstrates the flexibility of the solver. Unlike the infinite case, only a finite number of bound states exist, depending on the well depth. The numerical results capture this behavior correctly.

3.5 Scattering and resonant tunneling

In addition to bound states, we extend the solver to study one-dimensional scattering from finite potentials. For a given energy E , the wavefunction is propagated across the domain and decomposed into incoming and reflected components on the left and a transmitted component on the right.

The transmission and reflection coefficients $T(E)$ and $R(E)$ are computed from the amplitudes of these components and satisfy the conservation relation

$$T(E) + R(E) \approx 1.$$

For a single finite rectangular barrier, the numerical transmission is compared to the known analytical result, providing an additional validation of the method.

For a double-barrier potential, the transmission spectrum exhibits sharp resonance peaks corresponding to quasi-bound states in the central well. These resonances are clearly visible in $T(E)$ and illustrate quantum tunneling and interference effects.

3.6 Convergence study

We analyze numerical errors by varying both the grid spacing h and the domain size $x_{\max}$.

Grid convergence. For fixed $x_{\max}$, the energy error follows approximately

$$\Delta E \propto h^p,$$

where p is estimated from log-log fits and reported in the figure legends. The observed values $p \approx 3$ – 4 are consistent with high-order Numerov accuracy, with mild degradation due to boundary truncation and root-finding tolerances.

Dependence on domain size. For problems on unbounded domains (e.g., the harmonic oscillator), truncation at finite $x_{\max}$ introduces boundary error. For small $x_{\max}$ the wavefunction is artificially cut off, producing large errors. As $x_{\max}$ increases, the error decreases until it reaches a plateau set by discretization, root-finding tolerance, and floating-point precision. Thus this plot measures sensitivity to domain size rather than pure discretization convergence.

Convergence is primarily established using systems with known analytic solutions (square well and harmonic oscillator), and similar stable behavior is observed for more complex potentials such as the double well, although quantitative error estimates are less accessible there.

A comparison with a fourth-order Runge–Kutta (RK4) integrator was also performed for the harmonic oscillator. Both methods solve the same shooting problem, but RK4 evolves the wavefunction and its derivative simultaneously, whereas Numerov exploits the specific structure of the Schrödinger equation. Initially, RK4 appeared more accurate due to a lower-order derivative approximation in the Numerov boundary condition. After correcting this issue, both methods exhibit comparable convergence behavior, with Numerov achieving similar or better accuracy for the same grid spacing. This highlights the importance of treating boundary conditions with the same order of accuracy as the integration scheme.

Representative plots include wavefunctions, probability densities, root-finding diagnostics, convergence curves, and energy splitting as a function of potential parameters.

4 Discussion

The correctness of the code is verified through multiple checks. Analytical benchmarks confirm the accuracy of the computed eigenvalues, while normalization ensures that the wavefunctions remain physically meaningful. Convergence studies provide strong evidence that the numerical results are stable with respect to discretization parameters.

The double-well analysis highlights a key physical phenomenon: tunneling-induced energy splitting. As the barrier height increases, the splitting decreases, reflecting reduced overlap between the localized states. This demonstrates that the numerical method captures not only correct energies but also meaningful physical behavior.

The results also illustrate typical numerical considerations, such as sensitivity to grid resolution and domain size, and emphasize the importance of matching the accuracy of boundary conditions to that of the integration scheme.

The agreement with analytical results, together with consistent convergence behavior and well-behaved root-finding diagnostics, demonstrates the correctness and robustness of the numerical implementation. Differences in observed convergence rates across states are attributed to boundary truncation and solver tolerances rather than defects in the underlying method.

5 Conclusion

The Numerov method, combined with parity-based shooting and robust root-finding, provides an efficient and accurate tool for one-dimensional bound-state problems, with controlled errors from discretization, domain truncation, and floating-point precision. In addition to reproducing known spectra, it captures physically interesting effects such as tunneling-induced energy splitting in a double well.

The extension to scattering problems further demonstrates the flexibility of the approach as a secondary component, allowing the study of transmission, reflection, and resonant tunneling phenomena within the same numerical framework.

References

- [1] Tao Pang. *An Introduction to Computational Physics*. 2nd ed. Cambridge University Press, 2006.




Figure 11: Wavefunctions for the first four harmonic oscillator states plotted together with the potential.




Figure 12: Comparison of Numerov and RK4 shooting for the harmonic oscillator. Both methods show fourth-order convergence, while Numerov gives a smaller maximum energy error at the same grid spacing.