

“Calculus 3”

Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo

Day 15

1

Any Reminders? Any Questions?

- Quiz on Friday on triple integrals in cartesian, cylindrical, and spherical coordinates.
- I will be away Wednesday-Thursday at a conference so no in-person class on 3/12 – watch videos instead!

2



ALVARO: Start the recording!



3

“Calculus 3”



Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo

Changes of Variables in Integrals



4



Today – Changes of Variables in Double and Triple Integrals

- Changes of Variables in Single Integrals
- Changes of Variables in Double Integrals
- Changes of Variables in Triple Integrals

5

Changes of Variables in Single Integrals (aka u-Substitution)



6

Example: Evaluate the (single) integral

$$\int_0^2 x e^{x^2} dx$$

7

Changes of Variables in Double Integrals

Example: Evaluate the (double) integral


$$\int_{-1}^1 \int_0^{\sqrt{1-x^2}} \sqrt{x^2 + y^2} dy dx$$

8

Changes of Variables in Double Integrals (in Polar)

9

Changes of Variables in Double Integrals (in General)



10

Example: Show that the change of variables $x = \frac{u}{2}, y = \frac{v}{2}$ sends the unit square (in x,y coordinates, in the first quadrant) to a square of side length 2 (in u,v).

11

Example: Show that the change of variables $x = \frac{u}{2}, y = \frac{v}{2}$ sends the unit square (in the first quadrant) to a square of side length 2.

[Extra]

12

Example: Evaluate the (double) integrals

$$\int_0^1 \int_0^1 e^{2x+2} dy dx \quad \text{and} \quad \int_0^2 \int_0^2 e^{u+v} dv du$$

13

Example: Use the change of variables $x + y = u, y = v$ instead???

14

Example: Evaluate the (double) integrals

$$\int_0^1 \int_0^1 e^{2x+2} dy dx \quad \text{and} \quad \int_0^1 \int_v^{v+1} e^{2u} du dv$$

15

Example: Show that the change of variables $x = R \cos(\theta)$, $y = R \sin(\theta)$ sends the square of side lengths R and $\pi/4$ to a quarter circle of radius R . (Then, compute the areas of both regions.)


16

[Extra]

Example: Show that the change of variables $x = R \cos(\theta)$, $y = R \sin(\theta)$ sends the square of side lengths R and $\pi/4$ to a quarter circle of radius R . (Then, compute the areas of both regions.)

17


Example: Show that the change of variables $x = u^2 - v^2$, $y = 2uv$ sends the region S to the region R , and vice-versa.



18


[Extra]

Example: Show that the change of variables $x = u^2 - v^2, y = 2uv$ sends the region S to the region R , and vice-versa.



19

Changes of Variables in Double Integrals (in General)



Suppose that T is a C^1 transformation whose Jacobian is nonzero and that T maps a region S in the uv -plane onto a region R in the xy -plane. Suppose that f is continuous on R and that R and S are type I or type II plane regions. Suppose also that T is one-to-one, except perhaps on the boundary of S . Then

$$\iint_R f(x, y) \, dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv$$

20

Changes of Variables in Double Integral: the Jacobian

$$\iint_R f(x, y) \, dA = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv$$

The **Jacobian** of the transformation T given by $x = g(u, v)$ and $y = h(u, v)$ is

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$



The Jacobian is named after the German mathematician Carl Gustav Jacob Jacobi (1804–1851). Although the French mathematician Cauchy first used these special determinants involving partial derivatives, Jacobi developed them into a method for evaluating multiple integrals.

21

Example: Find the jacobian of the change of variables
 $x = R \cos(\theta), y = R \sin(\theta)$

22


Example: Find the jacobian of the change of variables $x = \frac{u}{2}, y = \frac{v}{2}$

23

Example: Find the jacobian of the change of variables
 $x + y = u, y = v$

24

Example: Find the jacobian of the change of variables
 $x = u^2 - v^2, y = 2uv$




25

Example: Use the change of variables $x = u^2 - v^2, y = 2uv$ to evaluate

$$\iint_R y \, dA$$

 where R is the region bounded by the parabolas
 $y^2 = 4 - 4x$ and $y^2 = 4 + 4x$ and $y \geq 0$.




26

[Extra]

Example: Use the change of variables $x = u^2 - v^2, y = 2uv$ to evaluate

$$\iint_R y \, dA$$

where R is the region bounded by the parabolas $y^2 = 4 - 4x$ and $y^2 = 4 + 4x$ and $y \geq 0$.



27

Example: Evaluate the integral $\iint_R e^{\frac{x+y}{x-y}} \, dA$

where R is the trapezoidal region with vertices $(1,0)$, $(2,0)$, $(0,-2)$, $(0,-1)$.

28

[Extra]

Example: Evaluate the integral $\iint_R e^{\frac{x+y}{x-y}} dA$
 where R is the trapezoidal region with vertices
 $(1,0)$, $(2,0)$, $(0,-2)$, $(0,-1)$.

29

Example: Use the change of variables $x = 2u$, $y = 3v$ to evaluate the integral

$$\iint_R x^2 dA,$$

where R is region bounded by ellipse $9x^2 + 4y^2 = 36$.

30

[Extra]

Example: Use the change of variables $x = 2u, y = 3v$ to evaluate the integral

$$\iint_R x^2 dA,$$

where R is region bounded by ellipse $9x^2 + 4y^2 = 36$.

31

Example: Use the change of variables to evaluate the integral

$$\iint_R \sin(4x^2 + y^2) dA,$$

where R is the region in the first quadrant bounded by ellipse $4x^2 + y^2 = 1$.

32

[Extra]

Example: Use the change of variables to evaluate the integral

$$\iint_R \sin(4x^2 + y^2) dA,$$

where R is the region in the first quadrant bounded by ellipse $4x^2 + y^2 = 1$.

33

“Calculus 3”

Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo

Changes of Variables in Triple Integrals

34

Changes of Variables in Triple Integrals (in General)

$$\iiint_R f(x, y, z) \, dV = \iiint_S f(x(u, v, w), y(u, v, w), z(u, v, w)) \left| \frac{\partial(x, y, z)}{\partial(u, v, w)} \right| \, du \, dv \, dw$$

$$x = g(u, v, w) \quad y = h(u, v, w) \quad z = k(u, v, w)$$

$$\frac{\partial(x, y, z)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

35

Example: Spherical Coordinates

$$x = \rho \sin \phi \cos \theta \quad y = \rho \sin \phi \sin \theta \quad z = \rho \cos \phi$$

We compute the Jacobian as follows:

$$\begin{aligned} \frac{\partial(x, y, z)}{\partial(\rho, \theta, \phi)} &= \begin{vmatrix} \sin \phi \cos \theta & -\rho \sin \phi \sin \theta & \rho \cos \phi \cos \theta \\ \sin \phi \sin \theta & \rho \sin \phi \cos \theta & \rho \cos \phi \sin \theta \\ \cos \phi & 0 & -\rho \sin \phi \end{vmatrix} \\ &= \cos \phi \begin{vmatrix} -\rho \sin \phi \sin \theta & \rho \cos \phi \cos \theta \\ \rho \sin \phi \cos \theta & \rho \cos \phi \sin \theta \end{vmatrix} - \rho \sin \phi \begin{vmatrix} \sin \phi \cos \theta & -\rho \sin \phi \sin \theta \\ \sin \phi \sin \theta & \rho \sin \phi \cos \theta \end{vmatrix} \\ &= \cos \phi (-\rho^2 \sin \phi \cos \phi \sin^2 \theta - \rho^2 \sin \phi \cos \phi \cos^2 \theta) \\ &\quad - \rho \sin \phi (\rho \sin^2 \phi \cos^2 \theta + \rho \sin^2 \phi \sin^2 \theta) \\ &= -\rho^2 \sin \phi \cos^2 \phi - \rho^2 \sin \phi \sin^2 \phi = -\rho^2 \sin \phi \end{aligned}$$

36


Questions?



37

Thank you

Until next time.



38