

“Calculus 3”

Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo

Day 19

1

Any Reminders? Any Questions?

- Quiz on Friday (discussion) covers 13.1-13.3 (vector functions, their integrals and derivatives, and arc length)
- Office hours Mondays 1-2pm online
- Office hours Thursday 3:30-4:30 at MONT 233
- Calc Night Thursday 6:30-8:30 at MONT 104

2



ALVARO: Start the recording!



3

“Calculus 3”



Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo





Line Integrals



4



Line Integrals



$$\int_C f(x, y) ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*, y_i^*) \Delta s_i$$

5

Line Integrals



$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

6

Example: Evaluate $\int_C (2 + x^2 y) ds$ where C is the upper half of the unit circle.

7

[Extra Space]

Example: Evaluate $\int_C (2 + x^2 y) ds$ where C is the upper half of the unit circle.

8

Example: Evaluate $\int_C (2x) \, ds$ where C consists of the arc of the parabola $y = x^2$ from $(0,0)$ to $(1,1)$ followed by a line segment from $(1,1)$ to $(1,2)$.

9

[Extra Space]

Example: Evaluate $\int_C (2x) \, ds$ where C consists of the arc of the parabola $y = x^2$ from $(0,0)$ to $(1,1)$ followed by a line segment from $(1,1)$ to $(1,2)$.

10

Application of Line Integrals: Center of Mass

$$\bar{x} = \frac{1}{m} \int_C x \rho(x, y) ds \quad \bar{y} = \frac{1}{m} \int_C y \rho(x, y) ds$$

11

Example: A wire takes the shape of the top half of a semi-circle of radius 1 and it is thicker near its base than near the top. Find the center of mass of the wire if the linear density at any point is proportional to its distance from the line $y = 1$.

12

[Extra Space]

Example: A wire takes the shape of the top half of a semi-circle of radius 1 and it is thicker near its base than near the top. Find the center of mass of the wire if the linear density at any point is proportional to its distance from the line $y = 1$.

13

Line integrals with respect to x and y

$$\int_C f(x, y) dx = \int_a^b f(x(t), y(t)) x'(t) dt$$

$$\int_C f(x, y) dy = \int_a^b f(x(t), y(t)) y'(t) dt$$

$$\int_C P(x, y) dx + \int_C Q(x, y) dy = \int_C P(x, y) dx + Q(x, y) dy$$

14

Example: Evaluate $\int_C y^2 dx + x dy$ where C is the line segment from $(-5,-3)$ to $(0,2)$.

15

[Extra Space]

Example: Evaluate $\int_C y^2 dx + x dy$ where C is the line segment from $(-5,-3)$ to $(0,2)$.

16

Example: Evaluate $\int_C y^2 dx + x dy$ where C is the arc of the parabola $x = 4 - y^2$ from $(-5,-3)$ to $(0,2)$.

17

Example: Evaluate $\int_C y^2 dx + x dy$ where C is the line segment from $(-5,-3)$ to $(0,2)$.

[Extra Space]


18

Example:



19

Questions?



20



ALVARO: Start the recording!



21

“Calculus 3”



Multi-Variable Calculus



Instructor: Álvaro Lozano-Robledo

Line Integrals... in Space



22

Line Integrals



$$\int_C f(x, y, z) \, ds = \int_a^b f(x(t), y(t), z(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} \, dt$$

23

Line Integrals

$$\int_C f(x, y, z) \, ds = \int_a^b f(x(t), y(t), z(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} \, dt$$

$$\int_C P(x, y, z) \, dx + Q(x, y, z) \, dy + R(x, y, z) \, dz$$

24

Example: Evaluate $\int_C y \sin(z) ds$ where C is the helix given by
 $(\cos(t), \sin(t), t)$, $0 \leq t \leq 2\pi$

25

[Extra Space]

Example: Evaluate $\int_C y \sin(z) ds$ where C is the helix given by
 $(\cos(t), \sin(t), t)$, $0 \leq t \leq 2\pi$

26

Example: Evaluate $\int_C y \, dx + z \, dy + x \, dz$ where C consists of the line segment from $(2,0,0)$ to $(3,4,5)$, followed by the line segment from $(3,4,5)$ to $(3,4,0)$.

27

[Extra Space]

Example: Evaluate $\int_C y \, dx + z \, dy + x \, dz$ where C consists of the line segment from $(2,0,0)$ to $(3,4,5)$, followed by the line segment from $(3,4,5)$ to $(3,4,0)$.

28

Questions?

29



ALVARO: Start the recording!



30

“Calculus 3”

Multi-Variable Calculus


Instructor: Álvaro Lozano-Robledo

Vector Fields


31

Vector Fields

Velocity vector fields showing San Francisco Bay wind patterns on a particular spring day



(a) 6:00 PM



(b) 6:00 AM

32

Vector Fields




33

Vector Fields

Let D be a set in $\mathbb{R}^2$ (a plane region). A **vector field on $\mathbb{R}^2$** is a function $\mathbf{F}$ that assigns to each point (x, y) in D a two-dimensional vector $\mathbf{F}(x, y)$.

Let E be a subset of $\mathbb{R}^3$. A **vector field on $\mathbb{R}^3$** is a function $\mathbf{F}$ that assigns to each point (x, y, z) in E a three-dimensional vector $\mathbf{F}(x, y, z)$.



34

Example: Sketch a vector field on the plane given by

$$F(x, y) = (-y, x) = -y \cdot i + x \cdot j$$

35

[Extra Space]

Example: Evaluate $\int_C y \sin(z) ds$ where C is the helix given by $(\cos(t), \sin(t), t)$, $0 \leq t \leq 2\pi$

36


Questions?



37

Thank you

Until next time.



38