


Extending the reach of ab initio theory: Valence space IMSRG


Ragnar Stroberg
TRIUMF

ARIS In the Mountains
Keystone, Colorado
May 30, 2017






Outline



1. In-medium SRG for a valence space
2. Ensemble normal ordering
3. Selected results
4. Outlook



Work in collaboration with: A. Calci, J. Holt, P. Navrátil, S. Bogner, H. Hergert, N. Parzuchowski, K. Hebeler, R. Roth, A. Schwenk, J. Simonis, C. Stumpf, G. Hagen, and T. Morris

- $H|\Psi\rangle = E|\Psi\rangle$ is too difficult to solve.

$$H|\Psi\rangle = E|\Psi\rangle$$

complicated!

- Perform unitary transformation U (implicit change of basis) to make it easier to solve.

- Iterative/guess-and-check approach.

$$U \equiv e^{\Omega} = e^{\Omega_n} e^{\Omega_{n-1}} \dots e^{\Omega_2} e^{\Omega_1}$$

- Alternatively, $\Omega_n \rightarrow \eta ds \Rightarrow$ flow equation

- Computational effort dominated by commutator evaluation.


- $H|\Psi\rangle = E|\Psi\rangle$ is too difficult to solve.
- Perform unitary transformation $\tilde{H} = U H U^\dagger$ (implicit change of basis) so SE is easier to solve.

- Iterative/guess-and-check approach.

$$U \equiv e^\Omega = e^{\Omega_n} e^{\Omega_{n-1}} \dots e^{\Omega_2} e^{\Omega_1}$$


- Alternatively, $\Omega_n \rightarrow \eta ds \Rightarrow$ flow equation

- Computational effort dominated by commutator evaluation.



- $H|\Psi\rangle = E|\Psi\rangle$ is too difficult to solve.
- Perform unitary transformation $\tilde{H} = U H U^\dagger$ (implicit change of basis) so SE is easier to solve.
- Iterative/guess-and-check approach.


$$U \equiv e^\Omega = e^{\Omega_n} e^{\Omega_{n-1}} \dots e^{\Omega_2} e^{\Omega_1}$$



- Alternatively, $\Omega_n \rightarrow \eta ds \Rightarrow$ flow equation

$$\frac{dH(s)}{ds} = [\eta(s), H(s)].$$

- Computational effort dominated by commutator evaluation.



- $H|\Psi\rangle = E|\Psi\rangle$ is too difficult to solve.
- Perform unitary transformation $\tilde{H} = U H U^\dagger$ (implicit change of basis) so SE is easier to solve.



- Iterative/guess-and-check approach.

$$U \equiv e^\Omega = e^{\Omega_n} e^{\Omega_{n-1}} \dots e^{\Omega_2} e^{\Omega_1}$$



- Alternatively, $\Omega_n \rightarrow \eta ds \Rightarrow$ flow equation

$$\frac{dH(s)}{ds} = [\eta(s), H(s)].$$


- Computational effort dominated by commutator evaluation.



$$\frac{dH(s)}{ds} = [\eta(s), H(s)]$$



What we would like:




Why “in-medium”?

$$H = \underbrace{E_0}_{0\text{-body}} + \underbrace{\sum_{ij} H_{ij} \{a_i^\dagger a_j\}}_{1\text{-body}} + \underbrace{\frac{1}{4} \sum_{ijkl} H_{ijkl} \{a_i^\dagger a_j^\dagger a_l a_k\}}_{2\text{-body}} + \underbrace{\frac{1}{36} \sum_{ijklmn} H_{ijklmn} \{a_i^\dagger a_j^\dagger a_k^\dagger a_n a_m a_l\}}_{3\text{-body}} + \dots$$

- In general, the transformation U will induce 4-body, 5-body, etc. forces 😡


- Write H in terms of excitations out of reference $|\Phi_0\rangle$
- Normal ordering: $\langle \Phi_0 | \{a_1^\dagger \dots a_N^\dagger a_N \dots a_1\} | \Phi_0 \rangle = 0$
- If $|\Phi_0\rangle \approx |\Psi\rangle$, higher-body terms are negligible
- **Truncate all operators at 2 body level (NO2B)**



Why “in-medium”?


$$H = \underbrace{E_0}_{0\text{-body}} + \underbrace{\sum_{ij} H_{ij} \{a_i^\dagger a_j\}}_{1\text{-body}} + \underbrace{\frac{1}{4} \sum_{ijkl} H_{ijkl} \{a_i^\dagger a_j^\dagger a_l a_k\}}_{2\text{-body}} + \underbrace{\frac{1}{36} \sum_{ijklmn} H_{ijklmn} \{a_i^\dagger a_j^\dagger a_k^\dagger a_n a_m a_l\}}_{3\text{-body}} + \dots$$


- In general, the transformation U will induce 4-body, 5-body, etc. forces 😡
- Write H in terms of excitations out of reference $|\Phi_0\rangle$
- Normal ordering: $\langle \Phi_0 | \{a_1^\dagger \dots a_N^\dagger a_N \dots a_1\} | \Phi_0 \rangle = 0$
- If $|\Phi_0\rangle \approx |\Psi\rangle$, higher-body terms are negligible
- Truncate all operators at 2 body level (NO2B)



Why “in-medium”?


$$H = \underbrace{E_0}_{0\text{-body}} + \underbrace{\sum_{ij} H_{ij} \{a_i^\dagger a_j\}}_{1\text{-body}} + \underbrace{\frac{1}{4} \sum_{ijkl} H_{ijkl} \{a_i^\dagger a_j^\dagger a_l a_k\}}_{2\text{-body}} + \underbrace{\frac{1}{36} \sum_{ijklmn} H_{ijklmn} \{a_i^\dagger a_j^\dagger a_k^\dagger a_n a_m a_l\}}_{3\text{-body}} + \dots$$

- In general, the transformation U will induce 4-body, 5-body, etc. forces 
- Write H in terms of excitations out of reference $|\Phi_0\rangle$
- Normal ordering: $\langle \Phi_0 | \{a_1^\dagger \dots a_N^\dagger a_N \dots a_1\} | \Phi_0 \rangle = 0$
- If $|\Phi_0\rangle \approx |\Psi\rangle$, higher-body terms are negligible
- **Truncate all operators at 2 body level (NO2B)**




Tskukiyama et al (2011)


Solving the many-body problem




- Decouple a 1×1 sub-block
- Use SRG to suppress excitations out of $|\Phi_0\rangle$
- After decoupling, energy is $E_0 = \langle \Phi_0 | \tilde{H} | \Phi_0 \rangle$




- Open shell systems: multiple (quasi-) degenerate configurations \Rightarrow strong mixing, $|\Phi_0\rangle \approx |\Psi\rangle$
- Single Slater determinant may not have good total angular momentum J
- Large rotation angle \Rightarrow induced many-body forces
- Strategies:
 - Break symmetries and restore afterward
 - Construct multi-configuration reference, then decouple (multi-reference IM-SRG)
 - Decouple a subset of configurations, then construct state from them using standard shell model machinery, e.g NuShellX (valence-space IMSRG)




- Open shell systems: multiple (quasi-) degenerate configurations \Rightarrow strong mixing, $|\Phi_0\rangle \approx |\Psi\rangle$
- Single Slater determinant may not have good total angular momentum J
- Large rotation angle \Rightarrow induced many-body forces
- Strategies:
 - Break symmetries and restore afterward
 - Construct multi-configuration reference, then decouple (multi-reference IM-SRG)
 - Decouple a subset of configurations, then construct state from them using standard shell model machinery, e.g NuShellX (valence-space IMSRG)






What reference should be used when decoupling a valence space?
(i.e. what is the “medium”?)




Obvious choice: the inert core, e.g. ^{16}O .








Reference:
inert core




Reference:
inert core


Reference:
nearest
closed shell

(Ensemble gives
very similar results)











	EM 500/400	EM 1.8/2.0
NN	N^3LO	same
	$\Lambda_{2N} = 500$ MeV	same
	non-local regulator	same
	fit to NN scattering, 2H	same
	$\lambda_{SRG} = 1.88 \text{ fm}^{-1}$	\approx same
3N	N^2LO	same
	$\Lambda_{3N} = 400$ MeV	\approx same
	local regulator	non-local regulator
	fit to 3H BE, $t_{1/2}$	fit to 3H BE, 4He r_{ch}
	consistently SRG evolved	no SRG for 3N








	EM 500/400	EM 1.8/2.0
NN	N^3LO $\Lambda_{2N} = 500 \text{ MeV}$ non-local regulator fit to NN scattering, 2H $\lambda_{SRG} = 1.88 \text{ fm}^{-1}$	same same same same \approx same
3N	N^2LO $\Lambda_{3N} = 400 \text{ MeV}$ local regulator fit to 3H BE, $t_{1/2}$ consistently SRG evolved	same \approx same non-local regulator fit to 3H BE, 4He r_{ch} no SRG for 3N













	EM 500/400	EM 1.8/2.0
NN	N^3LO $\Lambda_{2N} = 500$ MeV non-local regulator fit to NN scattering, 2H $\lambda_{SRG} = 1.88$ fm $^{-1}$	same same same same \approx same
3N	N^2LO $\Lambda_{3N} = 400$ MeV local regulator fit to 3H BE, $t_{1/2}$ consistently SRG evolved	same \approx same non-local regulator fit to 3H BE, 4He r_{ch} no SRG for 3N













- Neither interaction is fully consistent however...
- Saturation properties are important for finite nuclei




























































Baumann et al. Nature (2007), Möller et al. (1995), Samyn et al. (2004), Holt et al. (in prep.)


- Only difference: choice of initial NN force.
- Identical procedure for fitting 3N contact terms.
- Based on few-body data, all interactions are equally good.
- Big differences for finite nuclei.
- 1.8/2.0 EM interaction is “magic”, i.e. lucky.















Limited by truncation of 3N matrix elements


$$E3_{max} = e_1 + e_2 + e_3$$





✓ converged




≈ almost converged

✗ not converged









Bogner et al. PRL (2014), SRS et al. PRC(R) (2016),



Bogner et al. PRL (2014), SRS et al. PRC(R) (2016), Simonis et al arXiv:1704.02915 (2017),



Electroweak Observables




Bogner et al. PRL (2014), SRS et al. PRC(R) (2016), Simonis et al arXiv:1704.02915 (2017), Parzuchowski et al. arXiv:1705.05511 (2017)

What does the future hold?

(Technical developments)


- Quantification of many-body uncertainty
 - Perturbative estimation of omitted 3-body terms
 - Invariant trace
 - Full IMSRG(3): Include 3-body terms throughout the calculation
- Heavy-mass frontier
 - Improve handling of 3N forces
- Decoupling arbitrary valence spaces
 - Island(s) of inversion
 - Parity-changing transitions, e.g. $E1$
- Improved basis
 - Two-frequency oscillator basis for halo systems
 - Explicit inclusion of collective modes



What does the future hold?

(Technical developments)


- Quantification of many-body uncertainty
 - Perturbative estimation of omitted 3-body terms
 - Invariant trace
 - Full IMSRG(3): Include 3-body terms throughout the calculation
- Heavy-mass frontier
 - Improve handling of 3N forces
- Decoupling arbitrary valence spaces
 - Island(s) of inversion
 - Parity-changing transitions, e.g. $E1$
- Improved basis
 - Two-frequency oscillator basis for halo systems
 - Explicit inclusion of collective modes



What does the future hold?

(Technical developments)




- Quantification of many-body uncertainty
 - Perturbative estimation of omitted 3-body terms
 - Invariant trace
 - Full IMSRG(3): Include 3-body terms throughout the calculation
- Heavy-mass frontier
 - Improve handling of 3N forces
- Decoupling arbitrary valence spaces
 - Island(s) of inversion
 - Parity-changing transitions, e.g. $E1$
- Improved basis
 - Two-frequency oscillator basis for halo systems
 - Explicit inclusion of collective modes



What does the future hold?

(Technical developments)


- Quantification of many-body uncertainty
 - Perturbative estimation of omitted 3-body terms
 - Invariant trace
 - Full IMSRG(3): Include 3-body terms throughout the calculation
- Heavy-mass frontier
 - Improve handling of 3N forces
- Decoupling arbitrary valence spaces
 - Island(s) of inversion
 - Parity-changing transitions, e.g. $E1$
- Improved basis
 - Two-frequency oscillator basis for halo systems
 - Explicit inclusion of collective modes



What does the future hold? (Observables)

- Radii / isotope shifts
- $E0$ transitions
- Chiral currents for $M1$, Gamow-Teller operators
- Neutrinoless double beta decay
- Structure factors for dark matter detection
- Superaligned Fermi decays
- Suggestions?

- Valence space IM-SRG with ensemble normal ordering allows access to all nuclei up to $A \sim 100$
- Reach in A is presently limited by E_{3max} truncation
- Consistent operators for other observables can be obtained
- Chiral interactions still need work (magic notwithstanding)
- Next goal: how to reliably estimate truncation error?



Collaborators:



A. Calci, J. Holt, P. Navrátil, C. Payne, O. Drozdowski,
D. Fullerton, C. Gwak, L. Kemmler, S. Leutheusser, D. Livermore



NSCL/MSU S. Bogner, H. Hergert, N. Parzuchowski



TU Darmstadt K. Hebeler, R. Roth, A. Schwenk, J. Simonis, C. Stumpf



ORNL/UT G. Hagen, T. Morris