

<b>Mastitis</b>	<p><i>Redness, tenderness, and increased heat, in a localized area. Palpable resistance. High fever. CRP↑</i></p> <p>Breast feeding (empty the breast) pump if needed  Culture if wound  Flukloxacillin (Heracillin) 1g x3  If abscess: Ultrasound drainage and culture.</p>
<b>Endometritis</b>	<p><i>Abd. pain. Tender uterus. Malodorous bloody discharge. CRP↑</i></p> <p>Cervical culture (streptococcus), blood culture  Methergin + antibiotics:  Within days: pip/tazo 4g x4  Late: amoxi/klav 500mg/125mg x3 + metronidazol 400mg x3 5-10d</p>
<b>Wound infection</b>	<p><i>Redness, pus</i></p> <p>Wound culture  Debridement</p>
<b>Pneumonia</b>	<p><i>Coughing, pleural pain</i></p> <p>Clinical examination, X-ray if needed  If uncomplicated: PcG</p>
<b>Urinary tract inf.</b>	<p><i>Urinary urgency and tenderness over kidneys or bladder</i></p> <p>Urine culture  Antibiotics, e.g. Selexid (CAVE Furadantin)</p>
<b>Thrombosis</b>	<p><i>Signs of pulmonary embolus or DVT</i></p> <p>Ultrasound legs  Pulmonary CT if needed</p>