

Two methods of solving Poisson's equation

Anna Lina P. Sjur and Jan-Adrian H. Kallmyr

September 5, 2018

Abstract

State problem. Briefly describe method and data. Summarize main results.

1 Introduction

Physics is a field concerned with the behaviour of nature, and nature is everchanging. It is therefore no surprise that differential equations appear everywhere in physics. From global climate dynamics to statistical mechanics, what we find is that differential equations, often many and coupled, are required to explain or model the phenomena. For such large models, efficiency is important, as we would, for example, like to have timely weather forecasts. One way to make a model more efficient is by using an efficient algorithm for solving differential equations.

In this report, we compare two different numerical methods of solving linear second-order differential equations with the Dirichlet boundary conditions. To do this, we will solve the one-dimensional Poisson's equation:

$$\frac{d^2\phi}{dr^2} = -4\pi\rho(r). \quad (1)$$

In the methods section, we develop an approximation for the 2nd derivative to the 2nd order. We will then solve eq. 1 numerically, using gaussian elimination and lower-upper decomposition. As for the former, we will further specialise it to solve eq. 1 more efficiently. Next, in the results section we present the comparison between our numerical solutions and the analytical solution, as well as the error. We

then compare the efficiency of all three algorithms, and finally, in the discussion section we will consider the advantages and disadvantages of each algorithm, and discuss their uses.

2 Methods

3 Results




Figure 1: text

n	t_g/t_s	t_{LU}/t_s
10	2.08	
10 ²	1.89	
10 ³	1.48	
10 ⁴	1.43	
10 ⁵	1.39	
10 ⁶	1.41	
10 ⁷	1.39	

Table 1:

4 Discussion

References

- [Górski et al.(1994)] Górski, K. M., Hinshaw, G., Banday, A. J., Bennett, C. L., Wright, E. L., Kogut, A., Smoot, G. F., and Lubin, P. 1994, ApJL, 430, 89