

# **Analysis 2 - Thomas Mountford**

Alp Ozen

Spring 2019

## Contents

<b>1</b>	<b>Reviewing <math>\mathbf{R}^n</math></b>	<b>2</b>
1.1	Introducing topological properties on $\mathbf{R}^n$ . . . . .	2

# 1 Reviewing $\mathbb{R}^n$

Let's recall that  $\mathbb{R}^n$  is a Euclidean vector space. We define a scalar product on  $\mathbb{R}^n$  as follows:

**Definition 1.**

$$\langle \cdot, \cdot \rangle: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$$

1.  $\langle x, x \rangle \geq 0$
2.  $\langle x, y \rangle = \langle y, x \rangle$
3.  $\langle ax + by, z \rangle = a \langle x, z \rangle + b \langle y, z \rangle$

## 1.1 Introducing topological properties on $\mathbb{R}^n$

A *norm* is defined as function that maps some real vector space  $E$  to  $\mathbb{R}$  and satisfies:

- 1)  $|x| \geq 0 \forall x \in E, |x| = 0 \iff x = 0$
- 2)  $|\lambda \cdot x| = |\lambda| \cdot |x|$
- 3)  $|x + y| \leq |x| + |y|$

In our intuitive understanding of  $\mathbb{R}^n$  we are actually thinking about the Euclidian space  $\mathbb{R}^n$  equipped with the Euclidian norm.

**Definition 2. Euclidian norm**

$$|x|_2 = \sqrt{\langle x, x \rangle} = \left( \sum_i^n x_k^2 \right)^{\frac{1}{2}}$$

And from this naturally follows the definition of Euclidian distance:

**Definition 3.**

$$d(x, y) = |x - y|$$

We note that  $d$  satisfies the same 3 properties as the norm. Thus, the couple  $(E, d)$  is called a metric space.

And now more definitions:

**Definition 4. Open sets**

1. **Open ball**  $B(a, r) := \{x \in \mathbb{R}^n : d(x, a) < r\}$
2. **Open subset** Some subset  $S \subset \mathbb{R}^n$  is open if  $\forall x \in \mathbb{R}^n, \exists \epsilon > 0 B(x, \epsilon) \subset S$
3. **Closed subset** Some  $S$  is closed if  $\mathbb{R}^n - S$  is open, note that the empty set and  $\mathbb{R}^n$  are both open and closed.
4. **The interior and boundary of a set**  $a$  is in the interior of  $S$  if  $\exists \epsilon > 0 B(a, \epsilon) \subset S$  and  $b$  is in the boundary of a set  $S$  if any  $B(a, \epsilon)$  contains points from both  $S$  and  $\mathbb{R}^n - S$ . The set of all interior points is denoted  $(^\circ S)$  and set of all boundary points is denoted  $\partial S$
5. **Closure of a set**  $a$  is a closure of  $S$  if for any  $B(a, \epsilon)$  we have  $B(a, \epsilon) \cap S \neq \emptyset$