

Suitability prevalence area index in late quaternary explains genetic diversity in Tassel eared Squirrels

Norma Hernandez, Angel Robles Nathan Upham

2023-06-27

Abstract

The current distributions of species do not always correspond to their historical distributions over evolutionarily significant periods. This is because environmental conditions are not static over time; species tend to distribute where conditions are most favorable.

It follows from this relationship between environment and species distribution that population size also varies with time. This variation in population size is related to the effective population size N_e , so that an index reflecting changes in environmental conditions in geography might be expected to be related to N_e and thus to indicators of population structure such as the fixation index F_{st} . Thus, it is possible to relate patterns of changes in the distribution of species to the genetic structure of their populations using a statistical model that explains this relationship.

With this approach we can predict the geographic pattern of population structure from environmental information. This approach is strongly driven by advances in currently available climate simulations (Leonardi et al. 2023; Krapp et al. 2021), as well as next generation sequencing data, and supported by both ecological niche (Thorup et al. 2021; Nogués-Bravo 2009) and population genetics theories (Lira-Noriega and Manthey 2014).

In this work we propose a method to find the Suitability Prevalence Area (SPA) as an index with a double purpose: 1) to find endemic areas to delimit the historical distribution of the species and 2) to explain the patterns of genetic diversity.

To obtain the SPA, we performed a historical reconstruction of the geographical range back to 120 000 BC at 2 000 year intervals and recorded the environmental suitability at each site.

Subsequently, we delimited historical endemic areas to locations where the prevalence of suitability remained at 90% during this period. From the fixation index calculated with respect to populations in the historical endemic areas, a statistical model of the fixation index as a function of SPA was performed. With this statistical model, the fixation index values were projected to the current distribution to obtain a map with the geographic pattern of this index.

As a case study we consider squirrels (*Sciurus aberti*), a species currently distributed in disjunct patches from the southern Rocky Mountains in the United States to the northern Sierra Madre Occidental in Mexico for which it is possible to find reliable information on both the genetic structure of its populations and its current distribution (Bono et al. 2018; Burgin et al. 2018).

Our results reveal that suitability prevalence corresponds to the fixation index of *S. aberti* populations with respect to a source population. Populations closer to the historical endemic area present a higher genetic diversity and a lower F_{st} value. Finally, this study allows us to add a biogeographic explanation to the results obtained with population genetic methods and to generate maps of this structure as tools to support conservation with a perspective that integrates both population genetics and historical patterns of species distribution.

- Bono, Jeremy M, Helen K Pigage, Peter J Wettstein, Stephanie A Prosser, and Jon C Pigage. 2018. "Genome-Wide Markers Reveal a Complex Evolutionary History Involving Divergence and Introgression in the Abert's Squirrel (*Sciurus Aberti*) Species Group." *BMC Evolutionary Biology* 18 (1): 1–17.
- Burgin, Connor J, Jocelyn P Colella, Philip L Kahn, and Nathan S Upham. 2018. "How Many Species of Mammals Are There?" *Journal of Mammalogy*. Oxford University Press US.
- Krapp, Mario, Robert M Beyer, Stephen L Edmundson, Paul J Valdes, and Andrea Manica. 2021. "A Statistics-Based Reconstruction of High-Resolution Global Terrestrial Climate for the Last 800,000 Years." *Scientific Data* 8 (1): 228.
- Leonardi, Michela, Emily Y Hallett, Robert Beyer, Mario Krapp, and Andrea Manica. 2023. "Pastclim 1.2: An r Package to Easily Access and Use Paleoclimatic Reconstructions." *Ecography*, e06481.
- Lira-Noriega, Andrés, and Joseph D Manthey. 2014. "Relationship of Genetic Diversity and Niche Centrality: A Survey and Analysis." *Evolution* 68 (4): 1082–93.
- Nogués-Bravo, David. 2009. "Predicting the Past Distribution of Species Climatic Niches." *Global Ecology and Biogeography* 18 (5): 521–31.
- Thorup, Kasper, Lykke Pedersen, Rute R Da Fonseca, Babak Naimi, David Nogués-Bravo, Mario Krapp, Andrea Manica, et al. 2021. "Response of an Afro-Palearctic Bird Migrant to Glaciation Cycles." *Proceedings of the National Academy of Sciences* 118 (52): e2023836118.