

# Stellar occultation predictions for 10 irregular satellites of giant planets and Triton for 2015-2017

A. R. Gomes-Júnior<sup>1</sup>, M. Assafin<sup>1,\*</sup>, L. Beauvalet<sup>2</sup>, R. Vieira-Martins<sup>1,2,3,\*\*</sup>, J. I. B. Camargo<sup>2,3</sup>, B. E. Morgado<sup>1,2</sup>

<sup>1</sup> Observatório do Valongo/UFRJ, Ladeira Pedro Antônio 43, CEP 20.080-090 Rio de Janeiro - RJ, Brazil  
e-mail: altair08@astro.ufrj.br

<sup>2</sup> Observatório Nacional/MCT, R. General José Cristino 77, CEP 20921-400 Rio de Janeiro - RJ, Brazil  
e-mail: rvm@on.br

<sup>3</sup> Laboratório Interinstitucional de e-Astronomia - LIneA, Rua Gal. José Cristino 77, Rio de Janeiro, RJ 20921-400, Brazil

Received ; accepted

## ABSTRACT

**Context.** Due to their orbital configurations, it is common belief that irregular satellites are probably captured by the giant planets in the early solar system. It is important to know their physical parameters, such as size, albedo and composition, to trace back their true origin. The best ground-based technique to constrain those parameters is the observation of stellar occultations by these objects.

**Aims.** We aim to predict stellar occultations for the eight biggest irregular satellites of Jupiter: Himalia, Elara, Pasiphae, Carme, Lysithea, Sinope, Ananke and Leda; the irregular satellite of Saturn: Phoebe; and the satellites of Neptune: Nereid and Triton.

**Methods.** We identified candidates to stellar occultations by the irregular satellites from the UCAC4 catalogue and from a catalogue of stars in the sky-path of Neptune obtained from observations made with the ESO2p2/WFI (2.2 m Max-Planck ESO telescope with the Wide Field Imager) instrument. These catalogues were crossed with the ephemeris of the satellites to identify stellar occultations. We used a new ephemeris based solely on the observations from Gomes-Júnior et al. (2015) to generate quick predictions for the short-time future for the satellites of Jupiter. For Phoebe, Nereid and Triton we used the JPL ephemeris.

**Results.** We managed to identify 687 candidates of stellar occultations between the period of January, 2015 and December, 2017. We made observational tests for the predictions of the events of Himalia in March 03, 2015 and Elara in March 30, 2015. The stars and objects were observed close to the date predicted and in the same field to minimize errors and the obtained relative satellite-star positions were used to evaluate the predictions.

**Conclusions.** The comparisons between the predictions and the observation tests show a good agreement. We discuss how the successful observation of a stellar occultation by these objects is quite possible and present some of those potential occultations.

**Key words.** Occultations - Planets and satellites: general - Planets and satellites: individual: Jovian and Saturnian irregular satellites

## 1. Introduction

Irregular satellites revolve around giant planets at large distances, on eccentric, highly inclined and frequently retrograde orbits. Because of these peculiar orbits, it is largely accepted that these objects did not form by accretion around of their planet, but were captured by their planets in the early solar system (Sheppard 2005).

There are several of capture mechanisms of objects by giant planets suggested in the literature. For example the Gas Drag in the primordial circumplanetary nebulae (Sheppard 2005) where the object would be affected by the gas drag and its velocity slowed down until it be captured by the planet. Another mechanism is called pull-down capture (Sheppard 2005), where the mass of the planet would increase while the object was temporarily captured.

A mechanism based in the Nice model (Morbidelli et al. 2005; Tsiganis et al. 2005; Gomes et al. 2005) was proposed by Nesvorný et al. (2007) and, in the specific case of Jupiter with the modern Nice model, by Nesvorný et al. 2014. In this

scenario, during the early solar system instability, encounters between the outer planets occurred. These planetary encounters could exchange energy and angular momentum between planets and the objects nearby allowing the capture of irregular bodies by the giant planets. In this scenario, the survival rate of prior-LHB (Late Heavy Bombardment) satellites is very small.

Another possible mechanism is the capture through collisional interactions (Sheppard 2005). A collision between two small bodies in the Hill's sphere of the planet could generate fragmented objects and the dissipated energy could be such that some of these objects could be captured.

Some of these objects are in dynamical groups with similar semi-major axis, eccentricities and inclinations, called families, similar to families found in the Main Asteroid Belt. These families may have been created by a parent body disrupted by collisions with comets or other satellites (Nesvorný et al. 2004). These collisions are more likely to have occurred during the LHB (Gomes et al. 2005).

Nesvorný et al. (2003) studied the collision rates between irregular satellites and concluded that some satellites could have been removed by collision with a bigger satellite. The collision rate between satellites of the Himalia Group (Himalia, Elara, Lysithea and Leda, mainly), for instance, was found to be more

Send offprint requests to: A. R. Gomes-Júnior

\* Affiliated researcher at Observatoire de Paris/IMCCE, 77 Avenue Denfert Rochereau 75014 Paris, France

\*\* Affiliated researcher at Observatoire de Paris/IMCCE, 77 Avenue Denfert Rochereau 75014 Paris, France

**Table 1.** Estimated diameter of the satellites and correspondent apparent diameter

Satellite	Diameter of the satellites		References
	mas <sup>a</sup>	km	
Ananke	8	29	1
Carme	13	46	1
Elara	24	86	1
Himalia	41	$(150 \times 120) \pm 20^b$	2
Leda	5	20	1
Lysithea	10	36	1
Pasiphae	17	62	1
Sinope	10	37	1
Phoebe	32	$212 \pm 1.4^b$	3
Nereid	15	$340 \pm 50^c$	4
Triton	124	$2707 \pm 2.0^c$	5

**References.** (1) Rettig et al. (2001); (2) Porco et al. (2003); (3) Thomas (2010); (4) Thomas et al. (1991); (5) Thomas (2000).

<sup>(a)</sup> Using a mean distance from Jupiter of 5 AU, from Saturn of 9 AU and from Neptune of 30 AU. <sup>(b)</sup> From Cassini observation.

<sup>(c)</sup> From Voyager-2 observation.

than one during the solar system age suggesting that their current structure was originated by satellite-satellite collision.

For Phoebe, ejected material from its surface caused by impacts could evolve due to Poynting-Robertson drag and collide with Iapetus causing the large variation in albedo observed on it (Nesvorný et al. 2003). Indeed, Cassini detected in Phoebe an absorption feature at  $2.42 \mu\text{m}$  (probably CN combinations) that was also detected in the dark side of Iapetus (Clark et al. 2005).

The region of origin of these object is not well known. Grav et al. (2003) and Grav & Bauer (2007) showed that the irregular satellites from the giant planets have their colors and spectral slopes similar to C-, D- and P-type asteroids, Centaurs and trans-neptunian objects (TNOs) suggesting that they come from different locations in the early solar system.

In order to obtain precise fundamental physical parameters like size, shape and albedo for the irregular satellites and thus to contribute to the study of their origin, we want to observe stellar occultations, which provide more accurate results than other ground-based techniques (Sicardy et al. 2011; Ortiz et al. 2012; Braga-Ribas et al. 2014).

No observation of a stellar occultation by an irregular satellite was published up to date. Since their estimated sizes are very small (see Table 2), this may have prevented earlier tries. But, in fact, given their distances to us, current ephemeris and star positions are already good enough for the prediction of the exact location and instant where the shadow of the occultation will cross the Earth. For instance, Himalia, supposedly the largest irregular satellite of Jupiter has an estimated size of 150 km (Porco et al. 2003), which is equivalent to an apparent size of about 40 mas (milliarcseconds). Thus, in this case, the cumulated uncertainty of ephemeris and star position must be around 40 mas to get good chances of observing a stellar occultation, which is quite feasible today.

From what we have just presented, we can think that the observation of an stellar occultation by an irregular satellite should be likelier than occultations by TNOs. The orbits of their host planets are well known and their observation time-span covers many orbital periods, contrary to TNOs. Moreover, the irregular satellites are closer to Earth which means a minor localization error in kilometer.

Gomes-Júnior et al. (2015) obtained 6523 suitable positions for 18 irregular satellites between 1992 and 2014 with an estimated error in the positions of about 60 to 80 mas. For some satellites the number of positions obtained is comparable to the number used in the numerical integration of orbits by the JPL (Jacobson et al. 2012). They pointed out that the ephemeris of the irregular satellites have systematic errors that may reach 200 mas for some satellites. For an object at the distance of Jupiter, this represents an error bigger than 700 km in the shadow path. Using the positions obtained by Gomes-Júnior et al. (2015) we produced a specific ephemeris for short-time future for the satellites of Jupiter and better predict stellar occultations for these objects.

We present in this paper stellar occultation predictions for the 8 major irregular satellites of Jupiter (Himalia, Elara, Pasiphae, Lysithea, Carme, Ananke, Sinope and Leda), Phoebe from Saturn and Nereid and Triton from Neptune. In the section 2 we explore the scientific rationale for the study of the irregular satellites and the possibility of having a common origin with TNOs. In section 3 we show the the process of the production of the new ephemeris. In section 4, we present the predictions of the stellar occultations by irregular satellites and how they were made. Some tests made to confirm the predictions are presented in section 5 and the final discussion are in section 6.

## 2. Scientific Rationale

There is no consensus for a single model explaining where the irregular satellites were formed. Ćuk & Burns (2004) showed that the progenitor of the Himalia group may have originated in heliocentric orbits similar to the Hilda asteroid group. Sheppard (2005) stated that the irregular satellites may be some of the objects that were formed within the giant planets region.

Phoebe is the most studied irregular satellite. Clark et al. (2005) suggest that its surface is probably covered by material of cometary origin. It was also stated by Johnson & Lunine (2005) that if the porosity of Phoebe is 15%, Phoebe would have an uncompressed density similar to those of Pluto and Triton.

Sheppard (2005) and Jewitt & Haghighipour (2007) also expose the possibility for the irregular satellites having their origin as comets or TNOs. TNOs are highly interesting objects that, due to their large heliocentric distances, may be highly preserved having their physical properties similar to those they had when they were formed (Camargo et al. 2014). This is even more true for the smaller objects, since in principle larger sizes favour physical differentiation processes in the body and vice-versa. However, due to the distance, the smaller TNOs from this region are more difficult to observe. A clever way to overcome this difficulty is to study the much closer irregular satellites, under the hypothesis that they share a common origin with the small TNOs' population.

Triton is a uncommon satellite. Its orbit is retrograde and inclined, but quasi-circular and very close to the planet compared to the irregular ones. Because Triton's orbit size is very small and its precession is not dominated by Solar perturbations, Triton is frequently excluded from the irregular satellites' class, but studied together by many authors (Sheppard 2005; Jewitt & Haghighipour 2007).

Similarly to the irregular satellites, Triton was probably captured in the early solar system and may have the same origin as the TNOs (Agnor & Hamilton 2006). Differently, Triton is bigger than the irregular satellites by an order of magnitude and has an atmosphere. The main motivation to study Triton by stellar occultations is to understand the evolution of its atmosphere due

to Triton's complicated and extreme seasonal cycle (McKinnon & Kirk 2007).

## Introduction V2

Irregular satellites revolve around giant planets at large distances, on eccentric, highly inclined and frequently retrograde orbits. Because of these peculiar orbits, it is largely accepted that these objects did not form by accretion around their planet, but were captured by their planets in the early solar system (Sheppard 2005).

There is no consensus for a single model explaining where the irregular satellites were formed. Ćuk & Burns (2004) showed that the progenitor of the Himalia group may have originated in heliocentric orbits similar to the Hilda asteroid group. Sheppard (2005) stated that the irregular satellites may be some of the objects that were formed within the giant planets region.

Grav et al. (2003) and Grav & Bauer (2007) showed that the irregular satellites from the giant planets have their colors and spectral slopes similar to C-, D- and P-type asteroids, Centaurs and trans-neptunian objects (TNOs). This suggests that they may have come from different locations in the early solar system.

Sheppard (2005) and Jewitt & Haghighipour (2007) also expose the possibility for the irregular satellites having their origin as comets or TNOs. TNOs are highly interesting objects that, due to their large heliocentric distances, may be highly preserved having their physical properties similar to those they had when they were formed (Camargo et al. 2014). This is even more true for the smaller objects, since in principle larger sizes favour physical differentiation processes in the body and vice versa. However, due to the distance, the smaller TNOs from this region are more difficult to observe. A clever way to overcome this difficulty is to study the much closer irregular satellites, under the hypothesis that they share a common origin with the small TNOs' population.

Phoebe is the most studied irregular satellite. Clark et al. (2005) suggest that its surface is probably covered by material of cometary origin. It was also stated by Johnson & Lunine (2005) that if the porosity of Phoebe is 15%, Phoebe would have an uncompressed density similar to those of Pluto and Triton.

In order to obtain precise fundamental physical parameters like size, shape and albedo for the irregular satellites and thus to contribute to the study of their origin, we want to observe stellar occultations, which provide more accurate results than other ground-based techniques (Sicardy et al. 2011; Ortiz et al. 2012; Braga-Ribas et al. 2014).

We present in this paper stellar occultation predictions for the 8 major irregular satellites of Jupiter (Himalia, Elara, Pasiphae, Lysithea, Carme, Ananke, Sinope and Leda), Phoebe from Saturn and Nereid and Triton from Neptune. Phoebe, being the most studied object with a good measured size, can be used to calibrate and evaluate the technique for similar objects.

Triton is a uncommon satellite. Its orbit is retrograde and inclined, but quasi-circular and very close to the planet compared to the irregular ones. Because Triton's orbit size is very small and its precession is not dominated by Solar perturbations, Triton is frequently excluded from the irregular satellites' class, but studied together by many authors (Sheppard 2005; Jewitt & Haghighipour 2007).

Similarly to the irregular satellites, Triton was probably captured in the early solar system and may have the same origin as the TNOs (Agnor & Hamilton 2006). Differently, Triton is bigger than the irregular satellites by an order of magnitude and has

**Table 2.** Estimated diameter of the satellites and correspondent apparent diameter

Satellite	Diameter of the satellites		References
	mas <sup>a</sup>	km	
Ananke	8	29	1
Carme	13	46	1
Elara	24	86	1
Himalia	41	$(150 \times 120) \pm 20^b$	2
Leda	5	20	1
Lysithea	10	36	1
Pasiphae	17	62	1
Sinope	10	37	1
Phoebe	32	$212 \pm 1.4^b$	3
Nereid	15	$340 \pm 50^c$	4
Triton	124	$2707 \pm 2.0^c$	5

**References.** (1) Rettig et al. (2001); (2) Porco et al. (2003); (3) Thomas (2010); (4) Thomas et al. (1991); (5) Thomas (2000).

(<sup>a</sup>) Using a mean distance from Jupiter of 5 AU, from Saturn of 9 AU and from Neptune of 30 AU. (<sup>b</sup>) From Cassini observation.

(<sup>c</sup>) From Voyager-2 observation.

an atmosphere. The main motivation to study Triton by stellar occultations is to understand the evolution of its atmosphere due to Triton's complicated and extreme seasonal cycle (McKinnon & Kirk 2007).

No observation of a stellar occultation by an irregular satellite was published up to date. Since their estimated sizes are very small (see Table 2), this may have prevented earlier tries. But, in fact, given their distances to us, current ephemeris and star positions are already good enough for the prediction of the exact location and instant where the shadow of the occultation will cross the Earth. For instance, Himalia, supposedly the largest irregular satellite of Jupiter has an estimated size of 150 km (Porco et al. 2003), which is equivalent to an apparent size of about 40 mas (milliarcseconds). Thus, in this case, the cumulated uncertainty of ephemeris and star position must be around 40 mas to get good chances of observing a stellar occultation, which is quite feasible today.

Gomes-Júnior et al. (2015) obtained 6523 suitable positions for 18 irregular satellites between 1992 and 2014 with an estimated error in the positions of about 60 to 80 mas. For some satellites the number of positions obtained is comparable to the number used in the numerical integration of orbits by the JPL (Jacobson et al. 2012). They pointed out that the ephemeris of the irregular satellites have systematic errors that may reach 200 mas for some satellites. In object at the distance of Jupiter, this represents an error bigger than 700 km in the shadow path. Using the positions obtained by Gomes-Júnior et al. (2015) we produced a specific ephemeris for short-time future for the satellites of Jupiter and better predict stellar occultations for these objects.

From what we have just presented, we can think that the observation of an stellar occultation by an irregular satellite should be likelier than occultations by TNOs. The orbits of their host planets are well known and their observation time-span covers many orbital periods, contrary to TNOs. Moreover, the irregular satellites are closer to Earth which means a minor localization error in kilometer.

In section 3 we show the the process of the production of the new ephemeris. In section 4, we present the predictions of the stellar occultations by irregular satellites and how they were



made. Some tests made to confirm the predictions are presented in section 5 and the final discussion are in section 6.

### 3. Special tailored Ephemerides (STE) for Jupiter irregular satellites

#### 3.1. Observations used

The last observations used to develop JPL current ephemeris of the irregular satellites of Jupiter were obtained in 2012 (Jacobson et al. 2012). As a result, the errors on the residuals of recent observations are too important for the ephemeris to accurately predict stellar occultations without any corrections. In order to have efficient ephemerides for predicting the stellar occultations by irregular satellites, we need predictions based on recent observations. This is why we decided to develop our own ephemeris based on the observations published in Gomes-Júnior et al. (2015).

Gomes-Júnior et al. (2015) published 6523 precise positions for 18 irregular satellites from observations made at the Observatório do Pico dos Dias (OPD), Observatoire Haute-Provence (OHP) and European Southern Observatory (ESO) between 1992 and 2015. Their offsets of the observations relative to the JPL ephemeris could be up to 200 mas for some satellites. At the distance of Jupiter, this represents an difference bigger than 700 km in the shadow path of an occultation.

#### 3.2. Description of the model

Our numerical model describes the dynamical evolution of the irregular satellites of Jupiter in a jovicentric reference frame. The satellites are submitted to the influence of the Sun and the rest of the solar system, as well as those of the Galilean satellites and the first harmonics of Jupiter's gravity field. The axis of the reference frame are those of the equatorial reference frame J2000.

We use the following notations:

- $i$  and  $l$  one of the irregular satellites of Jupiter
- $J$  Jupiter
- $j$  another body of the Solar System
- $M_j$  the mass of the  $j$  body, not an irregular satellite
- $m_i$  the mass of the irregular satellite  $i$
- $\mathbf{r}_i$  the position of the  $i$  body with respect to the barycentre of Jupiter System
- $r_{ij}$  the distance between bodies  $i$  and  $j$
- $R_J$  the radius of Jupiter
- $J_n$  the dynamic polar oblateness of the  $n$ th order for Jupiter's gravity field
- $U_{ij}$  potential generated by the oblateness of Jupiter on the satellite  $i$
- $\Phi_i$  is the inclination of the  $i$  satellite with respect to Jupiter's equator.

For an irregular satellite  $i$ , under the gravitational influence of Jupiter, the  $N' - 1$  other irregular satellites, the regular Jovian satellites and the rest of the Solar System ( $N$  bodies), the

equation of motion is:

$$\begin{aligned} \ddot{\mathbf{r}}_i = & -GM_J \frac{\mathbf{r}_J - \mathbf{r}_i}{r_{iJ}^3} - \sum_{l=1, l \neq i}^{N'} Gm_l \frac{\mathbf{r}_l - \mathbf{r}_i}{r_{il}^3} \\ & - \sum_{j=1}^N GM_j \left( \frac{\mathbf{r}_j - \mathbf{r}_i}{r_{ij}^3} - \underbrace{\frac{\mathbf{r}_j - \mathbf{r}_J}{r_{jJ}^3}}_{\text{undirectperturbations}} \right) \\ & + GM_J \nabla U_{ij} - \sum_{l=1}^N \underbrace{Gm_l \nabla U_{il}}_{\text{undirectperturbations}} \end{aligned} \quad (1)$$

where the oblateness potential seen by the body  $i$  because of Jupiter is:

$$\begin{aligned} U_{ij} = & -\frac{R_J^2 J_2}{r_{iJ}^3} \left( \frac{3}{2} \sin^2 \Phi_i - \frac{1}{2} \right) \\ & -\frac{R_J^4 J_4}{r_{iJ}^5} \left( \frac{35}{8} \sin^4 \Phi_i - \frac{15}{4} \sin^2 \Phi_i + \frac{3}{8} \right) \\ & -\frac{R_J^6 J_6}{r_{iJ}^7} \left( \frac{231}{16} \sin^6 \Phi_i - \frac{315}{16} \sin^4 \Phi_i + \frac{105}{16} \sin^2 \Phi_i - \frac{5}{16} \right) \end{aligned} \quad (2)$$

The expressions of  $\nabla_i U_{ij}$  and  $\nabla_i U_{il}$  have been developed in Lainey et al. (2004). The equations of motion are integrated with the numerical integrator RADAU (Everhart 1985). Our model was fitted to the observations through a least-squares procedure. The satellites were integrated one dynamical family at a time, to gain computing time, while losing minimum precision. Indeed, the interactions between satellites not belonging to the same dynamical family are negligible considering the short timespan of our integration.

The initial osculating elements at the origin of integration are presented in Table 3, while the residuals are presented in Fig 1 for Himalia and Carme as representatives.

All the orbits determined for the satellites show satisfying residuals. The residuals are lower than those obtained with JPL ephemeris, but the accuracy of an ephemeris decreases when we get further from the time of observations. The main risk of divergence over time comes from the possible absence of long-term effects when fitting to a short timespan of observations. If that were the case, our ephemeris would diverge too quickly to be of any use. JPL ephemerides are fitted over all the available observations. As a result, they will diverge less quickly than our own. Though they are no longer precise enough for our use, they remain a precious reference to identify whether our own model presents a quick divergence. We compared our ephemeris to the JPL for all the Jupiter satellites we fitted, until 2017. The divergence between 2015 and 2018 is at most 98 mas in  $\Delta \alpha \cos \delta$  and 58 mas in  $\Delta \delta$  for Himalia and 181 mas in  $\Delta \alpha \cos \delta$  and 152 mas in  $\Delta \delta$  for Carme. A example of this difference for satellites Himalia and Carme over time is presented in Figure 2.

The obtained ephemeris is hereafter referred as STE, for special tailored ephemeris.

### 4. Prediction of occultations

The prediction of the occultations was made by crossing the stellar coordinates and proper motions of the UCAC4 catalogue




Fig. 1. Residuals of the numerical integration for Himalia and Carme.

Table 3. Initial osculating elements for Jupiter irregular satellites at JD 2451545.0.

Satellite	a (km)	e	incl. (EQJ2000)	$\Omega$	$\omega$	$v$
Himalia	$11372100 \pm 500$	$0.166 \pm 0.002$	$45.14 \pm 0.15$	$39.77 \pm 0.19$	$351.48 \pm 0.46$	$97.35 \pm 0.48$
Elara	$11741170 \pm 690$	$0.222 \pm 0.002$	$28.64 \pm 0.18$	$68.42 \pm 0.43$	$179.82 \pm 0.56$	$339.08 \pm 0.82$
Lysithea	$11739900 \pm 1300$	$0.136 \pm 0.004$	$51.12 \pm 0.27$	$5.53 \pm 0.52$	$53.0 \pm 1.5$	$318.9 \pm 2.0$
Leda	$11140300 \pm 4300$	$0.173 \pm 0.007$	$16.15 \pm 0.75$	$272.6 \pm 1.7$	$212.2 \pm 3.6$	$218.8 \pm 3.2$
Pasiphae	$23425000 \pm 5000$	$0.379 \pm 0.001$	$152.44 \pm 0.10$	$284.59 \pm 0.21$	$135.96 \pm 0.19$	$236.97 \pm 0.16$
Sinope	$22968800 \pm 5200$	$0.316 \pm 0.002$	$157.76 \pm 0.12$	$256.62 \pm 0.55$	$298.38 \pm 0.55$	$167.57 \pm 0.19$
Carme	$24202924 \pm 4800$	$0.242 \pm 0.001$	$147.13 \pm 0.10$	$154.01 \pm 0.25$	$47.90 \pm 0.29$	$234.41 \pm 0.19$
Ananke	$21683800 \pm 7200$	$0.380 \pm 0.002$	$172.29 \pm 0.20$	$56.9 \pm 1.2$	$123.3 \pm 1.2$	$231.24 \pm 0.21$




Fig. 2. Differences between the JUP300 JPL ephemeris and the special-tailored ephemeris for Himalia and Carme.

(Zacharias et al. 2013) with the **special-tailored ephemeris** for Jupiter satellites (see Sec. 3) and JPL ephemeris for the others. The search for stellar candidates follows the same procedure as presented by Assafin et al. (2010, 2012) and Camargo et al. (2014).

We predicted occultation for the 8 major irregular satellites of Jupiter: Ananke, Carme, Elara, Himalia, Leda, Lysithea, Pasiphae and Sinope; Phoebe of Saturn and Triton and Nereid of Neptune.

For Triton and Nereid, the candidates for stellar occultations in 2015 and 2016 was searched using the WFI catalogue in the same way as the predictions for Centaurs and TNOs occultations by Assafin et al. (2010, 2012) and Camargo et al. (2014). This catalogue contains the stars in the path of Neptune in the sky

up to mid-2016. The catalogue was generated by observations made at the ESO 2p2 telescope (IAU code 809) using the Wide Field Imager (WFI) CCD mosaic detector. The filter used was the broad-band R filter ESO#844 with  $\lambda_c = 651.725$  nm and  $\Delta\lambda = 162.184$  nm.

A total of 687 events were identified between January 2015 and December 2017. In table 4 we present the number of stellar occultations predicted by year for each satellite. Table 5 shows a sample of the catalogue of occultations generated and their parameters, which are necessary to produce occultation maps (see Fig. 3 as an example). Since these objects are very small, the duration of each event is a few seconds.

The ESA astrometry satellite GAIA (de Bruijne 2012) catalogue is expected to be released in the end of 2016. The precise

**Table 5.** Stellar occultation sampled predictions for Pasiphae

d m Year	h m s	RA (ICRS)	Dec	C/A	P/A	$\nu$	$D$	$R^*$	$\lambda$	LST	$\mu_{\alpha^*}$	$\mu_{\delta}$
09 04 2016	03:58:19.	11 14 36.7707	+07 39 20.7610	1.003	17.9	-12.88	4.54	14.9	271.	22:03	12.	-33.
13 06 2016	00:16:12.	11 12 48.5020	+07 06 43.3520	0.661	30.0	+14.32	5.50	13.9	262.	17:45	-1.	1.
27 06 2016	13:56:09.	11 18 03.4160	+06 23 45.1940	1.707	28.0	+20.29	5.74	11.7	44.	16:53	4.	-10.
18 07 2016	15:07:24.	11 28 15.5076	+05 05 31.8060	0.942	26.7	+27.80	6.05	14.0	8.	15:40	4.	4.
22 07 2016	16:15:07.	11 30 30.4310	+04 48 43.4340	0.644	206.5	+29.04	6.11	14.6	348.	15:27	23.	-24.
24 07 2016	01:37:34.	11 31 17.8471	+04 42 49.0540	0.029	206.6	+29.46	6.12	15.1	206.	15:22	2.	-8.
24 07 2016	17:37:18.	11 31 40.7472	+04 39 57.5060	0.840	26.5	+29.46	6.13	14.9	326.	15:20	-11.	-1.

**Notes.** Entries included: day of the year and UTC time of the prediction; right ascension and declination of the occulted star - in the original table; C/A: the geocentric closest approach, in arcseconds; P/A: the planet position angle with respect to the occulted star at C/A, in degrees;  $\nu$ : velocity in plane of sky, in  $\text{km s}^{-1}$ : positive = prograde, negative = retrograde;  $D$ : planet range to Earth, in AU;  $R^*$ : normalized magnitude to a common shadow velocity of  $20 \text{ km s}^{-1}$  by the relationship  $R^* = R_{\text{actual}} + 2.5 \times \log_{10} \left( \frac{\text{velocity}}{20 \text{ km s}^{-1}} \right)$ ;  $\lambda$ : east longitude of subplanet point in degrees, positive towards east; LST: UT +  $\lambda$ : local solar time at subplanet point, hh:mm;  $\mu_{\alpha^*}$  and  $\mu_{\delta}$ : proper motions in right ascension and declination, respectively (mas/year).

**Table 4.** Number of stellar occultations for each satellite from 2015 up to 2017.

Satellite	2015	2016	2017	Total
Ananke	21	12	16	49
Carme	26	20	14	60
Elara	32	14	16	62
Himalia	26	15	12	53
Leda	13	8	24	45
Lysithea	22	16	11	49
Pasiphae	25	20	19	64
Sinope	19	15	21	55
Phoebé <sup>a</sup>	13 <sup>b</sup>	32 <sup>b</sup>	98 <sup>b</sup>	143 <sup>b</sup>
Nereid <sup>a</sup>	41 <sup>b</sup>	11 <sup>b</sup>	1 <sup>b</sup>	53 <sup>b</sup>
Triton <sup>a</sup>	37 <sup>b</sup>	16 <sup>b</sup>	1 <sup>b</sup>	54 <sup>b</sup>

Occultations predicted using the UCAC4 catalogue and STE ephemeris. <sup>(a)</sup> Using JPL ephemeris. <sup>(b)</sup> Using the WFI catalogue as explained in Sec. 4.

star positions to be derived by GAIA will provide better predictions with the main source of error being the ephemeris. Astrometric reduction of observations published in Gomes-Júnior et al. (2015) will be revised with the GAIA catalogue and the predictions from 2017 onwards will be better. Because of that, we only publish stellar occultations with UCAC4 until 2017.

## 5. Occultation tests

Observing a stellar occultation demands a great effort. And, in our case, the shadow covers a very restricted area on Earth because of the size of the irregular satellites. So, before we start a large observational campaign, we tested some occultation predictions for large targets, to assess the quality of the predictions.

The tests consisted in observing the object and star to be occulted near the date of the event predicted when the two objects were present in the same field of view (FOV), preferably when the objects were close to each other. Thus, the relative positions between the two objects had minimal influence of the errors of the reference catalogue of stars used and possible field distortions (Peng et al. 2008, and references therein). The relative positions of the star and of the satellites were used to check the original predictions.

Two occultation tests were performed, one by Himalia that occurred on March 3, 2015 and the second by Elara that occurred


on March 30, 2015. For each event, four situations were considered:

1. Prediction with the STE ephemeris (see Sec. 3), and the nominal UCAC4 position of the star;
2. Prediction with the JPL ephemeris and the nominal UCAC4 position of the star;
3. From star and satellite offsets calculated from observations made a few days before the occultation when the objects were separated (different FOVs);
4. Same as 3 but with the star and the satellite close in the same FOV.

Figure 4 shows four zoomed in occultation maps corresponding to each of the four situations for Himalia. The map 4a is for the situation 1 in the list above. Map 4b using JPL ephemeris (situation 2). The map 4c was made from obtained positions on February 22 observed with the Zeiss telescope (diameter = 0.6m; FOV = 12'6; pixel scale = 0'37/px) at the Observatório do Pico dos Dias (OPD, IAU code 874, 45°34'57''W, 22°32'04''S, 1864m). On that day, Himalia and the star were observed in separate FOVs as they were still far apart (situation 3). On the night of the event, March 3, the objects were observed with Perkin-Elmer telescope (diameter = 1.6m; FOV = 5'8; pixel scale = 0'17/px) at OPD just over an hour after the time scheduled for the event. Satellite and star were separated by about 16 arcsec, so very close to each other (situation 4). From the calculated offsets, the map 4d was generated. Notice that the shadow path was not predicted to cross the OPD (green point at the bottom of the maps). This was not necessary for testing the prediction.

In this event, it is possible to see that the shade does not vary by much among the four maps suggesting that there was a good probability of observing the event. In fact, the largest differences between the shadows of the four maps were 36s, which correspond to 120 mas, in time along the shadow path and 101km (31 mas) in the direction perpendicular to the shadows (see table 6).

The second test was with the satellite Elara, which is the second largest irregular satellite of Jupiter. The event was predicted to occur at March 30, 2015. The observations were made on March 25 and April 2, 2015 with the Zeiss telescope. On the night of April 2 the star and satellite could still be observed in the same FOV. Due to Elara being fainter than Himalia by 2 magnitudes, dispersions of the satellite positions on both nights ended up being higher than for Himalia. The largest differences between them were 144s (186 mas) in time along the shadow path and 525 km (153 mas) perpendicular to it (see table 7).



**Fig. 3.** Occultation map for Pasiphae regarding to the first event sampled in Table 5

**Table 6.** Comparison between the predictions of the Himalia occultation at March 03, 2015.

Method	Difference from STE prediction		
	Instant of C/A	C/A	Map
STE	00:39:51 UTC	0 <sup>h</sup> 703	4a
JPL	-26s	+11mas (36km)	4b
Feb. 22 Obs.	-14s	-20mas (65km)	4c
Mar. 03 Obs.	-36s	-09mas (29km)	4d

**Table 7.** Comparison between the predictions of Elara occultation at March 30, 2015.


Method	Difference from STE prediction	
	Instant of C/A	C/A
STE	01:45:13 UTC	1 <sup>h</sup> 082
JPL	+02s	+57mas(195km)
Mar. 25 Obs.	-70s	+20mas (68km)
Apr. 02 Obs.	+74s	-96mas (330km)

## 6. Discussion

We presented stellar occultations for the period of 2015-2017 for eight irregular satellites of Jupiter: Ananke, Carme, Elara, Himalia, Leda, Lysithea, Pasiphae, and Sinope; one satellite of Saturn: Phoebe; and two satellites of Neptune: Triton and Nereid. The procedure used was the same as that for the prediction of stellar occultations by Pluto and its satellites in Assafin et al. (2010) and by Centaurs and TNOs in Assafin et al. (2012) and Camargo et al. (2014).

The candidate stars **was** searched in the UCAC4 catalogue, except for the candidates in 2015 and 2016 for Triton and Nereid. In this case, we used the WFI catalogue that was generated from observations made with ESO2p2/WFI CCD mosaic that covered the path of Neptune in the sky-**up** to 2016 (see Sec. 4). From this, a total of 687 events **were** foreseen.

The probability of successfully observing an occultation is roughly the ratio of the satellite's diameter by the budget error of **ep** and star position. Thus, UCAC4 errors ranging between 20 mas - 50 mas combined with a mean accuracy in the **position** of 30 mas for Himalia and 150 mas for Leda published



**Fig. 4.** Predictions for Himalia: The straight lines show the size of the shadow. The big red dot show the geocentric closest approach of the shadow. The black "+" marks at maps (b,c,d) are the STE prediction closest approach for reference. The small red ones are the center of the shadow separated by one minute. 4a is the map using the STE ephemeris showed in Sec. 3. 4b shows the shadow using the JPL ephemeris. In 4c we apply offsets to the positions of star and satellite accordingly the observations made at February 22. 4d is as in 4c but with observations made at March 03 when the objects were close to each other. The green dot at the bottom of the maps is the location of Observatório do Pico dos Dias.

in Table 2 of Jacobson et al. (2012) would give 80%-50% probability of observing such an event by Himalia and  $\approx 3\%$  for Leda, the smallest irregular satellite in the sample. Observations a few days before the date of occultation predicted may improve the combined accuracy to 40-80 mas, depending on the magnitude of the objects.

The tests made with an occultation expected to happen in March 03, 2015 for Himalia showed that this event would probably have been observed successfully in case there were observers available in the shadow area. A similar test was made for an occultation of Elara in March 30, 2015. The results show satisfying small offsets. ~~We can expect similar results for other satellites and our method should allow successful stellar occultation.~~

Continuous observations of the satellites and candidates are recommended and fitting of our dynamical model to those ob-

servations are expected to be the respective position errors. The GAIA catalogue, which is to be released at the end of 2016 will improve the position error of the stars and will allow for the detection of occultations candidates of fainter stars, which are not present in the UCAC4 catalogue. The release of the GAIA catalogue should have a positive impact on both the astrometric precision of occulted stars and the reduction of astrometric positions of occulting satellites. As a result, stellar occultations by irregular satellites should increase in number as well as success.

*Acknowledgements.* ARG-J thanks the financial support of CAPES. MA thanks the CNPq (Grants 473002/2013-2 and 308721/2011-0) and FAPERJ (Grant E-26/111.488/2013). RV-M thanks grants: CNPq-306885/2013, Capes/Cofecub-2506/2015, Faperj/PAPDRJ-45/2013. JIBC acknowledges CNPq for a PQ2 fellowship (process number 308489/2013-6). FB-R acknowledges PAPDRJ-FAPERJ/CAPES E-43/2013 number 144997, E-26/101.375/2014. BEM thanks the financial support of CAPES.



## References

- Agnor, C. B. & Hamilton, D. P. 2006, *Nature*, 441, 192–194
- Assafin, M., Camargo, J. I. B., Vieira Martins, R., et al. 2010, *Astronomy & Astrophysics*, 515, A32
- Assafin, M., Camargo, J. I. B., Vieira Martins, R., et al. 2012, *Astronomy & Astrophysics*, 541, A142
- Braga-Ribas, F., Sicardy, B., Ortiz, J. L., et al. 2014, *Nature*, 508, 72–75
- Camargo, J. I. B., Vieira-Martins, R., Assafin, M., et al. 2014, *Astronomy & Astrophysics*, 561, A37
- Clark, R. N., Brown, R. H., Jaumann, R., et al. 2005, *Nature*, 435, 66–69
- de Bruijne, J. H. J. 2012, *Astrophysics and Space Science*, 341, 31–41
- Everhart, E. 1985, in *Dynamics of Comets: Their Origin and Evolution*, Proceedings of IAU Colloq. 83, held in Rome, Italy, June 11–15, 1984. Edited by Andrea Carusi and Giovanni B. Valsecchi. Dordrecht: Reidel, *Astrophysics and Space Science Library*. Volume 115, 1985, p.185, ed. A. Carusi & G. B. Valsecchi, 185
- Gomes, R., Levison, H. F., Tsiganis, K., & Morbidelli, A. 2005, *Nature*, 435, 466–469
- Gomes-Júnior, A. R., Assafin, M., Vieira-Martins, R., et al. 2015, *Astronomy & Astrophysics*
- Grav, T. & Bauer, J. 2007, *Icarus*, 191, 267–285
- Grav, T., Holman, M. J., Gladman, B. J., & Aksnes, K. 2003, *Icarus*, 166, 33–45
- Jacobson, R., Brozović, M., Gladman, B., et al. 2012, *The Astronomical Journal*, 144, 132
- Jewitt, D. & Haghighipour, N. 2007, *Annual Review of Astronomy and Astrophysics*, 45, 261–295
- Johnson, T. V. & Lunine, J. I. 2005, *Nature*, 435, 69–71
- Lainey, V., Duriez, L., & Vienne, A. 2004, *Astronomy and Astrophysics*, 420, 1171–1183
- McKinnon, W. & Kirk, R. 2007, *Encyclopedia of the Solar System*, 483–502
- Morbidelli, A., Levison, H. F., Tsiganis, K., & Gomes, R. 2005, *Nature*, 435, 462–465
- Nesvorný, D., Alvarillos, J. L. A., Dones, L., & Levison, H. F. 2003, *AJ*, 126, 398–429
- Nesvorný, D., Beaugé, C., & Dones, L. 2004, *AJ*, 127, 1768–1783
- Nesvorný, D., Vokrouhlický, D., & Deienno, R. 2014, *ApJ*, 784, 22
- Nesvorný, D., Vokrouhlický, D., & Morbidelli, A. 2007, *AJ*, 133, 1962–1976
- Ortiz, J. L., Sicardy, B., Braga-Ribas, F., et al. 2012, *Nature*, 491, 566–569
- Peng, Q., Vienne, A., Lainey, V., & Noyelles, B. 2008, *Planetary and Space Science*, 56, 1807–1811
- Porco, C. C., West, R. A., McEwen, A., et al. 2003, *Science*, 299, 1541–1547
- Rettig, T., Walsh, K., & Consolmagno, G. 2001, *Icarus*, 154, 313–320
- Sheppard, S. S. 2005in (Cambridge University Press (CUP)), 319
- Sicardy, B., Ortiz, J. L., Assafin, M., et al. 2011, *Nature*, 478, 493–496
- Thomas, P. 2000, *Icarus*, 148, 587–588
- Thomas, P. 2010, *Icarus*, 208, 395–401
- Thomas, P., Veverka, J., & Helfenstein, P. 1991, *J. Geophys. Res.*, 96, 19253
- Tsiganis, K., Gomes, R., Morbidelli, A., & Levison, H. F. 2005, *Nature*, 435, 459–461
- Zacharias, N., Finch, C. T., Girard, T. M., et al. 2013, *The Astronomical Journal*, 145, 44
- Čuk, M. & Burns, J. A. 2004, *Icarus*, 167, 369–381